

Original Research

# Effects of Lime Application on Activities of Related Enzymes and Protein Expression of Saponin Metabolism of *Panax notoginseng* under Cadmium Stress

Zuran Li<sup>1</sup>, Yueqin Jiang<sup>2</sup>, Yanqun Zu<sup>2\*</sup>, Xinyue Mei<sup>2\*\*</sup>, Li Qin<sup>2</sup>, Bo Li<sup>2</sup>

<sup>1</sup>College of Horticulture and Landscape, Yunnan Agriculture University, P.R. China

<sup>2</sup>College of Resources and Environment, Yunnan Agriculture University, P.R. China

Received: 7 February 2020

Accepted: 14 April 2020

## Abstract

*Panax notoginseng* was used as material with exogenous Cd (0, 0.6, 3.0, 6.0, 9.0, 12 mg kg<sup>-1</sup>, prepared with CdCl<sub>2</sub>) and lime (0, 0.75, 1.5 t ha<sup>-2</sup>, prepared with Ca(OH)<sub>2</sub>) application to study the effect of cadmium stress on growth and saponins contents of *P. notoginseng*, the relative enzymes activities of saponins metabolism (SS, mevalonate kinase(MVK), P450,  $\beta$ -amylin synthase ( $\beta$ -AS)) and protein expression used with iTRAQ proteome analysis technique to understand the response mechanism of *P. notoginseng*. The results showed the contents of soil available Cd and Cd contents in main root decreased under lime application. The available contents of soil Cd decreased by 9.2%-38.1% with 1.5 t ha<sup>-2</sup> lime application under Cd treatments. The biomass, root length, root surface area and root volume increased with lime application under Cd stress. The activities and expression levels of MVK, P450 and  $\beta$ -amylin synthase decreased, resulting in decrease 8.6%-23.6% in saponins contents and ratio of Rg1 to Rb1. In general, the results indicate excessive application of lime would lead to a certain reduction in the content of saponin under Cd stress and inhibition of conversion of diol-saponins to triol-saponins with single-peak model to Ca/Cd ratio. It could be recommended that the ratio of Ca to Cd is less than 680 for reduction of cadmium content and maintaining saponin contents of main root of *P. notoginseng* for the production of high quality.

**Keywords:** *Panax notoginseng*, lime, metabolism, heavy metal, iTRAQ proteome analysis

## Introduction

The soil Cd pollution has become one of the most important environmental problems [1]. Due to a series

of activities of humans, including irrigation of farmland sewage and industrial wastes, the contents of heavy metals in the soil increased [2-4]. Cadmium in soil could be absorbed by plants, which affects plant growth and physiological metabolism and reduces the yield and quality of crops [5,6]. Plants reduce the toxic effects of heavy metals by altering their own structure

\*e-mail: 649332092@qq.com

\*\*e-mail: 339257631@qq.com

or secondary metabolic pathways [7,8]. Protein change mechanism and protein-protein interactions could be studied through proteome technology. Isobaric tags for relative and absolute quantitation (iTRAQ) are more commonly used proteomics techniques in recent years. The differences in proteome expression of plants in response to Cd stress have been studied in rice (*Oryza sativa*), *Populus tremula*, *Kandelia candel*, *Thlaspi caerulescens*, *Phytolacca americana* and *Sedun alfredii* and *Perilla frutescens* (L.) Britt. [3,9], in which ATP activity-related regulatory proteins, detoxification and anti-oxidation and signaling-related proteins, ion transporters, sulfur and glutathione-related metabolic proteins showed differences in expression to cadmium stress [9-12].

*Panax notoginseng* (Burk.) F.H.Chen is a perennial herb of the *Panax* genus of Araliaceae family. Saponin is the main medicinal ingredient of *P. notoginseng*, and belongs to the dammarane tetracyclic triterpene substance, which the functions of saponin are related to promote blood circulation, immunity and anti-tumor. The saponin includes 20(S)-protopanaxadiol and 20(S)-protopanaxatriol [13]. There are more than 70 kinds of monomeric saponins including notoginsenoside R1 and ginsenosides Rb1 and Rg1, in which notoginsenoside R1 is unique to *P. notoginseng*, and contents of ginsenoside Rg1 and Rb1 are the highest [14-16]. The synthesis process of saponins in plants includes: (1) synthesis of isopentenyl pyrophosphate and dimethylallyl pyrophosphate; (2) catalytic synthesis of isopentenyl transferase and anthracycline cyclase 2,3-oxidation squalene; (3) cyclized, hydroxylated, glycosylated modified to form dammarane type notoginsenoside. Key enzymes include squalene synthase (SS), farnesyl diphosphate synthase (FPS), squalene epoxidase (SE), and dammarenediol synthase (DS), cytochrome P450, glycosyl transferase (GT), mevalonate kinase (MVK) and  $\beta$ -amylin synthase ( $\beta$ -AS) [13]. The origin and main production area of *P. notoginseng* was located in Wenshan prefecture, Yunnan Province, China. The yield of *P. notoginseng* in Wenshan prefecture accounts for more than 90% of total yield in China. The background values of heavy metals in soil of *P. notoginseng* production area are high, which threatens the quality of *P. notoginseng* [17]. Zu et al. (2017) investigated soil heavy metals contents in Wenshan prefecture, indicating the over-standard rate of soil Cd content was 53.3% [18]. Control measurements should be taken to reduce Cd contents in soil and *P. notoginseng*. Environmental stress can change the synthesis of saponin and the expression of relative genes [19]. Other studies showed that contents of notoginsenoside R1 reached a maximum when the soil Cd content was 6 mg kg<sup>-1</sup>, and the contents of each monomer saponin and total saponin was significantly decreased when Cd was 30 mg kg<sup>-1</sup> [20]. Reducing the Cd content and maintaining the contents of saponins are important for high quality of *P. notoginseng*.

Application of lime can increase soil pH and reduce the bioavailability of heavy metals and the absorption and accumulation of heavy metals by plants [21,22]. Calcium is an essential element of plant growth. The Ca<sup>2+</sup> can maintain cell structure stability and cell viability, regulate photosynthesis, signal transduction, transpiration and respiration [19, 23-25]. Under cadmium stress, the supply of calcium can alleviate the degree of chlorophyll decline [26]. The vacuole is the main storage place of the Ca<sup>2+</sup>. The calcium ion in the vacuole mainly depends on the ions Ca<sup>2+</sup> channel to participate in the formation of calcium signal. The Ca<sup>2+</sup> in the chloroplast and mitochondria is related to the cell senescence and death [27], the transport of heavy metal ions [28], and abiotic stress [29]. The valence of Cd<sup>2+</sup> and Ca<sup>2+</sup> is the same and the ionic radius is close. The ions Ca<sup>2+</sup> and Cd<sup>2+</sup> compete for the adsorption site of the soil. Application of lime significantly reduced the Cd contents of corn [9,21] and rice [30], and increased the biomass of *Arabidopsis* seedlings [31].

Therefore, the hypothesis include: (1) application of lime under cadmium stress reduces Cd content in *P. notoginseng*, changes the growth and morphology of *P. notoginseng* roots; (2) Under certain Ca/Cd ratio, key enzyme activities and genes expressions related to saponin metabolism change, affecting the monomer heterogeneity and total saponins contents. The iTRAQ proteomics technology was used to study the response mechanism of lime-mediated root growth and saponin metabolism under cadmium stress.

## Materials and Methods

### Treatments and Sampling

One -year old *P. notoginseng* seedling was cultivated in upland red soil. Background values of experiment soil were pH 5.42, CEC 19.4 cmol kg<sup>-1</sup>, alkali nitrogen 124.61 mg kg<sup>-1</sup>, available phosphorus 28.77 mg kg<sup>-1</sup>, available potassium 318.92 mg kg<sup>-1</sup>, total nitrogen 0.18 %, total phosphorus 0.10%, total potassium 1.21% and Cd content 0.53 mg kg<sup>-1</sup>. The experiment site was located at 25°31'21.80"N, 103°17'12.87"E and altitude of 1871 m.

Three lime application levels were 0 (L<sub>0</sub>), 0.75 (L<sub>1</sub>) and 1.5 (L<sub>II</sub>) t ha<sup>-2</sup>, formulated with Ca(OH)<sub>2</sub>. The 0-15 cm layer soil of each plot was mixed with differently treated lime. Six Cd treatment levels were 0 (Cd<sub>0</sub>), 0.6 (Cd<sub>0.6</sub>), 3.0 (Cd<sub>3</sub>), 6.0 (Cd<sub>6</sub>), 9.0 (Cd<sub>9</sub>), 12 mg kg<sup>-1</sup> (Cd<sub>12</sub>), formulated with CdCl<sub>2</sub>·2.5H<sub>2</sub>O. After CdCl<sub>2</sub>·2.5H<sub>2</sub>O mixed with 2 mm soil based on the ratio of 1:40 of CdCl<sub>2</sub>·2.5H<sub>2</sub>O : soil, it was evenly spread in each plot, and then mixed with the soil of 0-15 cm. Each treatment was 3 replicates. After 15 days, one -year old *P. notoginseng* seedling were transplanted in February 2016. The experiment plots was randomly arranged. The area of the plot was 1.8 m<sup>2</sup> (1.2 m × 1.5 m). 96 seedlings were transplanted in each

plot, with a plant spacing of 12 cm × 15 cm. General management was carried through growth periods.

During the vigorous flowering stage of *P. notoginseng* (August 23, 2016), 15 plants of *P. notoginseng* were collected in each plot. The ten plants were used for the determination of Cd content, root morphology and saponin content. The soil and plants were sampled separately. Four treatments including  $L_0Cd_0$  (0 t ha<sup>-2</sup> lime + 0 mg kg<sup>-1</sup> Cd),  $L_0Cd_6$  (0 t ha<sup>-2</sup> lime + 6.0 mg kg<sup>-1</sup> Cd),  $L_LCd_0$  (0.75 t ha<sup>-2</sup> lime + 0 mg kg<sup>-1</sup> Cd) and  $L_LCd_6$  (0.75 t ha<sup>-2</sup> lime + 6.0 mg kg<sup>-1</sup> Cd) with 3 replicates were selected for proteome analysis of the main root of *P. notoginseng*. The five plants of each treatment used for the enzyme activity and proteomic analysis were washed with ultrapure water and stored in a cryopreservation tube with the liquid nitrogen quickly frozen at -80°C.

The soil was air dried and passed through a 1 mm sieve for the determination of soil pH and available Cd content. The plants were first washed with tap water, then washed with deionized water. The roots were scanned by the root scanning analyzer (the Epson Expression 10000XL1.0 root system scanner (Japan)), using the root analysis software WinRHIZO Pro V2007d. (Regina, Canada) to analyze the root morphological characteristics (total root length, root surface area and root volume) of *P. notoginseng*. After the root scanned, the each part of *P. notoginseng* was placed in a blast drying oven, dried at 105°C for 30 min, then dried at 65°C to a constant weight. After cooling, the biomass of each part of *P. notoginseng* was measured. After grinding through a 1 mm sieve, it was used to determine the contents of Cd and saponin of *P. notoginseng*.

#### Cadmium (Cd) Content in Root of *P. notoginseng*

Some 0.2 g of the main root of *P. notoginseng* was accurately weighed, and placed in a teflon tank and added 2 mL nitric acid and 1 mL hydrogen peroxide, then tighten with a stainless steel cover, and place in an oven at 160°C for 4 h. After cooling to room temperature, the digestive the mixture was moved to a 10 mL volumetric flask. The Cd content of solution was measured by ICP-MS (Nexion 350X, Perkin Elmer, USA).

#### Saponin Contents in Root of *P. notoginseng*

Some 0.6 g main root powder was accurately weighed, and dissolved in 50 mL methanol and incubation 12 hr and 1 hr in an 80°C water bath. After cooling to room temperature, the mixture was moved to a 50 mL volumetric flask with adding methanol and filtered for analysis contents of saponin Rb1, saponin Rg1 and notosaponin R1, 10 µL using HPLC (Prominence LC-20A, Japan) and the acetonitrile mobile phase A and aqueous mobile phase B, chromatographic analysis was performed on an octadecylsilyl column

under the detection wavelength 203 nm [16]. Standard reagents of 0.4 mg mL<sup>-1</sup> saponin Rg1, saponin Rb1 and 0.1 mg mL<sup>-1</sup> notosaponin R1 were used after standard reagents dissolved in 1 mL methanol (The Ministry of Health of Pharmacopoeia Committee 2000). Content of saponin in main root was expressed in unit % of dry weight.

Samples for HPLC analysis was not necessary to purify according to standard analysis method in China. The parameter of column to HPLC was C18 with particle diameter 5 µm and length 200 mm. The detector was SPD. The reagents to HPLC were chromatographic pure from J&K Scientific LTD, Beijing, China.

The total saponin content of three monomers (*Panax notoginseng* saponins, PNS) = Rb1 + Rg1 + R1.

Saponin yield of main root = mean of main root biomass × PNS.

#### Extraction and Measurement of Enzymes Activities

Some 0.500 g of the main root powder was weighed, and put into the frozen mortar. 5 mL of phosphate buffer was added to grind homogenate, then poured into the centrifuge tube for refrigerated centrifugation. The liquid supernatant was collected. The standard solution was diluted 5 times with a 2-fold dilution gradient in the kit as required. Activities of squalene synthase (SS), mevalonate kinase (MVK), P450 reductase and β-amylin synthase (β-AS): Enzymatic assay was carried out using relative assay kits following manufactory's instruction (Tszelisa, USA) on a ELISA Reader (DNM-9602, Beijing Perlong New technology Limited Company). Enzymatic activity was expressed in units per gram of fresh weight (U g<sup>-1</sup>).

#### iTRIQ Transcriptome Analysis

The main root fresh tissues were ground into powder in liquid nitrogen, homogenized in 1 mL lysis buffer (pH 7.6, 10% w/v SDS, 0.1 M dithiothreitol in 0.1 M Tris-HCl). The mixture was moved to an ice bath for 15 cycles (work 5s, stop 2s) for sonicating at 35 kHz. After super-centrifugation (30000 g, 4°C for 15 min), 10% trichloroacetic acid/acetone was added to the supernatant with three acetone washes. Protein concentration was measured by Bradford assay protein assay kit (Bio-Rad, United States) after dissolved in the lysis buffer and 100 µg aliquots of each sample were used for proteomic experiments. It was used to prepare peptide samples for iTRAQ experiments, including the Filter-aided sample preparation-method (FASP) by Wisniewski et al. (2009), employing ultrafiltration devices to perform SDS removal, buffer exchange, chemical modification, and protein digestion to obtain peptides mixtures.

Trypsin digestion of enzyme to protein ratio 1:50 was carried out at 37°C for 4 h. 4-plex iTRAQ labeling experiments were performed according to the manual

provided by AB SCIEX, Pte. Ltd. (Redwood City, CA, United States) with cation exchange conducted in a GeminiNX 5u C18 110A 150 mm × 4.6 mm column (Phenomenex, Guangzhou, China) by using a LC-20AB HPLC Pump system (Shimadzu, Japan) [33]. The condition of process included: uv wavelength: 214 nm; flow rate: 1000 μL/min; washing-gradient of liquid chromatography: Time(min), B%: 1, 5%; 18, 30%; 20, 80%; 24, 80%; 24.1, 5%; 30, stop. Reversed-phase liquid chromatography tandem mass spectrometry were performed by SAGENE, Co., Ltd. (Guangzhou, China) in C18 enriching column (3 μm, ID 100 μm, 20 mm length) and separation column (1.9 μm, ID75 μm, 100 mm length) by using a Triple TOF 6600 (Applied Biosystems, United States). The condition of process included: flow rate: 300 nL min<sup>-1</sup>; washing-gradient of liquid chromatography: Time(min), B%: 0.1, 8%; 55, 25%; 65, 80%; 70, 80%; 70.1, 2%; 75, 2%; stop. The peptides were identified by ProteinPilot 5.0 (AB Sciex) and matched to the reference transcripts of protein sequences of *P. notoginseng* in SwissProt/UniProt database. The statistically significant differentially-accumulated proteins between treatment groups were determined based on protein abundance difference multiple >1.3 or <0.77 ( $P < 0.05$ , t-test) [34]. KEGG pathway annotation was performed, and pathway overlaps with DEGs were identified.

### Statistical Analysis

The data of Cd content, saponin content and enzyme activities in the main root of *P. notoginseng* were statistically analyzed using one-way variance (ANOVA) with SPSS 19 software. Least significant difference (LSD) ( $P = 0.05$ ) were calculated.

## Results

### Soil pH, Soil Available Cd Content and Cd Content of *Panax notoginseng*

The application of lime had a greater impact on soil pH. With the amount of lime applied increased, the pH gradually increased. Under 0, 0.6, 3.0, 6.0, 9.0, 12.0 mg kg<sup>-1</sup> Cd treatments, the soil pH increased by 14.4-22.0%, 8.7-10.7%, 8.5-13.0%, 10.6-15.2%, 8.9-16.5%, and 17.0-19.0%, respectively, with 0.75 and 1.5 t ha<sup>-2</sup> lime application (Table 1).

When the application level of lime was consistent, the soil effective Cd contents increased significantly with the increase in Cd treatment levels. When the treatment level of Cd was the same, the soil effective Cd content decreased significantly with increase in the amount of lime applied (Table 1). The effective Cd content decreased by 3.3-38.0% with 0.75 and 1.5 t ha<sup>-2</sup> lime application compared to the non-lime treatment. Correlation analysis showed that there was a significant negative correlation between soil available Cd content and Ca/Cd treatment ratio ( $R = -0.738$ ,  $F = 9.589$ ,  $P < 0.05$ ,  $N = 10$ ).

With the amount of Cd treatment increased, the Cd content of the main root of *P. notoginseng* increased significantly. Under the same treatment amount of Cd, the Cd content of the main root of *P. notoginseng* decreases with the increase in lime application levels. Under 0, 3.0 and 12.0 mg kg<sup>-1</sup> Cd treatments, the Cd content of the main root of *P. notoginseng* was significantly decreased by 13.3%, 12.5% and 17.5% with 1.5 t ha<sup>-2</sup> lime application. Under 0.6, 6.0 and 9.0 mg kg<sup>-1</sup> Cd treatments, there was no significant changes in the Cd content in the main root of *P. notoginseng* (Table 1). Correlation analysis showed that there was a significant positive correlation between the Cd content of the main root and the soil effective Cd

Table 1. Soil pH and available Cd contents, Cd content in main root of *Panax notoginseng*.

Parameters	Lime /t ha <sup>-2</sup>	Cd/mg kg <sup>-1</sup>					
		0	0.6	3.0	6.0	9.0	12.0
Soil pH	0	5.42±0.17c	5.77±0.10b	5.75±0.03c	5.66±0.03b	5.64±0.12c	5.41±0.03b
	0.75	6.20±0.09b	6.27±0.10a	6.24±0.04b	6.26±0.21a	6.14±0.05b	6.33±0.07a
	1.5	6.61±0.1a	6.39±0.09a	6.50±0.14a	6.52±0.12a	6.57±0.11a	6.44±0.07a
Soil available Cd contents/mg kg <sup>-1</sup>	0	0.24±0.02a	0.79±0.04a	2.47±0.21a	4.58±0.18a	7.09±0.22a	8.32±0.16a
	0.75	0.18±0.02b	0.61±0.04b	2.24±0.16ab	4.43±0.34a	6.56±0.24a	7.74±0.19ab
	1.5	0.17±0.01b	0.49±0.02c	1.99±0.13b	4.16±0.32a	5.76±0.27b	7.13±0.40b
Cd contents in main root/mg kg <sup>-1</sup>	0	0.15±0.01a	0.31±0.02a	0.48±0.04a	0.89±0.03a	1.13±0.07a	1.43±0.10a
	0.75	0.14±0.00ab	0.31±0.02a	0.45±0.03ab	0.87±0.03a	1.05±0.08a	1.31±0.02a
	1.5	0.13±0.00b	0.29±0.02a	0.42±0.01b	0.84±0.03a	1.00±0.03a	1.18±0.04b

Note: The data are mean ± standard deviation. Different lowercase letters indicate difference significantly between lime treatments at  $P < 0.05$  levels with t-test,  $n = 3$ .



Table 2. Biomass of main root of biennial *Panax notoginseng* at flowering period (g plant<sup>-1</sup> dry weight).

Lime /t ha <sup>-2</sup>	Cd/mg kg <sup>-1</sup>					
	0	0.6	3.0	6.0	9.0	12.0
0	0.17±0.03a	0.18±0.03a	0.18±0.03c	0.27±0.01a	0.24±0.04a	0.25±0.04a
0.75	0.17±0.02a	0.26±0.08a	0.22±0.03b	0.25±0.02a	0.22±0.01a	0.20±0.04a
1.5	0.21±0.05a	0.17±0.02a	0.31±0.02a	0.21±0.04a	0.29±0.01a	0.27±0.03a

content ( $R = 0.992$ ,  $F = 502.665$ ,  $P < 0.001$ ,  $N = 10$ ), and there was a significant negative relationship between the Cd content of the main root and the Ca/Cd treatment ratio ( $R = -0.709$ ,  $F = 8.072$ ,  $P < 0.05$ ,  $N = 10$ ).

### Biomass and Morphological Characteristics of Roots

Under 3.0 mg kg<sup>-1</sup> Cd treatment, the biomass of main root increased by 22-72% with the increase in lime




Fig. 1. Effects of Cd and lime on root length, surface area and volume of *Panax notoginseng*.

application levels. Compared with other treatments, the biomass of main root of *P. notoginseng* was the highest with 3.0 mg kg<sup>-1</sup> Cd and 1.5 t ha<sup>-2</sup> lime treatment (Table 2).

Under the same lime treatment, the root length increased first and then decreased, and the root surface area and root volume decreased with the increase in Cd treatment levels. Under the 9.0 and 12.0 mg kg<sup>-1</sup> Cd treatments, the root length, root surface area and root volume under lime 1.5 t ha<sup>-2</sup> treatment were greater than those non-lime treatment (Fig. 1). Under 3.0 and 6.0 mg kg<sup>-1</sup> Cd treatments, the root surface area and root volume showed a decreasing trend with lime application. Correlation analysis showed that there were significant negative correlations between root length and soil available Cd content ( $R = -0.745$ ,  $F = 9.959$ ,  $P < 0.05$ ,  $N = 10$ ), Cd content of main root ( $R = -0.784$ ,  $F = 12.769$ ,  $P < 0.01$ ,  $N = 10$ ).

### Content of Saponins of *Panax notoginseng* and its Related Enzyme Activities

#### Saponins Contents

Under 0.6, 6.0 and 12.0 mg kg<sup>-1</sup> Cd treatments, the contents of saponins in the main root of *P. notoginseng* decreased with the increase in 0.75 and 1.5 t ha<sup>-2</sup> lime treatments. The content of saponins decreased by 4.8- 23.6% with lime application. The saponin content decreased by 23.6% under 0.6 mg kg<sup>-1</sup> Cd and 1.5 t ha<sup>-2</sup> lime application (Fig. 2).

The contents of three monomeric saponins showed in order: Ginsenoside Rg1 > ginsenoside Rb1 > notoginsenoside R1, three monomeric saponins and total saponin content reached the maximum at 0.75 t ha<sup>-2</sup> lime + 3.0 mg kg<sup>-1</sup> Cd treatment, which was 0.45% notoginsenoside R1, 1.62% ginsenoside Rg, 0.97% ginsenoside Rb1 and 3.03% total saponin. The ratio of Rg1 to Rb1 was the highest under 0.6 mg kg<sup>-1</sup> Cd treatment. With 0 and 0.75 t ha<sup>-2</sup> lime applications, the ratio of R1/(Rg1+Rb1) was the highest under 0.6 mg kg<sup>-1</sup> Cd treatment. The ratio of R1/(Rg1+Rb1) was the highest with 0.75 t ha<sup>-2</sup> lime application and 6 mg kg<sup>-1</sup> Cd treatment (Fig. 2). Correlation analysis showed that there were significant negative correlations between the ratio of Rg1/Rb1 and soil effective Cd content ( $R = -0.677$ ,  $F = 6.767$ ,  $P < 0.05$ ,  $N = 10$ ), Cd content of main root ( $R = -0.644$ ,  $F = 5.671$ ,  $P < 0.05$ ,




Fig. 2. Saponins contents and ratios of Rg1/Rb1 and R1/(Rg1+Rb1) in main root of *Panax notoginseng*.

$N = 10$ ); There was a significant quadratic correlation between the ratio of Rg1/Rb1 and the Ca/Cd treatment ratio ( $Y = 1.28 + 0.009X - 1.32 \times 10^{-5}X^2$ ,  $R = 0.815$ ,  $F = 6.925$ ,  $P < 0.05$ ,  $N = 10$ ). (The ratio of Rg1/Rb1 was the highest when Ca/Cd treatment ratio 682); There was a significant logarithm correlation between R1/(Rg1+Rb1) ratio and Ca/Cd treatment ratio ( $Y = 0.022 + \ln X$ ,  $R = 0.689$ ),  $F = 7.217$ ,  $P < 0.05$ ,  $N = 10$ ).

#### Related Enzyme Activities

The SS enzyme activity was the lowest under 12.0 mg kg<sup>-1</sup> Cd treatment. The SS activity in the main roots increased significantly under 0.6, 3.0 and 12 mg kg<sup>-1</sup> Cd treatment and 0.75 t ha<sup>-2</sup> lime application. With 1.5 t ha<sup>-2</sup> application, SS activity was significantly increased in the main root under 0.6 and 3.0 mg kg<sup>-1</sup> Cd treatments. The SS activity in the main root was the highest under 12.0 mg kg<sup>-1</sup> Cd treatment and 0.75 t ha<sup>-2</sup> lime application (Table 3). Correlation analysis

showed that there were negative relationships between SS activity and soil effective Cd content ( $R = -0.747$ ,  $F = 10.102$ ,  $P < 0.05$ ,  $N = 10$ ), Cd content of main root ( $R = -0.765$ ,  $F = 11.317$ ,  $P < 0.05$ ,  $N = 10$ ). There were significant positive correlations between SS activity and R1 content ( $R = 0.723$ ,  $F = 8.745$ ,  $P < 0.05$ ,  $N = 10$ ), Rg1 content ( $R = 0.646$ ,  $F = 5.717$ ,  $P < 0.05$ ,  $N = 10$ ).

Mevalonate kinase (MVK) activities increased under 0.6 mg kg<sup>-1</sup> Cd treatment without application of lime. MVK activity was inhibited under other Cd treatments, and the inhibitory effect increased with increase in Cd treatment concentrations. Under the same Cd treatment, the activities of mevalonate kinase gradually decreased with the increase in lime application levels. There was a significant difference in the MVK activity under 0 and 3.0 mg kg<sup>-1</sup> Cd treatments with lime application comparing with non-lime application. There was no significant difference of MVK activity between 0.75 t ha<sup>-2</sup> and 1.5 t ha<sup>-2</sup> lime applications. Under 6.0 mg kg<sup>-1</sup> Cd treatment, there was no significant difference between the applications of 0.75 t ha<sup>-2</sup> lime and non-lime. The MVK activity decreased significantly when 1.5 t ha<sup>-2</sup> lime application. Under 9.0 mg kg<sup>-1</sup> Cd treatment or more, the MVK activities significantly decreased. There were significant difference of MVK activities between 0.75 t ha<sup>-2</sup> and 1.5 t ha<sup>-2</sup> application.

P450 reductase activity decreased with the increase in Cd treatment concentrations under the same lime application. Under the same Cd treatment level, P450 reductase activity decreased with the increase in lime application levels. Under 0, 0.6, 3.0, 6.0, 9.0 and 12.0 mg kg<sup>-1</sup> Cd treatments, the P450 reductase activities with 0.75 t ha<sup>-2</sup> and 1.5 t ha<sup>-2</sup> lime application reduced by 27.7-40.8%, 10.0-37.2%, 38.6-41.9%, 24.6-32.6%, 40.5-60%, and 45.2-63.3% compared with non-lime application, respectively. Correlation analysis showed that there were significant negative correlations between P450 reductase activity and soil effective Cd content ( $R = -0.770$ ,  $F = 11.658$ ,  $P < 0.01$ ,  $N = 10$ ), Cd content of main root ( $R = -0.750$ ,  $F = 10.257$ ,  $P < 0.05$ ,  $N = 10$ ). There was a significant quadratic correlation between P450 reductase activity and Ca/Cd treatment ratio ( $Y = 0.85 + 0.006X - 8.62 \times 10^{-6}X^2$ ,  $R = 0.809$ ,  $F = 6.612$ ,  $P < 0.05$ ,  $N = 10$ ) (P450 reductase activity was highest when Ca/Cd treatment ratio was 696). There were significant positive correlations between P450 reductase activity and R1 content ( $R = 0.716$ ,  $F = 8.433$ ,  $P < 0.05$ ,  $N = 10$ ), ratio of R1/(Rg1+Rb1) ( $R = 0.709$ ,  $F = 8.109$ ,  $P < 0.05$ ,  $N = 10$ ).

When lime was not applied, the  $\beta$ -amyrin synthase ( $\beta$ -AS) activity decreased with increase in 0-6.0 mg kg<sup>-1</sup> Cd treatment concentration. The  $\beta$ -AS activities gradually decreased with increase in Cd treatment concentrations under 0.75 t ha<sup>-2</sup> and 1.5 t ha<sup>-2</sup> lime application. The  $\beta$ -AS activities decreased with increase in lime application levels under the same Cd treatment concentration. The  $\beta$ -AS activity was 75.95 U g<sup>-1</sup>, which was the lowest under 0.75 t ha<sup>-2</sup> application and 9.0 mg kg<sup>-1</sup> Cd treatment. The  $\beta$ -AS


Table 3. Activities of SS, MVK, P450 and  $\beta$ -AS in main root of *Panax notoginseng*.

Enzymes/U g <sup>-1</sup>	Lime /t ha <sup>-2</sup>	Cd/mg kg <sup>-1</sup>					
		0	0.6	3.0	6.0	9.0	12.0
SS	0	0.17±0.01a	0.17±0.01b	0.18±0.01b	0.21±0.02a	0.18±0.01a	0.15±0.01b
	0.75	0.17±0.01a	0.22±0.07a	0.25±0.06a	0.18±0.01a	0.17±0.01a	0.18±0.01a
	1.5	0.17±0.01a	0.20±0.02a	0.20±0.01a	0.19±0.01a	0.18±0.02a	0.15±0.02b
MVK	0	9.45±0.82a	10.33±0.76a	8.50±0.39a	7.69±0.59a	7.47±0.88a	6.05±0.88a
	0.75	7.04±0.55b	7.48±0.44b	6.28±0.28b	6.85±1.06a	5.06±0.69b	6.82±0.34b
	1.5	5.64±1.47b	8.13±0.45b	5.30±0.12b	5.19±1.06b	7.68±0.33c	3.40±0.23c
P450	0	3.11±0.13a	2.31±0.20a	2.15±0.03a	1.75±0.16a	2.15±0.21a	1.77±0.42a
	0.75	2.25±0.45b	2.08±0.21a	1.32±0.14b	1.32±0.36ab	1.28±0.43b	0.97±0.22b
	1.5	1.84±0.03b	1.45±0.21b	1.25±0.48b	1.18±0.16b	0.86±0.08b	0.65±0.14b
$\beta$ -AS	0	115.57±19.03a	107.87±13.40a	101.11±20.10a	90.30±10.58a	137.08±1.12a	125.71±20.40a
	0.75	110.83±16.86a	98.85±10.86ab	92.76±14.71a	77.06±16.29a	75.95±17.81b	80.44±7.84b
	1.5	86.10±4.46a	81.01±5.76c	52.68±9.29b	53.01±5.18b	37.71±3.90c	23.74±10.75c

activity was 23.74 U g<sup>-1</sup>, which was the lowest under 1.5 t ha<sup>-2</sup> application and 12.0 mg kg<sup>-1</sup> Cd treatment. There were significant positive correlations between  $\beta$ -AS activity and R1 content ( $R = 0.866$ ,  $F = 24.082$ ,  $P < 0.01$ ,  $N = 10$ ), ratio of R1/(Rg1 + Rb1) ( $R = 0.629$ ,  $F = 5.230$ ,  $P < 0.05$ ,  $N = 10$ ).

#### Expression of Related Proteins in the Synthesis of Saponins of *Panax notoginseng*

When lime was applied under Cd stress, 328122 total spectra were obtained by iTRAQ analysis with unique spectra 5918, unique peptide 2862, and 1181

Fig. 3. Saponins biosynthesis pathway of *Panax notoginseng*.

Note: mevalonate kinase (MVK), geranyl pyrophosphate synthase (GPS), farnesyl diphosphate synthase (FPS), squalene synthase (SS), squalene epoxidase (SE), dammarenediol synthase (DS), cytochrome P450,  $\beta$ -amyrin synthase ( $\beta$ -AS), glycosyl transferase (GT).

Table 4. Functional classification of differentially expressed proteins (DEPs).

DEPs	Protein ID	Coverage	Unique spectrum	Unique peptide	Difference multiple $\pm$ SD	P value	Up/Down	Comparison group
P450( <i>CYP72A129</i> )	Unigene0012926	0.095	1	1	0.66 $\pm$ 0.08	0.000	Down	L <sub>L</sub> Cd <sub>6</sub> /L <sub>0</sub> Cd <sub>0</sub>
	Unigene0050708	0.024	1	1	0.58 $\pm$ 0.23	0.006	Down	L <sub>0</sub> Cd <sub>6</sub> /L <sub>0</sub> Cd <sub>0</sub>
P450 ( <i>CYP72A129</i> -like)	Unigene0047723	0.045	1	1	0.59 $\pm$ 0.09	0.000	Down	L <sub>L</sub> Cd <sub>6</sub> /L <sub>0</sub> Cd <sub>0</sub>
P450( <i>CYP736A54</i> )	Unigene0014832	0.051	1	1	1.75 $\pm$ 0.58	0.025	Up	L <sub>L</sub> Cd <sub>6</sub> /L <sub>0</sub> Cd <sub>0</sub>
		0.051	1	1	1.54 $\pm$ 0.39	0.024	Up	L <sub>0</sub> Cd <sub>6</sub> /L <sub>0</sub> Cd <sub>0</sub>
P450 reductase	Unigene0031564	0.074	1	1	0.66 $\pm$ 0.07	0.002	Down	L <sub>L</sub> Cd <sub>6</sub> /L <sub>0</sub> Cd <sub>0</sub>
$\beta$ -AS	Unigene0011395	0.028	1	1	0.62 $\pm$ 0.05	0.000	Down	L <sub>L</sub> Cd <sub>6</sub> /L <sub>0</sub> Cd <sub>0</sub>
glycosyl transferase UDP84B1	Unigene0032248	0.022	1	1	2.14 $\pm$ 0.76	0.015	Up	L <sub>L</sub> Cd <sub>6</sub> /L <sub>0</sub> Cd <sub>0</sub>
UDP-glycosyl transferase	Unigene0049934	0.108	6	5	0.63 $\pm$ 0.08	0.003	Down	L <sub>L</sub> Cd <sub>6</sub> -/L <sub>0</sub> Cd <sub>0</sub>
UDP-glycosyl transferase ( <i>UGTPg34</i> )	Unigene0003621	0.035	1	1	0.64 $\pm$ 0.28	0.024	Down	L <sub>0</sub> Cd <sub>6</sub> /L <sub>0</sub> Cd <sub>0</sub>
		0.35	1	1	0.66 $\pm$ 0.18	0.029	Down	L <sub>L</sub> Cd <sub>6</sub> /L <sub>0</sub> Cd <sub>0</sub>
		0.35	1	1	0.61 $\pm$ 0.26	0.015	Down	L <sub>L</sub> Cd <sub>6</sub> /L <sub>0</sub> Cd <sub>0</sub>
UDP-glycosyl transferase( <i>UGTPg19</i> )	Unigene0040383	0.025	1	1	0.64 $\pm$ 0.09	0.000	Down	L <sub>L</sub> Cd <sub>6</sub> /L <sub>0</sub> Cd <sub>0</sub>

Note: L<sub>0</sub> and L<sub>L</sub> were 0 and 0.75 t ha<sup>-2</sup> lime application. Cd<sub>0</sub> and Cd<sub>6</sub> were 0 and 6.0 mg kg<sup>-1</sup> Cd treatment.

differential proteins identified, in which the expression of 10 proteins related to saponin synthesis was significantly different ( $P < 0.05$ ), including P450,  $\beta$ -AS and glycosyl transferase (GT) based on GO and KEGG analysis (Table 4).

The expression of P450 reductase,  $\beta$ -AS, glycosyl transferase UGTPg34, and glycosyl transferase UDP84B1 were down-regulated under 0.75 t ha<sup>-2</sup> lime application compared to non lime application (Table 4). The glycosyl transferase UDP84B1 protein homologous to the grape *Vitis vinifera* was down-regulated. The UGTPg19 protein homologous to ginseng was down-regulated.

The expression of P450 (*CYP72A129*) protein homologous to *Panax ginseng* was down-regulated under 6.0 mg kg<sup>-1</sup> Cd treatment. The expression of P450 (*CYP736A54*) protein homologous to *Bupleurum chinense* was up-regulated. The expression of the glycosyl transferase UGTPg34 protein homologous to *Panax ginseng* was down-regulated.

When treated with 6.0 mg kg<sup>-1</sup> Cd + 0.75 t ha<sup>-2</sup> lime, the expression of P450 (*CYP72A129* and *CYP72A129*-like) proteins homologous to *Panax ginseng* was down-regulated. The expression of P450 (*CYP736A54*) protein homologous to *Bupleurum chinense* was up-regulated. The expression of glycosyl transferase UGTPg19 and UGTPg34 protein homologous to *Panax ginseng* was down-regulated, and the expression of glycosyl transferase 84B1 protein was up-regulated.

## Discussion

### Effects of Cd Stress on the Growth and Saponin Content in Roots of *Panax notoginseng* under Lime Application

Soil pH is one of the most important factors affecting soil Cd formation and bioavailability. The change of pH has a great influence on the effective fraction of Cd in soil. The OH<sup>-</sup> ion in Ca(OH)<sub>2</sub> not only increases the pH value of the soil, but also promotes the formation of hydroxyl metal [35]. The hydroxyl metal ion has a strong affinity with the adsorption site of the soil and can be effectively adsorbed on the soil surface. The increase of OH<sup>-</sup> in the soil causes Cd<sup>2+</sup> to combine with CO<sub>3</sub><sup>2-</sup> and OH<sup>-</sup> to form insoluble precipitates, which are fixed in the soil and reduce the effective Cd content in the soil [36, 37]. The increase of pH will lead to the decrease of hydrogen ions and free metal cations, reduce the competition with Cd<sup>2+</sup> on soil adsorption, facilitate the adsorption of Cd and soil surface, and reduce the content of available Cd in soil. However, the reduction of available Cd in the soil by lime depends on the form and dose of lime, especially in the case of high Cd content in the soil. Application of lime increases soil pH and reduces soil available Cd content. Correlation analysis showed that there was a significant negative correlation between the ratio of Ca/Cd treatment concentration and soil available Cd content and Cd content of main root ( $P < 0.05$ ). It is indicated that the concentration ratio of Ca/Cd treatment be an important



factor affecting the available Cd content in soil and the Cd content in the main root of *P. notoginseng*. With the increase of calcium ion concentration, the desorption of Cd increases, while the high concentration of Cd in soil solution can reduce the absorption and transport of calcium by roots, increase the absorption of Cd, reduce the absorption of calcium by plants [38, 39].

Under the same lime application, the effect of cadmium stress on biomass, root length, root surface area and root volume of *P. notoginseng* had a dose effect. Correlation analysis showed that there were significant negative correlations between root length and soil effective Cd content ( $P < 0.05$ ), main root Cd content ( $P < 0.01$ ). Under certain cadmium stress, the application of lime can increase the biomass, root length, root surface area and root volume of roots to some extent. Ca plays an important role in reducing the absorption of Cd by plants and reducing the toxicity of Cd to plants [38,40]. Exogenous Ca treatment alleviated the DNA damage of Arabidopsis seedling root cells induced by Cd stress. Ca-binding protein is involved in the regulation of biological gene expression, DNA synthesis, repair and transcription to promote plant root growth[41]. Calcium not only acts as a plant's essential nutrient element, but more importantly, as a second intracellular signal transduction, initiates a series of physiological and biochemical processes and activates multiple resistance mechanisms in plants [42]. High concentrations of exogenous Ca treatment may increase solution concentration, cause osmotic stress and ionic toxicity, and may even aggravate Cd stress damage [31].

Under the same Cd stress, the saponin contents decreased with lime application. There may be competition between calcium and cadmium ions with lime application. With low Cd concentration treatments, the effect of lime on the synthesis of saponins was greater. With high Cd concentration treatments, the effect of Cd on the synthesis of saponin was dominant. There was a significant negative correlation between the content of saponins and Cd treatment concentration ( $R = -0.734$  and  $-0.579$ ,  $P < 0.05$ ,  $n = 6$ , respectively, for 0.75 and 1.5 t ha<sup>-2</sup> lime application treatments). Exogenous stress affects the synthesis of secondary metabolites by affecting the expression of key enzyme genes. For example, methyl jasmonate and ethyl jasmonate induced the expressions of SS and SE genes in *P. notoginseng*, and increased the accumulation of total saponins of *P. notoginseng* [37]. Liang et al (2011) found that high concentration B and Mn stress inhibited the expression of SS gene in key glycyrrhizic acid synthesis [43]. A certain concentration of Zn and Mo stress promoted the expression of SS gene. Zhu et al. (2017) showed a negative correlation between saponin content and soil Cd content in *Paris polyphylla* var. *Yunnanensis* [44]. Under treated with 30 mg kg<sup>-1</sup> Cd, notoginsenoside R1, ginsenoside Rb1, ginsenoside Rg1 and total saponins decreased [20]. The optimal application of lime may alleviate the toxicity of Cd stress to *P. notoginseng*.

The heterogeneity of ginsenoside Rg1, ginsenoside Rb1 and notoginsenoside R1 content was related to the balance of calcium and cadmium. Correlation analysis showed that there were significant negative correlations between Rg1/Rb1 ratio and soil available Cd content ( $P < 0.05$ ), Cd content of main root ( $P < 0.05$ ). There was a significant negative quadratic correlation ( $P < 0.05$ ) between the ratio of Rg1/Rb1 and Ca/Cd treatment, which was a signal-peak modal. It could be suggested that the conversion of the diol saponin Rb1 to the triol saponin Rg1 is not only inhibited by cadmium, but also affected by the balance of calcium and cadmium. Under the ratio of Ca/Cd treatment < 680, the conversion of the diol saponin Rb1 to the triol saponin Rg1 was promoted. Under the ratio of Ca/Cd treatment > 680, the conversion might be inhibited. There was a significant logarithmic correlation between the R1/(Rg1+Rb1) ratio and the Ca/Cd treatment ratio ( $P < 0.05$ ). Therefore, the higher the ratio of Ca/Cd treatment was, the greater the ratio of the content of notoginsenoside R1 to the total saponin content was. The heterogeneity of saponin monomer was affected by the balance of calcium and cadmium. The inhibition of saponin synthesis by cadmium and calcium occurred at different stages of metabolism. The inhibitory effect of cadmium on the conversion process of glycolic saponins to triol saponins was more prominent.

#### Enzymology and Proteomics Mechanisms of the Metabolism of Saponin under Cd Stress with Lime Application

Saponin is a dammarane-type tetracyclic triterpene whose synthesis pathway includes the mevalonate pathway and the 2C-methyl-4-phosphate-4D-erythritol pathway [45,46]. The mevalonate pathway is dominant, which is co-catalyzed by a number of enzymes. Mevalonate is catalyzed by three consecutive ATP-dependent enzymes (mevalonate kinase (MVK), phosphomevalonate and mevalonate pyrophosphate decarboxylase) to form isopentenyl pyrophosphate [21]. Isopentenyl pyrophosphate ultimately produces triterpenoid saponins by catalysis and modification of a series of enzymes. Mevalonate kinase is one of the rate-limiting enzymes in the mevalonate pathway. It is present in the cytoplasm and has a highly conserved domain. It has two conserved regions that bind to the substrate and ATP, and undergoes catalytic reaction. MVK is the first ATP-dependent enzyme [47, 48]. The activity of mevalonate kinase increased under 0.6 mg kg<sup>-1</sup> Cd treatment. The activity of mevalonate kinase decreased with the increase in Cd treatment concentrations, indicating the dose effect. This result was similar to findings of Zhu et al. (2014) [20]. It may be due to the low concentration of Cd initiating its own defense system. The mevalonate kinase activity was inhibited under 0.75 t ha<sup>-2</sup> lime application. Under the same Cd treatment, the activity of mevalonate

kinase gradually decreased with the increase in lime application levels. The inhibitory effect of Ca and Cd on mevalonate kinase activity may be due to an ATP-dependent enzyme of MVK. Calcium and Cadmium reduced the absorption of phosphorus by plants, inhibited the energy metabolism of plants and the supply of ATP. The ions  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Al}^{3+}$ , and  $\text{Fe}^{3+}$  in the soil were susceptible to chemical precipitation with phosphate, which caused most of the soil phosphorus to be converted into insoluble fractions, reduced the absorption and utilization of phosphorus by plants [49,50].

SS is a key enzyme in the synthesis of saponins. There were significant positive correlations ( $P < 0.05$ ) between SS activity and R1 content, Rg1 content. Under the application of lime, the SS activity in the main root was increased to some extent, which suggested that SS activity be promoted by Ca. Cd stress decreased SS enzyme activity. There were significant negative correlations ( $P < 0.05$ ) between SS activity, soil effective Cd content, and Cd content of main root.

In the catalytic process, cytochrome P450 reductase combines with heme to form a thiol-ferrous-carbon monoxide complex. The complex has the largest characteristic absorption spectrum at 450 nm, which is dependent on Heme, a family of multifunctional metal-containing enzymes that catalyze a variety of complex oxidation reactions [51]. Cytochrome P450 (Cytochrome P450) catalyzes the hydroxylation and oxidation of the triterpene skeleton [52]. Cytochrome P450 is a type of oxidase encoded by a supergene family. Their expression products are similar and their substrates are different. It requires multiple cytochrome P450s to participate in the synthesis of the final product. In the case of same the amount of lime applied, the P450 reductase activity decreased with increase in Cd treatment concentrations. There were significant negative correlations between P450 reductase activity and soil available Cd content ( $P < 0.01$ ), Cd content of main root ( $P < 0.05$ ). The expression of P450 gene of *P. notoginseng* under the exogenous Cd stress up-regulated at the low concentration, down-regulated at concentration. Under the exogenous Cd stress, P450 gene expression did not show significant correlation with saponins R1, ginsenoside Rb1, ginsenoside Rg1 and total saponins. In this study, P450 (CYP72A129) protein homologous to *Panax ginseng* was down-regulated when treated with  $6.0 \text{ mg kg}^{-1} \text{ Cd}^{2+}$ . The expression of P450 (CYP736A54) protein homologous to *Bupleurum chinense* was up-regulated. It suggested that different P450 isoenzyme activities and expression levels have different responses to Cd stress. Saponin synthesis and metabolism could not be indicated by only one gene express of P450.

Under the same Cd treatment concentration, the inhibitory effect of cadmium on cytochrome P450 was alleviated with  $0.75 \text{ t ha}^{-2}$  lime application at low Cd treatment concentrations. There was a significant quadratic correlation between P450 reductase activity

and Ca/Cd treatment ratio ( $P < 0.05$ ), which was a single-peak model. Under the ratio of Ca/Cd treatment 696, the activity of P450 reductase was the highest. It suggested that the balance of Ca/Cd be important for the activity of P450 reductase. Under 0.75 and  $1.5 \text{ t ha}^{-2}$  lime application, there were significant positive correlations between the saponins content of main root and P450 reductase activity, P450 reductase activity and R1 content, ratio of  $\text{R1}/(\text{Rg1}+\text{Rb1})$  ( $P < 0.05$ ). Under  $6.0 \text{ mg kg}^{-1} \text{ Cd}^{2+}$   $0.75 \text{ t ha}^{-2}$  lime application treatment, protein expressions of P450 (CYP72A129 and CYP72A129-like) homologous to *Panax ginseng* was down-regulated. The protein expression of P450 (CYP736A54) homologous to *Bupleurum chinense* was up-regulated. In the presence of calcium, the difference in response of P450 isoenzymes was the similar as under Cd treatment. Therefore, the decrease in the content of saponins might be related to the decrease of P450 reductase activity and the down-regulation of P450 CYP72A129 protein expression. The P450 reductase was involved in the synthesis of triol type saponin, inhibiting the conversion of diol type saponin to triol type saponin. Since P450 contains  $\text{Fe}^{2+}$  ions, the inhibitory effect of cadmium and calcium on P450 activity may be related to decrease in uptake of  $\text{Fe}^{2+}$  due to cadmium and calcium replacing  $\text{Fe}^{2+}$  at the active site, resulting in a decrease in P450 activity. Because of changing the pH with lime application, the availability of iron in the soil decreased, limiting the absorption of iron from the soil by plants [53].

In the saponin synthesis pathway, the cyclization of 2,3-oxidosqualene is catalyzed by oxidosqualene cyclases (OSC) [54]. The  $\beta$ -AS belongs to the OSC superfamily, which catalyzes the cyclization of 2,3-oxidized squalene to  $\beta$ -aromatic alcohol and promotes the reaction into the triterpenoid synthesis tributary [55]. There were significant positive correlations between  $\beta$ -AS activity and R1 content ( $P < 0.01$ ),  $\text{R1}/(\text{Rg1}+\text{Rb1})$  ratio ( $P < 0.05$ ). The activities of  $\beta$ -AS decreased with increase in  $0\text{-}6.0 \text{ mg kg}^{-1} \text{ Cd}^{2+}$  treatment concentrations. Under the same Cd treatment concentration, the  $\beta$ -AS activity decreases with increase in lime application levels. The activities of  $\beta$ -AS was inhibited and its protein expression was down-regulated with the lime application. The activities of  $\beta$ -AS similar to P450 reductase, participates in the synthesis of triol saponin. The activities of  $\beta$ -AS and the synthesis of triol saponin were inhibited by both lime and Cd treatments. The decrease in the content of notoginsenoside might be related to the decreased activities of P450 reductase and  $\beta$ -AS.

The synthesis of triterpenoid saponins is accomplished by adding a glycosyl group to the saponin by glycosyl transferase (GT) [56]. UGTs are a subfamily of GTs whose main function is to be responsible for the glycosylation of saponins. The triterpenoid compound forms a sugar chain by glycosylation of a glycosyltransferase at positions C3, C16, C28, to produce saponin. Sugar chain donors

include UDP-glucose, UDP-rhamnose, UDP-galactose, UDP-xylose, and UDP-arabinose [4, 57, 58]. When Cd was treated and lime was applied, UGTPg34 was down-regulated. The expression of glycosyltransferases UGTPg19 and UGTPg34, which are homologous to *P. ginseng*, was down-regulated, and the expression of glycosyltransferase 84B1 was up-regulated under Cd treatments. The expression of the glycosyl transferase GTs was down-regulated due to both Cd and lime treatments, which affected the saponin content. Because Cd<sup>2+</sup> might be bound to the active site-thiol group and the imidazole-containing ligand in the enzyme molecule to form a stable complex, thereby competing with the substrate and inhibiting the enzyme activity.

Therefore, although calcium promoted SS activity, excessive cadmium and calcium led to decrease in the activities of mevalonate kinase, P450 reductase,  $\beta$ -fragrant alcohol synthase and glycosyltransferase and their protein expression in the saponin synthesis pathway. The Ca/Cd ratio met a single-peak model with the conversion of the diolic saponin to the triol saponin. The effects of cadmium/lime on saponin synthesis by the expression and activity of key enzymes in the upstream MEP/DOXP pathway (terpenoid backbone biosynthesis) should be paid more attention in the further research.

### Conclusion

With the increased in lime application levels under Cd stress, the soil available Cd content and the Cd content in the main root of *P. notoginseng* significantly reduced, the biomass and root length of the main root of *P. notoginseng* increased. The content of saponins decreased with the activities and expressions of mevalonate kinase (MVK), P450 reductase and  $\beta$ -AS decreased. When the Ca/Cd ratio less than 680, the synthesis of triol type saponin and P450 activity were promoted with the application of lime. It is recommended the lime application rate should be less than 0.948 and 4.59 t ha<sup>-2</sup> lime under 0.6 and 3.0 mg kg<sup>-1</sup> Cd treatments, respectively, based on quadratic correlation between the ratio of Rg1/Rb1 and the Ca/Cd treatment ratio. The results indicate that excessive application of calcium or cadmium stress might inhibit the conversion of glycol type to triol type saponin and glycosylation, change the heterogeneity of saponin monomer, and reduce the total amount of saponin.

### Acknowledgements

The authors would like to thank to The National Natural Science Foundation of China (Grant Nos. 31560163 and 41867055) and The Yunnan Key Research and Development Project (2019BC001-04).

### Conflict of Interest

The authors declare no conflict of interest.

### References

1. YANG H.F., WANG Y.B., LI J.L. Impacts of Cu and Zn pollution on rape (*Brassica chinensis* L.) growth and accumulation effect of heavy metals in paddy soil. *Ecol. Environ. Sci.*, **20** (10), 1470, **2011**.
2. WEI B.G., YANG L.S. A review of heavy metal contaminations in urban soils, urban road dusts and agricultural soils from China. *Microchem. J.*, **94** (2), 99, **2010**.
3. XUE L., LIU J.F., SI S.Q., WEI Y., CHANG E.M., GAO M., JIANG Z.P. A review on the progress of proteomic study on plant responses to heavy metals. *Acta Prataculturae Sin.*, **22** (4), 300, **2013**.
4. ZHANG R.R., ZHANG P., DU S.T. Oxidative stress related signals and their regulation under Cd stress: A review. *Chin. J. Appl. Ecol.*, **27** (3), 981, **2016**.
5. HUANG B.F., XIN J.L. Mechanisms of heavy metal accumulation in plants: a review. *Acta Pratacu. Sin.*, **22** (1), 300, **2013**.
6. WANG X.H., DAI L. Physiology and biochemical mechanism of the cadmium in crop roots. *Sci. Agric. Sin.*, **49** (22), 4323, **2016**.
7. PEREZ-CHACA M.V., RODRÍGUEZ-SERRANO M., MOLINA A.S., PEDRANZANI H.E., ZIRULNIK F., SANDALIO L.M., ROMERO-PUERTAS M.C. Cadmium induces two waves of reactive oxygen species in *Glycine max* (L) roots. *Plant Cell Environ.*, **37**, 1672, **2014**.
8. PITERKOV J., LUHOV L., NAVTILOV B., SEDLÁŘOVÁ M., PETŘIVALSKÝ M. Early and long-term responses of cucumber cells to high cadmium concentration are modulated by nitric oxide and reactive oxygen species. *Acta Physio. Plant.*, **37**, 1-12, 19, **2015**.
9. XIE H.L., LIU J., CHEN S., WANG J.Y., FU W., LI Y.P., WANG H., XIAO Q.T., ZHENG X.Y., HUANG J.W., LIN R.Y., LIN W.X. Analysis of differentially expressed proteins in *Perilla frutescens* (L.) Britt. leaves under cadmium stress. *Chin. J. Ecol-Agric.*, **22** (10), 1207, **2014**.
10. ROTH U., VON ROEPENACK-LAHAYE E., CLEMENS S. Proteome changes in *Arabidopsis thaliana* roots upon exposure to Cd<sup>2+</sup>. *J. Exp. Bot.*, **57** (15), 4003, **2006**.
11. KIEFFER P., DOMMES J., HOFFMANN L., HAUSMAN J.F., RENAUT J. Quantitative changes in protein expression of cadmium-exposed poplar plant. *Proteomics.*, **8**, 2514, **2008**.
12. DOHERTY M.K., WHITFIELD P.D. Proteomics moves from expression to turnover: update and future perspective. *Expert. Rev. Proteomics.*, **8** (3), 325, **2011**.
13. LI Z.D., ZHAO R.H., ZHANG Z.C., YU J., GU W., HE S., CAO G.H. Advances in biosynthesis and regulation mechanism of notoginseng saponins. *Chin. J. Exp. Tradi. Med. Formulae.*, **24** (14), 71, **2018**.
14. ZHANG W.B., TAO S.G., ZENG H.C., ZHOU J.M. HPLC determination of Xuesaitong capsules ginsenoside Rg1, Rb1 and the content of notoginsenoside R1. *Spec. Wild Econ. Animal Plant Res.*, **4**, 49, **2010**.
15. SHI L., GE F., LIU D.Q., CHEN C.Y. Advances on biosynthesis of *Panax notoginseng* saponins and regulation of key enzymes. *Acta Bot. Boreali-Occidentalia Sin.*, **30** (11), 2358, **2010**.



16. ZU Y.Q., MEI X.Y., MIN Q., SU Y., MA N., FENG G.Q., LI Y. Effects of As stress on contents of saponin and flavonoid, key enzymes activities of *Panax notoginseng* and its proteomic analysis. *Chin. J. Appl. Ecol.*, **27** (12), 4013, **2016**.
17. ZHU H., TENG Y., ZHANG M.Y., MA W.T., LIU W.X., REN W.J., LUO Y.M., LI Z.G., GUO D. Pollutant accumulation and microbiological characteristics change in notoginseng-planting soils. *Soils*, **47** (1), 121, **2015**.
18. ZU Y.Q., CHENG S.C., KE H.L., GUO X.H., WU J., LI Y. Distribution of Pb, Cd, Cu and Zn in plant of *Panax notoginseng* and soil in Wenshan Prefecture, a *Panax notoginseng* growing region. *J. Ecol. Rural. Environ.*, **33** (4), 317, **2017**.
19. ZHANG Y.C., SONG F.R., LIU Z.Q., WANG S.M., HOU Y.B., WU C. Effect of treating ginseng seeds with plasma ray radiation/magnetization on content of ginsenosides. *Chin. J. Appl. Chem.*, **3**, 311, **2009**.
20. ZHU M.L., CHEN Z.J., JIANG Y., WEI F.G., CUI B., JIANG T.X., CAI H.B., ZHANG W.S. Cadmium stress of soil on cadmium enrichment in different tissues and its main active compositions of *Panax notoginseng*. *Chin. Tradit. Patent Med.*, **36** (2), 342, **2014**.
21. GUO X., LUO H.M., CHEN S.L. Cloning and analysis of mevalonate kinase (PnMVK1) gene in *Panax notoginseng*. *Acta Pharm. Sin.*, **47** (8), 1092, **2012**.
22. CHEN S.J., PAN W.J., MEN Y.S., ZONG X.F., WANG S.G. Effects of acid soils treated with lime and polyacrylamide on the growth and physiological characteristics of flue-cured tobacco. *Plant Nutr. Fert. Sci.*, **18** (5), 1243, **2012**.
23. POOVAIAH B.W. Biochemical and molecular aspects of calcium action. *Acta Hort.*, **326**, 139, **1993**.
24. SCHÖNKNECHT G. Calcium signals from the vacuole. *Plants*, **2**, 589, **2013**.
25. NOMURAA H., SHIINA T. Calcium signaling in plant endosymbiotic organelles: mechanism and role in physiology. *Mol. Plant.*, **7**, 1094, **2014**.
26. ZHENG Y., CHEN Z.J. Organellar calcium signaling in plants. *Plant Physiol. Commun.*, **51** (8), 1195, **2015**.
27. GIACOMELLO M., DRAGO I., PIZZO P., POZZAN T. Mitochondrial Ca<sup>2+</sup> as a key regulator of cell life and death. *Cell Death Differ.*, **14**, 1267, **2007**.
28. SEIGNEURIN BERNY D., GRAVOT A., AUROY P., MAZARD C., KRAUT A., FINAZZI G., GRUNWALD D., RAPPAPORT F., VAVASSEUR A., JOYARD J., RICHAUD P., ROLLAND N. HMA1, a new Cu-ATPase of the chloroplast envelope, is essential for growth under adverse light conditions. *J. Biol. Chem.*, **281**, 2882, **2006**.
29. LIN L.L., SHEN L., LIN Y.Z., ZHONG F.L., YE L.P., QU X.L. The role of calcium-signal in plant adaptation to abiotic stress. *Chin. Hortic. Abstract.*, **4**, 41, **2015**.
30. SI L., GUO C.H., LIANG F., PENG C., XIAO X.Y., FENG W.L. Effects of lime and water management on uptake and translocation of cadmium in rice. *Transactions Chin. Soc. Agric. Eng.*, **33** (24), 111, **2017**.
31. WAN C.Y., WANG Y.T., ZENG W.L., LI S.S. Alleviation of Cd Toxicity in *Arabidopsis thaliana* Seedlings by Exogenous Ca<sup>2+</sup> or K<sup>+</sup>. *Chin. Bull. Bot.*, **49** (3), 262, **2014**.
32. WISNIEWSKI J.R., ZOUGMAN A., NAGARAJ N., MANN M. Universal sample preparation method for proteome analysis. *Nature methods*, **6**, 359, **2009**.
33. ROSS P.L., HUANG Y. N., MARCHESE J.N., WILLIAMSON B., PARKER K., HATTAN S. Multiplexed protein quantitation in *Saccharomyces cerevisiae* using amine-reactive isobaric tagging reagents. *Mol. Cell. Proteom.*, **3**, 1154, **2004**.
34. SZKLARCZYK D., MORRIS J.H., COOK H., KUHN M., WYDER S., SIMONOVIC M. The STRING database in 2017: quality-controlled protein-protein association networks, made broadly accessible. *Nucleic Acids Res.*, **45**, 362, **2017**.
35. LI P., WANG X.Q., LANG M., ZHANG T.L. Effects of amendments on the fraction transform of heavy metals in soil contaminated by copper and cadmium. *Chin. Environ. Sci.*, **32** (7), 1241, **2012**.
36. ZHOU X.Y., FENG W.Q., QIN Y.S., YU H., LIAO M.L., LIU Y.C., WANG C.Q., TU S.H. Effect of magnesium, manganese, activated carbon and lime on soil pH and available cadmium. *J. Soil Water Conserv.*, **26** (6), 199, **2012**.
37. HU F., ZHONG J. Jasmonic acid mediates gene transcription of ginsenoside biosynthesis in cell cultures of *Panax notoginseng* treated with chemically synthesized 2-hydroxyethyl jasmonate. *Process Biochem.*, **43** (1), 113, **2008**.
38. SONG Z.G., XU M.G., LI J.M., JU X.H., TANG S.R. Effects of calcium on cadmium bioavailability in lateritic red soil and related mechanisms. *Chin. J. Appl. Ecol.*, **20**, 1705, **2009**.
39. VOEGELIN A., VULAVA V.N. Reaction-base model describing competitive sorption and transport of Cd, Zn, and Ni in acidic soil. *Environ. Sci. Technol.*, **35**, 1651, **2001**.
40. ABD-ALLAH E.F., HASHEM A., ALQARAWI A.A., WIRTH S., EGAMBERDIEVA D. Calcium application enhances growth and alleviates the damaging effects induced by Cd stress in sesame (*Sesamum indicum* L.). *J. Plant Interact.*, **12** (1), 237, **2017**.
41. WANG C.Q., WANG B.S. Ca<sup>2+</sup>-calmodulin is involved in betacyanin accumulation induced by dark in C3 halophyte *Suaeda salsa*. *J. Integr. Plant Biol.*, **49**, 1378, **2007**.
42. CHEN G.L., JIA K.Z. Effects of calcium and calmodulin antagonist on antioxidant systems of eggplant seedlings under high temperature stress. *Sci. Agric. Sin.*, **38**, 197, **2005**.
43. LIANG X.H., LUAN W.J., LIANG J., ZHANG F.X. RT-PCR Analysis on the differential expression of the key enzyme genes involved in glycyrrhizic acid synthetic metabolic pathway under B, Mn, Zn and Mo elements treatment. *Lishizhen Med. Mater. Med. Res.*, **22** (10), 2351, **2011**.
44. ZHU Y.X., HUANG Y.F., KE F. Study on correlation between content of polyphyllins and heavy metals in *Paris Polyphylla* var. *Yunnanensis*. *Eval. Anal. Drug-Use Hosp. Chin.*, **17** (10), 1315, **2017**.
45. SUN Y., ZHAO H.W., GE F., SHI L., LIU D.Q. The construction of over-expression vector for *Panax notoginseng* SS gene and its transformation. *Acta Pharm. Sin.*, **48** (1), 138, **2013**.
46. ZU Y.Q., LI Z.R., MEI X.Y., WU J., CHENG S.C., JIANG Y.Q., LI Y. Transcriptome analysis of main roots of *Panax notoginseng* identifies genes involved in saponin biosynthesis under arsenic stress. *Plant Gene*, **16**, 1, **2018**.
47. WANG M.L., FAN Q.J., LI Y.B., QU Z.C., CUI X.S. Research progress on mevalonate kinase genes. *J. Agric. Sci. Technol.*, **13** (3), 17, **2011**.
48. WANG P.J., CHEN D., CAO H.L., YAO X.Q., CHEN D., YANG G.Y., ZHENG Z.L., YE N.X. Cloning and expression of mevalonate kinase gene *CsMVK* in tea plant. *J. Southern Agric.*, **48** (10), 1741, **2017**.
49. HEDLEY M., MCLAUGHLIN M. Reactions of phosphate fertilizers and by-products in soils. In *Phosphorus:*



- Agriculture and the Environment. Editor Sims J.T., Sharpley A.N., Wiley, USA, **46**, 181, **2008**.
50. GAO Y.J., KANG L.F., CHU G.X. Polymerization degree and rate of polyphosphate fertilizer affected the availability of phosphorus, Fe, Mn and Zn in calcareous soil. *J. Plant Nutr. Fert.*, **24** (5), 1294, **2018**.
51. PAN L.X., ZHU J., WANG Q.Y., YANG D.F. Cytochromes P450 for the antibiotic biosynthesis in *Streptomyces*. *Chin. J. Antibiotics.*, **44** (2), 153, **2019**.
52. XU J.S., WEI J.H., TAO Y.W., SUN J., SUI C. Advances in studies on functions of plant cytochrome P450 in triterpenoid saponin biosynthesis. *Chin. Tradit. Herbal Drugs.*, **43** (8), 1635, **2012**.
53. LIU L.J., ZHANG X.M., SU Z.M., YI Y. Physiological and biochemical characteristics of plant iron nutrition in calcareous soil: research progress. *Chin. Agric. Sci. Bull.*, **34** (11), 6, **2018**.
54. LONG Y.H., LI F.F., YANG G., XIN C.B. Gene cloning and expression level of  $\beta$ -AS and their correlation with content of saponins in *Eleutherococcus senticosus*. *Chin. Tradit. Herbal Drugs.*, **46** (9), 1354, **2015**.
55. JIN M.L., LEE D.Y., UM Y., LEE J.H., PARK C.G. Isolation and characterization of an oxidosqualene cyclase gene encoding a beta-amyrin synthase involved in *Polygala tenuifolia* Willd. saponin biosynthesis. *Plant Cell Rep.*, **33** (3), 511, **2014**.
56. AUGUSTIN J.M., DROK S., SHINODA T., SANMIYA K., NIELSEN J.K., KHAKIMOV B., OLSEN C.E., HANSEN E.H., KUZINA V., EKSTRØM C.T., HAUSER T., BAK S. UDP-glycosyltransferases from the UGT3C subfamily in *Barbarea vulgaris* catalyze saponin 3-O-glucosylation in saponin-mediated insect resistance. *Plant Physiol.*, **160** (4), 1881, **2012**.
57. ZHANG Y.X., JIN L., FENG S.S., LIU J.G., CAI T.Y. Effects of Cd on activity and gene expression of antioxidant enzymes in hyperaccumulator *Solanum nigrum*. *J. Graduate. Univ. Chin. Acad. Sci.*, **30** (1), 11, **2013**.
58. ZHU M.L., ZENG X.C., JIANG Y.X., FAN X.T., CHAO S.H., CAO H.B., ZHANG W.S. Effect on the expression of key enzymes DS and P450 genes in the medicinal composition synthesis of *Panax notoginseng* under cadmium stress. *Chin. Tradit. Herbal Drugs.*, **47** (24), 4428, **2016**.