

vegetation [23]. It influences incident solar radiation, which in turn alters soil and air temperature, and soil moisture [24]. Preference for slope aspect as one of the indication of drought resistance might be useful in predicting response of the perennial species to drought. Kimball et al. [23] reported that perennial species occupying south facing slopes under Mediterranean climate had better performance under drought. The aspect preference of the *D. hastata* indicates both possible existence of variation in drought resistance and potentially presence of drought resistant genotypes. Therefore, *D. hastata* genotypes especially growing on steppe habitats with south aspects might possess higher drought resistance and be used for development of cultivars with better drought resistance.

It was found that *D. hastata* individuals occupied different habitat structures ranging from forests especially at lower altitudes to the steppe regions with limestone slopes and calcareous rocky habitat structures at the higher altitudes within the distribution regions. The 75% of the populations existed under or openings of forest dominated by *Pinus brutia* trees accompanied with rocky habitat structures (Table 1). When the distribution regions of *D. hastata* is evaluated with a floristic point of view, it is observed that the dune, maquis and forest characters dominate the coastal part of the research area including Antalya-Kumluca route. These floral elements and their dominance gradually decrease with the gradual ascent towards the upper altitudes. In the higher elevation, the sand dunes turn into maquis and forest accompanied with rocky habitat structures. The trees end at about 1800 m, leaving to the limestone slopes and stony parts as main habitats that are among the most important topographic and vegetative elements in the region.

Determination of the amount and distribution of genetic variation within and among populations of a given species might provide important basic information for breeding programs and for the establishment of programs to conserve genetic resources [25]. Analysis of variance revealed the existence of significant variation ($P < 0.0001$) both within and among populations (Table 2). Both vegetative and generative plant structures differed substantially among populations. Within variation observed in some populations was also noteworthy. The most morphologically diverse genotypes were in Gulluk and Ucoluk, followed by Hisarcandir and Alakir populations. For instance, the genotypes within Gulluk population, varied in almost all morphological characters (Table 2). On the contrary, genotypes in Sogutcumasi and Altinyaka populations differed from each other only for four and five of the 15 morphological characteristics, respectively. Plant height and diameter along with flower spike length and width showed the highest variations among genotypes within populations. Although some of the morphological variation observed among populations might be attributed to the environmental differences

Table 2. Analysis of variance between and within populations for morphological characteristics of *D. hastata* genotypes. (PH: Plant height (cm), PD: Plant Diameter (cm), LW: Leaf width (Leaf width measured from the widest part of the leaves (cm)), LL: Leaf length (Leaf length measured from the longest part of the leaves (cm)), PETL: Petiole length (cm), SW: Spike width (Spike width measured from the widest part of the spike (cm)), SL: Spike length (cm), PEDL: Peduncle length (cm), CW: Calyx width (mm), CL: Calyx length (mm), COW: Corolla width (mm), COL: Corolla length (mm), HUE: Leaf color)

Source of variation	df	Mean square and P values												
		PH	PD	LW	LL	PETL	SW	SL	PEDL	CW	CL	COW	COL	Hue
Between populations	15	1560.5	36849.7	6.02	17.39	4.57	0.53	75.068	5.79	1.2	3.13	7.57	5.38	40.516
P value		<.0001	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001
Error		571.9	2281	0.27	0.76	0.32	0.016	2.249	0.679	0.08	0.17	0.31	0.34	3.555
Within populations														
Tahtalı	3	6498.4	24198.4	0.594	2.458	0.457	0.059	10.58	1.18	0.058	0.46	4.23	0.06	9.189
		<.0001	0.004	0.022	<.0001	0.107	<.0001	<.0001	0.356	0.1751	0.0002	<.0001	0.238	<.0001
Kesmeboğazi	2	551.4	7100.0	0.595	3.374	0.51	0.035	1.069	0.72	0.02	0.31	1.49	0.95	18.613
		0.017	<.0001	0.198	0.026	0.418	0.49	0.87	0.253	0.845	0.262	<.0001	0.0108	0.0064

Table 2. Continued.

Gölcük	3	1266.7	6968.2	0.514	2.731	0.592	0.056	14.426	2.277	0.09	1.06	1.02	0.53	3.637
		0.016	0.008	<.0001	<.0001	0.0035	0.0192	<.0001	0.188	0.394	0.006	0.071	0.542	0.504
Beldibi	4	3420.8	39953.9	0.467	1.834	0.497	0.044	11.921	0.808	3.53	2.76	3.04	2.71	7.421
		0.0065	<.0001	0.004	<.0001	0.066	<.0001	<.0001	0.072	<.0001	<.0001	<.0001	<.0001	0.48
Beycik	3	1603	14110.9	0.989	2.123	1.811	0.088	48.224	3.76	0.16	0.57	2.35	3.18	12.44
		0.034	<.0001	0.366	0.476	0.015	0.0016	<.0001	0.002	0.0067	0.056	<.0001	<.0001	0.074
Altmyaka	2	204.2	35654.2	0.236	1.115	0.671	0.011	5.458	1.196	0.04	0.19	0.46	0.25	2.029
		0.037	0.0013	0.607	0.03	0.004	0.683	0.0006	0.216	0.289	0.144	0.225	0.637	0.804
Gulluk	3	927.3	19233.4	1.64	6.937	1.551	0.222	46.292	1.119	0.22	0.54	1.81	3.84	5.799
		0.001	0.0094	0.0017	<.0001	<.0001	<.0001	<.0001	0.473	<.0001	0.001	<.0001	<.0001	0.05
Hisarçandır	4	2492.4	1619.7	1.657	3.962	0.615	0.12	7.806	0.64	0.26	0.61	0.99	0.4	21.861
		0.0003	0.0004	<.0001	<.0001	0.003	<.0001	<.0001	0.07	0.603	0.0007	0.075	0.0374	0.0002
Üçoluk	3	757.5	12845.3	0.921	1.867	1.444	0.113	4.968	2.177	0.26	0.09	0.77	0.55	21.249
		0.21	0.002	<.0001	0.023	<.0001	<.0001	0.0011	0.0024	0.546	0.411	0.019	0.019	<.0001
Söğütçüması	2	678.7	14969.7	0.385	1.553	0.941	0.03	4.331	0.22	0.04	0.4	0.35	1.15	1.223
		0.02	0.015	0.447	0.062	0.176	0.237	0.209	0.211	0.157	0.036	0.548	0.088	<.0001
Sivridağ	3	951.3	1315.0	0.611	2.069	1.148	0.046	7.14	0.238	0.06	0.78	0.59	0.69	7.848
		0.0013	0.002	0.064	0.093	0.385	0.003	0.0001	0.006	0.002	<.0001	0.073	0.182	0.019
Fesitikan	3	438.8	325.9	0.352	3.016	0.206	0.064	3.557	0.845	0.04	0.79	1.52	0.9	4.812
		<.0001	<.0001	0.337	0.002	0.571	0.0002	0.122	0.016	0.099	0.019	0.0015	0.252	0.222
Tünektepe	2	441.7	8541.	0.313	2.656	0.219	0.023	25.893	0.457	0.1	0.2	0.31	1.1	5.27
		0.03	0.05	0.016	0.003	0.79	0.123	<.0001	0.837	0.082	0.309	0.709	0.033	0.46
Alakır	3	2530.0	18300.9	0.129	2.01	0.527	0.121	13.361	3.205	0.13	0.37	1.07	1.33	7.998
		0.007	0.0014	0.263	<.0001	0.0019	<.0001	<.0001	0.0094	0.222	0.0669	0.0036	0.0001	0.0105
Göynük	2	635.7	13850.2	0.121	0.73	0.304	0.061	2.016	0.981	0.02	0.23	2.39	1.69	11.156
		0.003	0.0008	0.378	0.058	0.172	<.0001	0.221	0.075	0.897	0.0169	0.0002	<.0001	0.0284
Hacısekiler	2	679.2	73887.5	0.967	2.88	0.395	0.012	3.606	0.13	0.04	0.14	1.34	0.75	5.676
		0.03	<.0001	0.0404	0.056	0.026	0.254	0.395	0.477	0.165	0.475	<.0001	0.0006	0.0243

Table 3. Averages and standard deviations of morphological characteristics

(*PH: Plant height (cm) PD: Plant diameter (cm), LW: Leaf width (L-leaf width measured from the widest part of the leaves (cm)), LL: Leaf length (L-leaf length measured from the longest part of the leaves (cm)), PETL: Petiole length (cm), SW: Spike width (Spike width measured from the widest part of the spike (cm)), SL: Spike length (cm), PEDL: Peduncle length (cm), CW: Calyx width (mm), CL: Calyx length (mm), COW: Corolla width (mm), COL: Corolla length (mm), HUE: Leaf color of *D. hastata* populations, Antalya-Turkey.

Population	PH	PD	LW*	LL	PETL	SW	SL	PEDL	CW	CL	COW	COL	HUE
Tahtali	91.9±54.8	194.4±109	2.16±0.59	4.50±1.03	1.86±0.57	1.02±0.17	7.44±2.09	1.67±1.03	2.00±0.21	3.69±0.45	3.70±1.21	5.34±0.70	118.279±2.052
Kesmebogazi	58.5±17.3	133.3±59.5	2.81±0.68	6.26±1.43	2.60±0.69	1.02±0.19	7.00±1.31	1.78±0.77	2.01±0.21	4.02±0.51	3.21±0.83	4.83±0.73	118.931±3.185
Golcuk	67.1±25.9	216.5±59.6	3.12±0.46	5.95±1.09	2.56±0.55	1.35±0.18	10.90±2.40	2.60±1.33	2.20±0.30	4.44±0.75	4.57±0.82	5.76±0.75	117.876±1.932
Beldibi	62.5±41.6	338.7±132.8	2.48±0.49	5.33±0.85	2.09±0.57	0.83±0.14	8.00±1.98	1.55±0.73	2.38±1.01	4.14±0.93	2.81±0.96	4.84±0.94	119.227±2.161
Beycik	58.5±29.9	251.9±80.7	4.16±0.95	7.86±1.46	3.54±1.01	1.38±0.21	14.90±4.45	2.50±1.36	2.26±0.30	4.71±0.61	4.74±0.96	5.83±1.10	118.866±2.574
Altinyaka	42.5±10.7	310.0±134.9	2.65±0.51	4.77±0.82	2.11±0.60	1.03±0.12	8.85±1.63	2.29±0.97	2.12±0.19	3.67±0.38	4.22±0.61	4.84±0.54	117.728±1.693
Gulluk	64.4±21.1	209.4±99.1	3.75±0.89	6.41±1.69	2.64±0.79	1.20±0.28	10.74±4.06	2.61±1.05	2.13±0.31	3.81±0.51	3.53±0.81	5.17±1.14	120.647±1.935
Hisarcandir	73.2±33.8	116.8±27.3	2.81±0.78	5.37±1.14	1.90±0.55	1.07±0.21	9.66±1.80	1.38±0.65	2.02±0.55	3.56±0.53	3.41±0.82	5.25±0.61	119.957±3.077
Ücoluk	51.1±23.3	181.3±79.1	1.58±0.62	3.62±1.04	1.26±0.76	0.75±0.21	7.83±1.53	1.51±1.04	1.71±0.53	3.22±0.31	2.82±0.67	4.54±0.56	119.823±2.91
Sogutcumasi	64.8±18.4	183.2±89.8	1.91±0.61	4.24±1.01	1.71±0.85	0.82±0.16	8.70±1.85	1.19±0.42	1.78±0.18	3.78±0.52	2.94±0.62	5.26±0.89	120.114±0.681
Sivri dag	51.3±21.4	82.5±25.8	1.87±0.63	4.18±1.19	1.64±1.03	0.80±0.15	7.67±1.77	1.08±0.35	1.87±0.17	3.68±0.57	3.00±0.62	4.88±0.07	121.8±2.125
Feslikan	51.3±14.2	60.0±12.2	2.38±0.56	4.49±1.22	1.78±0.47	0.97±0.17	7.10±1.60	1.85±0.69	2.00±0.18	3.89±0.68	3.30±0.86	4.88±0.87	116.794±1.961
Tunektepe	52.5±15.7	248.3±70.5	2.25±0.42	5.23±1.18	1.78±0.55	0.92±0.13	9.85±3.31	1.53±0.83	1.85±0.26	4.06±0.43	3.76±0.63	5.37±0.86	116.49±2.275
Alakir	75.0±35.7	196.9±94.1	2.82±0.33	4.73±0.88	2.30±0.51	1.05±0.21	9.85±2.22	2.88±1.32	2.17±0.33	3.73±0.49	3.89±0.74	4.84±0.76	117.424±2.09
Goyruk	58.0±18.1	226.0±83.8	2.12±0.33	4.20±0.69	1.71±0.48	0.80±0.17	7.78±1.27	1.30±0.81	1.80±0.19	3.14±0.37	3.91±1.06	4.37±0.87	115.834±2.601
Hactisekiler	57.5±19.4	246.7±190	2.50±0.78	4.96±1.37	1.65±0.49	0.82±0.10	9.91±1.83	0.86±0.36	1.96±0.18	4.06±0.39	3.00±0.76	6.08±0.61	119.437±1.843
All populations	62.03±29.1	198.2±111	2.61±0.91	5.16±1.55	2.08±0.86	1.00±0.26	9.17±3.08	1.81±1.09	2.03±0.45	3.85±0.68	3.54±1.01	5.12±0.91	118.735±2.682
Max.	185.0	620.0	6.2	10.6	6.369	2.089	26.2	8.315	5.05	6.26	6.25	7.48	126.246
Min.	14.0	40.0	0.43	1.4	0.25	0.438	3.054	0.272	1	2	1.52	2.32	109.281
LSD (0.05)	33.1	130.9	0.68	1.37	0.69	0.23	3.29	0.82	0.53	0.69	1.09	1.01	2.41

Table 4. Analysis of variance between and within populations for edaphic characters of *D. hastata*. (L: Lime, EC: Salinity, TX: Soil texture (<%30 = Sand, %30-%50 = Loam, %50-%70 = Loamy clay, %70-%110 = Clay, %110< = Heavy clay), OM: organic matter, P: Phosphorous (ppm), K: Potassium (ppm), Mg: Exchangeable magnesium (ppm), Fe: Available iron, Mn: Available manganese, Zn: Available zinc and *df: degrees of freedom)

Source of variation	df	Mean square and P values											
		pH	L* (%)	EC (%)	TX (%)	OM (%)	P (ppm)	K (ppm)	Mg (ppm)	Fe (ppm)	Mn (ppm)	Zn (ppm)	
Between Population	15	0.081	388.611	0.018	861.26	2.008	31.32	10853.129	118843.687	221.92	101.112	12.572	
P value		0.016	0.337	0.102	0.084	0.044	<.0001	0.0033	0.0084	0.322	0.205	0.0214	
Error		0.036	335.593	0.011	514.38	1.061	7.223	3919.179	48450.684	188.608	74.71	5.877	
Within Population		Average and standard deviations											
Tahtali		8.00±0.21	8.178±7.84	0.46±0.11	99.05±22.93	4.95±0.24	9.75±4.91	425.42±232.73	333.15±139.14	18.92±35.24	13.58±5.78	2.25±2.17	
Kesmebogazi		7.47±0.06	35.37±30.94	0.223±0.06	51.63±14.23	2.52±1.22	2.27±1.07	286.68±199.67	303.54±254.26	5.84±4.16	10.31±5.77	0.39±0.07	
Goleuk		7.77±0.09	1.96±0.90	0.35±0.05	77.00±9.70	4.60±0.31	14.72±2.46	197.75±86.88	648.65±601.24	28.64±30.03	18.28±14.50	0.59±0.25	
Beldibi		7.55±0.05	24.53±2.76	0.26±0.16	59.29±3.44	3.70±0.81	3.04±2.69	123.37±38.94	217.87±11.78	8.02±5.98	4.86±0.89	0.49±0.056	
Beycik		7.8±0.29	24.81±18.47	0.37±0.13	80.02±26.11	4.00±1.36	9.59±6.88	315.17±218.28	406.77±165.70	0.88±0.48	7.45±2.85	0.69±0.39	
Altinyaka		7.85±0.05	41.92±40.16	0.435±0.035	95.7±7.26	4.95±0.25	17.36±2.34	436.07±114.58	371.85±42.85	1.32±0.06	6.50±1.13	1.90±1.03	
Gulluk		7.85±0.19	27.66±19.20	0.39±0.07	86.68±17.47	4.86±0.42	18.85±0.63	395.31±225.16	211.25±61.07	6.70±10.19	16.20±14.87	4.03±2.95	
Hisarcandir		7.68±0.19	17.82±15.35	0.32±0.10	71.28±22.13	3.31±1.29	5.80±4.42	456.17±301.06	413.06±159.45	8.65±5.85	17.27±8.96	0.53±0.41	
Ücoluk		7.80±0.08	10.38±18.09	0.35±0.03	78.54±7.37	4.95±0.18	10.95±6.27	356.89±144.96	224.14±91.11	9.88±11.14	17.19±6.24	1.77±1.36	
Sogutcumasi		7.6±0.17	2.62±0.41	0.27±0.084	62.41±19.57	3.59±1.64	16.19±2.64	439.24±175.90	268.00±127.92	7.143±6.06	15.81±12.74	1.12±0.55	
Sivridag		7.65±0.10	24.61±28.32	0.31±0.07	67.02±11.58	4.34±1.06	13.97±5.28	249.82±136.89	127.73±48.66	8.96±7.73	11.26±6.07	0.98±0.59	
Feslikan		7.97±0.21	9.74±9.09	0.48±0.13	102.87±25.70	4.84±0.32	13.82±5.89	706.12±168.53	574.27±342.78	15.88±6.53	20.72±12.73	5.96±5.54	
Tunektepe		7.86±0.21	4.59±3.39	0.39±0.08	85.32±16.79	4.88±0.33	9.96±8.16	369.69±155.83	363.56±29.48	1.17±1.03	6.06±1.75	4.00±5.02	
Alakir		7.73±0.32	17.79±20.12	0.37±0.21	81.78±43.28	3.99±1.63	12.02±5.92	242.15±68.33	313.8±221.09	11.79±16.74	3.86±3.90	1.49±1.446	
Goyruk		7.6±0.26	14.11±3.77	0.27±0.15	61.23±30.69	2.75±2.07	0.73±0.61	166.15±110.21	811.63±268.36	1.89±0.67	11.09±11.08	0.77±0.93	
Hacisekiler		8.03±0.21	12.61±17.68	0.46±0.11	112.22±40.18	5.11±0.07	18.60±3.20	921.42±505.45	509.50±348.82	4.64±4.91	16.01±6.15	7.06±4.85	
Pop Mean		7.76±0.22	16.88±18.79	0.35±0.11	79.22±25.22	4.21±1.18	10.92±6.74	374.98±261.55	376.00±270.5	9.42±14.32	12.6±9.19	2.06±2.87	
Min		7.4	0.8	0.15	35.2	1.48	0.19	59.14	55.29	0.35	1.02	0.14	
Max		8.2	82.08	0.63	155.39	5.23	20.51	1299.17	1545	71.77	39.6	13.1	

Table 5. Correlation coefficients among some morphological and ecological characteristics of *D. hastata* genotypes (AIT: Altitude, PD: Plant diameter, LW: Leaf width, LL: Leaf length, PETL: Petiole length, SW: Spike width, SL: Spike length, PEDL: Peduncle length, CW: Calyx width, CL: Calyx length, COW: Corolla width, COL: Corolla length).

Characters	ALT	LW	LL	PETL	SW	SL	PEDL	CW	CL	COW
PD	-0.42**									
LL	-0.20*	0.88**								
PETL	-0.15*	0.82**	0.81**							
SW	ns	0.74**	0.65**	0.69**						
SL	ns	0.63**	0.68**	0.61**	0.55**					
PEDL	ns	0.62**	0.50**	0.58**	0.62**	0.47**				
CW	ns	0.29*	ns	0.32*	0.35**	0.27*	0.26*			
CL	ns	0.37**	0.40**	0.44**	0.52**	0.37**	ns	0.66**		
COW	ns	0.36**	0.29*	0.47**	0.60**	0.36**	0.41**	ns	0.34**	
COL	ns	ns	0.28*	ns	0.37**	0.44**	ns	ns	0.45**	0.41**

*, ** and ns indicating significance at probability of 0.05 and 0.01, and nonsignificant at probability of 0.05, respectively.

two components of PCA respectively explained 83.3 % and 9.1% of the total variation.

Using population means, a UPGMA dendrogram was also produced by the similarity index (Fig. 3). The 16

populations were grouped into five clusters, containing 1 to 8 populations. The first group divided into two subgroups and Sivridag population was separated from the other 9 populations. Feslikan, Sivridag, and




Fig. 1. Unweighted pair group method arithmetic average (UPGMA) dendrogram with similarity coefficients of *D. hastata* in genotype level. Data were based on the means of morphological and edaphic characters of *D. hastata* genotypes.

(Genotype codes: HC1: Hacisekiler1, HC2: Hacisekiler2, HC3: Hacisekiler3, S1: Sivridag1, S2: Sivridag2, S3: Sivridag3, F1: Feslikan1, F2: Feslikan2, F3: Feslikan3, F4: Feslikan14, TN1: Tunektepe1, TN2: Tunektepe2, TN3: Tunektepe3, H1: Hisarcandir1, H2: Hisarcandir2, H3: Hisarcandir3, H4: Hisarcandir4, H5: Hisarcandir5, UC1: Ucoluk1, UC2: Ucoluk1,2, UC3: Ucoluk3, UC4: Ucoluk4, BL1: Beldibi1, BL2: Beldibi2, BL3: Beldibi3, BL4: Beldibi4, BL5: Beldibi5, GY1: Goynuk1, GY2: Goynuk2, GY3: Goynuk3, K1: Kesmebogazi1, K2: Kesmebogazi2, K3: Kesmebogazi3, BY1: Beycik1, BY2: Beycik2, BY3: Beycik3, BY4: Beycik4, T1: Tahtali1, T2: Tahtali2, T3: Tahtali3, T4: Tahtali4, GL1: Golcuk1, GL2: Golcuk2, GL3: Golcuk3, GL4: Golcuk4, A1: Altinyaka1, A2: Altinyaka2, A3: Altinyaka3, SO1: Sogutcumasi1, SO2: Sogutcumasi2, SO3: Sogutcumasi3, AL1: Alakir1, AL2: Alakir2, AL3: Alakir3, AL4: Alakir4, GU1: Gulluk1, GU2: Gulluk2, GU3: Gulluk3, GU4: Gulluk4)




Fig. 3. Unweighted pair group method arithmetic average (UPGMA) dendrogram with similarity coefficients of *D. hastata* in population level. Data were based on the means of morphological and edaphic characters of *D. hastata* populations.

(Population codes: HC: Hacisekiler, S: Sivridag, F: Feslikan, TN: Tunektepe, H: Hisarcandir, UC: Ucoluk, BL: Beldibi, GY: Goynuk, K: Kesmebogazi, BY: Beycik, T: Tahtali, GL: Golcuk, A: Altinyaka, SO: Sogutcumasi, AL: Alakir, GU: Gulluk)




Fig. 4. Plot of principal components based on the means of morphological and edaphic characteristics of *D. hastata* populations.

(Population codes: HC: Hacisekiler, S: Sivridag, F: Feslikan, TN: Tunektepe, H: Hisarcandir, UC: Ucoluk, BL: Beldibi, GY: Goynuk, K: Kesmebogazi, BY: Beycik, T: Tahtali, GL: Golcuk, A: Altinyaka, SO: Sogutcumasi, AL: Alakir, GU: Gulluk)

more diverse and sustainable landscapes [35]. In this regard, the native species and genotypes with an ornamental plant potential need to be determined first, and then cultivated, improved through breeding and used at suitable ecologies. Although *D. hastata* is a local endemic species, it has been able to maintain its diversity with a wide morphological variation including growth habit ranging from short creeping types that can be used as ground cover to taller shrub formations. Some of the genotypes (such as from

Beycik) with their large and numerous flashy flower spikes and lush green vegetation have a unique aesthetic appearance that should be exploited in ornamental plant industry. The variations offer opportunities to breeders to introduce new forms to ornamental plant sector. Moreover, the future presence of the species is under risk of extinction because of uncontrolled mass collection due to pharmacological properties in addition to grazing pressure and housing constructions. Hence, the natural populations are progressively decreasing

