

factors. Internal causes are mostly associated with the nature of the stand development and mass reproduction of insects. The mass reproduction of insects (*xylophages* and *phyllophages*) leads to large-scale but fragmentary damage to forest areas [2]. The insect outbreak manifests as spots, which look like areas of dry trees, or areas with fallen trunks. A characteristic feature of insects' mass reproduction is the cyclicity of forest damage, associated with climatic conditions of a region and developmental stages of insects. Moreover, insects' outbreaks are presented as periodic travelling waves occurring in different regions or even continents [3].

A bark beetle (*Ips typographus* Linnaeus) is the most dangerous pest for spruce (*Picea abies* Linnaeus) trees [4]. While the major habitat of the bark beetle is spruce, it may also attack cedar, larch and other conifers, yet the damage may be less than with the spruce. Dry periods, which seem to extend each year, favor the breeding of the typographer beetle. Specifically, in the summer of 2010, in the European part of Russia (Moscow region), abnormally high temperatures caused the drying of some spruce forests and the development of a food supply suitable for bark beetles of the genus typographer [5]. Thus, a great amount of dry wood contributes to developing the second generation of insects, which led to their mass reproduction. The bark beetles of the genus typographer affect mainly old spruce trees of 70-80 years [6]. Breeding outbreaks occur regularly and have been reported since the 19th century [7]. The typographer beetle aside, the subfamily *Scolytinae* consists of 140 species that can be found in Europe, most of which feed on conifers.

The quality and viability of spruce stands depend not only on insects' mass reproduction but also moisture regime. The diameter of the spruce tree is directly related to the mass of needles [8]. According to various sources, about 20-30% of precipitation is retained in the crown and does not reach soil due to the dense crown [9]. A significant lack in soil moisture can be expected in sandy soils since water does not keep in the surface layer. The active growing season of spruce occurs from May to September; therefore, the amount of precipitation at this time is critical. According to studies, a lack of moisture should be expected in spruce trees older than 35 years, due to the difference between the moisture consumption for transpiration and the amount of moisture coming from the soil [10].

Recently, a progressive drying of spruce forests and stands occurred in Europe [11]. (DeMarco et al. 2017). This is primarily due to global warming, particularly, a decrease in precipitation. The decrease in precipitation causes more outbreaks of insects, affecting more and more areas of spruce forests and stands. Therefore, the method for monitoring spruce stands using modern technologies should be introduced, namely, the development of interactive spruce stands maps using geographic information system (GIS technologies) and unmanned aerial vehicles (UAVs) [12-15].

Generally, the UAVs have been previously used for exploration, monitoring of oil and gas pipelines, and identifying foci of forest fires. The cost of UAVs production was significantly reduced due to technical progress, thus it possible to use them in monitoring the ecosystems, including forest. The UAVs are commonly used in the EU countries, the USA, Japan and China to search forests for lost people, track the extent of forest fires, and detect large gatherings of insect pests. In China, drones are used in agriculture for irrigation purposes.

The study attempts to develop interactive spruce stands map of the Moscow region using data obtained with UAVs and GIS technologies. The method helps to carry out a full-fledged, express assessment of spruce stands, covering large areas. Moreover, the data from different years allows us to assess the dynamics of the succession or degradation processes occurring in spruce stands over the decades. Despite the great forest cover of the territory (from 40 to 50% on average), the Moscow region is a convenient model territory due to intensive development and the felling of spruce stands. The data obtained can be used to study the influence of the territory expand to a megalopolis (Moscow) on forest ecosystems and their structure, on the example of spruce stands.

The aim of the study is to develop interactive spruce stands map of two model landscapes of the Moscow region, considering the foci of insects' mass reproduction. The tasks of the research included: a) development of interactive spruce stands map and related tree species using GIS technologies; b) study the dynamics of succession and degradation changes in spruce stands of the Moscow region (Russia) during 1991-2016; c) development of a model map of the anthropogenic landscape with a predominance of spruce stands using UAV data.

Experimental Procedure

Study Location

The Klinsko-Dmitrovskaya Gryada has an elevation of 290-310 m above sea level. Gryada is made up of Cretaceous rocks, which are overlapped by rocks of the Quaternary age. Since, gryada was several times affected by repeated glaciations (the last occurred about 170-125 thousand years ago) its landscape characteristics are associated with moraines, determining the structural and mechanical composition of soils. The Istrinsky landscape is located in the lowlands of the gryada, at altitudes up to 170 m above sea level. The Istrinsky district is made up of outcrops of Jurassic clays, thus, there is high humidity. The moraine was eroded by the melting of the glacier. This landscape is characterized by intensive development, gardening, development of rural power, and deforestation.




Fig. 1. Dynamics of areas reduction occupied by forests in 1991-2016 in the Istrinsky a) and Luzganinsky landscapes b).

Results

GIS Mapping

A predominance of spruce and pine-spruce forests is observed in the Istrinsky landscape (Fig. 1a), which make up background plant communities. Spruce and pine-spruce forests draw up to 55% of all forests in the landscape. Particularly, spruce forests make up about 40% while spruce-pine forests about 15%.

Forests grow on depressions, mainly on the flat areas of moraine and water-glacial areas. They grow at altitudes of 180-220 m above sea level. Meanwhile, short-term productivity forests (61%) and conditional inference forests (5%) prevail among the plant communities in this area.

The high-resolution images helped to identify foci of spruce affected by bark beetles of the genus *typographer*. The lesions are located in the northern part of the ridge and occur on monodominant trees aged 100 years. Therefore, local forestry carried out clear-felling, which negatively affected the sustainability of forest communities. An analysis of the maps showed an increase in the fragmentation of forests and spruce stands.

The areas covered by a prime denote the forest area that has decreased from 1991 to 2016. The main figure is taken from satellite imagery data in natural colors. The figures indicate the types of plant communities:

figure – conditional inference forests; fractional – short-term productivity, with the dominance of spruce and pine; a figure with the letter – long-term productivity with the dominance of small-leaved species.

Coniferous (spruce) forests prevail in the Luzgarinsky landscape (Fig. 1b), making up 87% of the total forest area. The share of the Luzgarinsky landscape is about 70%. The average age of spruce trees is 90-100 years. The conditional inference forests (15%) and short-term productivity forests (70%) prevail. Fragmentation is not expressed, since forests are located densely. Forests grow in swampy lowlands, having the full the water balance in ecosystems.

The analysis of maps from 1991 to 2016 showed significant differences in a forest area in the Istrinsky landscape and insignificant in Luzgarinsky (Table 1).

According to Table 1, the forest area in the Istrinsky landscape has decreased by one-fifth, while changes in the Luzgarinsky are insignificant. Thus, the decrease in forest area has occurred in both landscapes.

Table 1. Change in forest cover in the Istrinsky and Luzgarinsky landscapes during 1991-2016 (in %).

| Landscapes | 1991 year | 2016 year |
|-------------|-----------|-----------|
| Istrinsky | 53.1 | 44.3 |
| Luzgarinsky | 74.5 | 70.1 |

of spruce forests and stands drying have become more frequent in many European countries [3, 25]. No cases of mass drying of spruce were observed, except forest areas affected by bark beetles of the genus typographer. The drying is generally associated with recent abnormal summer temperatures. With the mass death of trees, there is a loss of food supply for many invertebrates [26], namely, aphids and ants [27]. Considering the role of these insects, a decrease in their number contributes to further degradation of forest ecosystems. Moreover, according to some data, single-species tree stands also contain depleted insect communities, in particular, ants, in comparison with forests [28]. With the mass death of single-species trees, there is no alternative source of food for insects. Thus, their population decreases. On the one hand, red wood ants are useless against the typographer beetle, and cannot fully extinguish outbreaks of its mass reproduction. There are cases when dry or dying trees became their nesting place, in conditions of strong shading, as was shown on the example of *Formica rufa* [29].

The lack of reproduction of the next generations is another reason for the instability of single-species man-made spruce stands. The undergrowth in monospecific tree associations is not viable, since it is not adapted for growing under the canopy of mature spruce trees [7, 30, 31]. The Luzgarinsky landscape has a high forest cover and low susceptibility to bark beetles, since includes not only spruce but other tree species – oak, pine, aspen, birch.

The reasons for deforestation of short-term and conditional inference forests in the Moscow Region may be different. Along with deforestation, sanitary felling is also possible, e.g. to prevent outbreaks of bark beetles of the genus typographer, which mainly cover old spruce trees.

The data obtained with UAV allows establishing some additional parameters: a) the number of trees in the upper layer; b) type of forest (natural, artificial); c) determination of the state of the spruce forest (how much it is affected by bark beetles of the genus typographer, how many dry-topped trees per unit area, how many trees of different species suffered as a result of strong wind blows and snow breakers); d) age and species of trees in the upper layers of the forest, as well as their seedlings and young growth in felling. The high resolution of images (up to 10 cm) allows clarifying the necessary aspects of the study in detail. The listed data are not given in this article, except for paragraph B aimed at subsequent developments. The conducted research period did not allow processing all the information obtained by the UAV. The disadvantages of UAV are errors in determining the coordinates of the center of the image, up to 10-15 m, comparing to GIS maps. This is due to the side drift of the UAV, as well as the rolls when vehicle maneuvers. The large number of images obtained is another disadvantage, which troubles information processing or takes more time. Moreover, the use of UAVs is useful in small areas. However,

permission for shooting in private property is a separate obstacle.

Conclusions

The processing of satellite images showed an unequal decrease in forest cover for the spruce forests and stands during 1991-2016, namely from 53% to 44% and from 73% to 70% in the Istrinsky and Luzgarinsky landscape, respectively. The reason is that the Istrinsky landscape is close to the metropolis (Moscow) unlike Luzgarinsky one. The total area of long-term productivity forests is twice less comparing to short-term productivity and conditional inference forests in the Istrinsky landscape. The short-term productivity and conditional inference forests decreased in area by 67% during 1991-2016, while long-term productivity - only by 28%. The area of deforestation in the Istrinsky district was 53.5 km².

The data obtained with the UAV helped to determine the area affected by the bark beetle: 3.2 out of 93.6 hectares of the total area of spruce forests. A detailed map of the surveyed area was obtained. The UAV helps to develop maps of drying trees, forest areas damaged by the strong wind blows, snow breakers, as well as dividing the study area into quarters with a different composition of tree species. These forests are at least 50 years old. UAV allows studying the state of tree undergrowth in felling areas. Forests affected by bark beetle are at least 50 years old. The use of UAVs is justified for monitoring small areas, up to 200 hectares, since the processing of images takes lots of time. Meanwhile, the accuracy of the images is at least 10 cm per 1 pixel, which exceeds the capabilities of free satellite imagery. The disadvantage of UAV is inadequate orientation in the images since there is always an error caused by the maneuvers.

Aknowledgements

The study was prepared based on research within the framework of the state-ordered research theme of the Institute of Geography RAS, no. 0148-2019-0008 (AAAA-A19-119022190170-1) 'Problems and prospects of Russia's territorial development in terms of its unevenness and global instability'.

Conflict of Interest

The authors declare no conflict of interest.

References

1. WALSHE D., MCINERNEY D., VAN DE KERCHOVE R., GOYENS C., BALAJI P., BYRNE K.A. Detecting

