



Fig. 1. Geographical locations of the rivers studied along Chile (A: northern area; B: central area; C: southern area).

activities in its basin, it is one of the most important rivers of Chile. The Maipo River is located in an area where there are different deposits in the geology of the Cajón del Maipo, with fluvial, fluvio-glacial and alluvial deposits strongly linked to the formation of terraces throughout the Cajón. The Trancura River is a watercourse that flows in the Araucanía Region, which originates in the Quillehue Lagoon and flows into the Villarica Lake. The Correntoso River is a watercourse that originates from other tributaries in the Araucanía Region, which flows into the Pucón River. The Correntoso and Trancura rivers are located in an area that represents the evolution of the South American southwest margin, from the Mesozoic to the Holocene. The Marchant River is located in the Aysén Region (300 km south of Puerto Montt) is influenced by the Melimoyu volcano, and it originates from snow and rainfall, flowing into the Pacific Ocean. The Bronce, Robalo and Akika rivers, which flows into the Beagle Channel, are located on Navarino Island, Tierra del Fuego, Southern Chile, with the absolute absence of volcanoes. The relief of Navarino Island is of subglacial origin, where the summits rarely exceed 1,000 m a.s.l., with the Dientes de Navarino peak being the highest elevation summit on the island, reaching 1,118 m a.s.l.

At all sites, superficial sediment samples (~20 cm depth) were collected in the middle part of the rivers by means of a pre-cleaned plastic shovel ($n = 3$ per site), and then stored at -20°C in clean polypropylene bags (Whirl-Pak®) until their arrival at the laboratory.

Sample Preparation and Analysis

In the laboratory, all samples were freeze-dried until dry masses were constant, and then homogenized to a fine powder using a glass mortar previously cleaned with a 2% Conrad solution (Merck) for 24 h, washed with deionized water + 1M HCl, and then rinsed with distilled water [17]. Afterwards, the samples were heated (oven at 100°C for 12 h) to remove adsorbed water prior to analysis [18]. Then, the samples were analyzed for chemical elements at the Analytical Laboratory of the Faculty of Veterinary Sciences, Universidad de Concepción (Chile), using a portable battery-operated energy dispersive X-ray fluorescence spectrometer (Thermo Scientific Niton XL3t 950 He GOLDD+). The instrument was set up with the instrument tip up on a shielded laboratory test stand, which was remotely operated. The precision and accuracy were verified by means of international reference standards such as GSS-5, GSS-7, GXR-6 and GSR-3 [18], being the precision $<2\%$ and accuracy within 1-5%.

Data were statistically analyzed by means of nonparametric statistical methodologies because of the assumptions of normality and homoscedasticity were not met. Differences between chemical element concentrations were assessed by means of the Kruskal-Wallis analysis of variance and Mann-Whitney U tests. Post hoc tests were carried out by means of Kruskal-Wallis analyses, using the critical differences of mean rank. Spearman rank correlation

Table 2. Mean concentrations \pm SD of chemical elements in sediments of rivers from central Chile. Maximum and minimum values in parentheses ($\mu\text{g/g}$, d.w.).

Elements	Maipo River	Trancura River	Correntoso River
Ba	275.62 \pm 24.98 (293.66-247.11)	225.0 \pm 26.08 (254.39-204.62)	265.35 \pm 12.77 (279.4-254.44)
Ce	Bdl	194.57 \pm 32.90 (231.49-168.34)	242.60 \pm 55.59 (302.28-192.29)
La	Bdl	213.02 \pm 18.61 (227.74-192.11)	230.93 \pm 33.30 (269.37-210.86)
Nb	4.81 \pm 0.86 (5.74-4.04)	3.07 \pm 0.56 (3.71-2.69)	Bdl
Nd	Bdl	352.02 \pm 65.40 (414.84-284.31)	414.26 \pm 13.56 (426.98 – 400.0)
Pr	Bdl	397.50 \pm 31.52 (421.8-361.88)	360.80 \pm 44.35 (410.22-324.46)
Rb	28.02 \pm 1.43 (29.08-26.4)	8.53 \pm 0.60 (9.0-7.85)	8.00 \pm 0.61 (8.62-7.4)
Th	12.70 \pm 1.69 (14.37-10.99)	8.26 \pm 0.65 (8.99-7.75)	8.44 \pm 0.65 (9.16-7.88)
Y	18.47 \pm 2.19 (20.99-17.1)	20.94 \pm 0.94 (21.57-19.86)	18.65 \pm 0.82 (19.17-17.71)
Zr	128.96 \pm 3.15 (131.79-125.57)	86.53 \pm 4.19 (91.11-82.88)	73.43 \pm 3.77 (77.35-69.82)

Bdl: below detection limit.

Table 3. Mean concentrations \pm SD of chemical elements in sediments of rivers from southern Chile. Maximum and minimum values in parentheses ($\mu\text{g/g}$, d.w.).

Elements	Marchant River	Bronce River	Ukika River	Robalo River
Ba	434.84 \pm 20.15 (457.59 -419.22)	331.83 \pm 22.83 (356.79-311.99)	231.95 \pm 68.37 (278.33-153.43)	Bdl
Ce	233.03 \pm 37.84 (276.29-206.06)	152.71 \pm 42.30 (197.79-113.89)	Bdl	Bdl
La	220.25 \pm 6.64 (226.73-213.47)	159.40 \pm 45.34 (207.49-117.41)	Bdl	Bdl
Nb	8.52 \pm 0.35 (8.87-8.17)	3.36 \pm 0.47 (3.84-2.91)	3.85 \pm 0.63 (4.47-3.22)	7.74 \pm 0.80 (8.3-6.83)
Nd	440.85 \pm 14.55 (451.15-263.67)	259.25 \pm 8.05 (267.74-250.75)	Bdl	Bdl
Pr	432.95 \pm 10.45 (444.08-423.33)	270.61 \pm 41.85 (317.95-238.51)	Bdl	Bdl
Rb	18.98 \pm 1.41 (20.58-17.87)	22.17 \pm 0.86 (23.16-21.61)	21.12 \pm 0.22 (21.35-20.91)	18.76 \pm 2.96 (22.1-16.44)
Th	12.32 \pm 0.20 (12.52-12.12)	7.19 \pm 0.79 (8.1-6.69)	7.88 \pm 2.38 (10.25-5.5)	16.68 \pm 3.05 (20.07-14.16)
Y	24.81 \pm 1.68 (26.66-23.37)	17.28 \pm 1.29 (18.32-15.83)	16.35 \pm 1.23 (17.5-15.05)	18.76 \pm 2.96 (22.32-17.15)
Zr	163.0 \pm 18.5 (182.71-145.99)	80.57 \pm 1.55 (82.31-79.33)	80.06 \pm 3.12 (83.03-76.8)	75.17 \pm 8.30 (82.82-66.34)

Bdl: below detection limit.

- BAO Z., TAN C., WANG X., WAN D. Geochemistry of major and trace elements in sediments from inlets of the Xiangjiang and Yuanjiang River to Dongting Lake, China. *Environ. Earth Sci.* **77**, 16, **2018**.
19. WULANINGSIH T., HUMAIDA H., HARIJOKO A., WATANABE K. Major element and rare earth elements investigation of Merapi volcano, Central Java, Indonesia. *Procedia Earth Planet. Sci.* **6**, 202, **2013**.
20. MUNEMOTO T., SOLONGO T., OKUYAMA A., FUKUSHI K., YUNDEN A., BATBOLD T., NAGAO S. Rare earth element distributions in rivers and sediments from the Erdenet Cu–Mo mining area, Mongolia. *Appl. Geochemistry* **123**, 104800, **2020**.
21. TRIPODI E.E.M., RUEDA J.A.G., CÉSPEDES C.A., VEGA J.D., GÓMEZ C.C. Characterization and geostatistical modelling of contaminants and added value metals from an abandoned Cu–Au tailing dam in Taltal (Chile). *J. South Am. Earth Sci.* **93**, 183, **2019**.
22. BAYON G., TOUCANNE S., SKONIECZNY C., ANDRE L., BERMELL S., CHERON S., DENNIELOU B., ETOUBLEAU J., FRESLON N., GAUCHERY T., GERMAIN Y., JORRY S.J., MENOT G., MONIN L., PONZEVEVA E., RUET M.L., TACHIKAWA K., BARRAT J.A. Rare earth elements and neodymium isotopes in world river sediments revisited. *Geochim. Cosmochim. Acta* **170**, 17, **2015**.
23. ABDUSAMADZODA D., ABDUSHUKUROVA D.A., ZINICOVSCAIA I., DULIU O.G., VERGEL K.N. Assessment of the ecological and geochemical conditions in surface sediments of the Varzob river, Tajikistan. *Microchem. J.* **158**, 105173, **2020**.
24. BERMANEC V., FIKET Z., HRENOVIC J., KAZAZIC S., KNIEWALD G., VENTER CH., BOTHA A. Geochemical and microbiological characterization of sediments at the mouth of Río de Las Minas (Punta Arenas, Chile). *An. Inst. Patagon.* **47**, 49, **2019**.
25. ROMERO L., ALONSO H., CAMPANO P., FANFANI L., CIDU R., DADEA C., KEEGAN T., THORNTON I., FARAGO M. Arsenic enrichment in waters and sediments of the Río Loa (Second Region, Chile). *Appl. Geochemistry* **18**, 1399, **2003**.
26. LISTER T.R., FORDYCE F.M., FERREIRA A.M.P.J., EVERETT P., LAWLEY R.S. SW England rare earth elements (REE) stream sediment dataset user guide. *British Geological Survey*, 15, UK, **2020**.
27. XU Z., LIM D., CHOI J., YANG S., JUNG H. Rare earth elements in bottom sediments of major rivers around the Yellow Sea: implications for sediment provenance. *Geo-Mar. Lett.* **29**, 291, **2009**.
28. LIDE D.R. *Handbook of Chemistry and Physics*. CRC Press/Taylor and Francis, Boca Raton, FL, **2009**.
29. SALEEM M., IQBAL J., SHAH H. A study of seasonal variations and risk assessment of selected metals in sediments from Mangla Lake, Pakistan. *J. Geochem. Explor.* **125**, 144, **2013**.