

Original Research

Soil Environment Modulation by Varying Physicochemical Attributes Change the Population Dynamics of Fecal *Escherichia coli*

Chenghuan Qiao¹, Muhammad Azeem², Huadong Zhang¹,
Ling Yang¹, Shoujun Yang^{1*}

¹Yantai Institute of China Agricultural University, Yantai, Shandong, China

²Biosaline Research Laboratory, Department of Botany, University of Karachi, Karachi, Pakistan

Received: 26 April 2022

Accepted: 17 August 2022

Abstract

Present research reveals the effect of different soil environment modifications on population dynamic tendency of fecal *Escherichia coli* (*E. coli*). Various soil environments were created by varying the temperature regimes, relative water content, pH, C/N ratio and ammonium content. The result showed that population dynamics of *E. coli* with all treatments increased initially and then decreased with the passage of time. Maximum population of *E. coli* at higher temperature regime (37°C and 45°C) was observed on 8th day of incubation, whereas, at low temperature regime (0.6°C and 25°C), it reached the peak after 15 days of incubation, other treatments (soil relative water content, pH, C/N, ammonium content) reached the peak all after 8 days of incubation. At the end of the incubation period (60 days), maximum population of *E. coli* was observed in the soil treated with 0.6°C, 80% of relative soil water content, pH 7.0, 135 mg-ammonium/kg and C/N of 25:1. The results indicated that 45°C, 90% of relative water content, pH 9.0, 125 mg-ammonium/kg, C/N of 10:1 inhibited *E. coli* growth, whereas, 37°C, 80% of relative soil water content, pH 7.0, 135 mg-ammonium/kg and C/N of 25:1 stimulated *E. coli* growth.

Keywords: *Escherichia coli*, soil environment factors, population dynamics

Introduction

Livestock and poultry breeding industry are major contributors in agricultural productivity in China. They play a major role to fulfill the food demands of ever-increasing population around the globe by

providing protein rich diet. However it also produces a large amount of manure, it is reported that Chinese aquaculture industry generates 3.8 billion tons of manure every year. The livestock manure is considered as an excellent source of nutrients, which can increase the soil fertility by providing organic matter and essential nutrients. The soil fortification with livestock manure potentially improves soil physical and chemical properties, nutrient profiling and ionic contents which positively influence crop growth, quality and yield

*e-mail: sjyang-2008@163.com

[1-2], contrarily, livestock manure contains a large number of pathogenic microbes, which may cause soil environmental pollution and may cause zoonotic diseases if applying to field without treatment [3-5]. Therefore, different countries and organizations have designed different standards for pathogenic microbes in manure before returning to the field. Italy has the most restrictive limit on the number of *E. coli* in wastewater, which is 10^3 cfu L⁻¹, in WHO, Cyprus, Portugal and Spain (10^4 cfu L⁻¹), and in Greece (25000 cfu L⁻¹). China and France have the least restrictive limit on *E. coli* (10^5 cfu L⁻¹). [6] The application of livestock manure in field condition may enhance the population of pathogenic microbes including *E. coli* even if it has been treated. Additionally, various types of soil environments and farmland eco-systems can also favours the growth and population dynamics of microbes in field conditions which may increase the ecological risks.

E. coli is associated with the gastrointestinal tract of warm-blooded animals and has been considered as one of the most important pathogenic microbes in livestock manure. Various studies showed differential growth responses of *E. coli* due to different soil environment conditions. It is reported that *E. coli* survives better at 37°C [7-8], while some other studies find that temperature higher than 4°C inhibit the *E. coli* population in soil [9]. In case of soil water content, some studies indicated that higher soil moisture content (100% field capacity) enhanced the survival rate of *E. coli* [10], while some other studies indicated that moisture content did not significantly affect the inactivation rate of *E. coli* [11]. Adequate levels of carbon and nitrogen positively correlated with growth of microorganisms, hence, suitable soil C/N ratio enhance the *E. coli* growth in soil medium. Most of studies claim that C/N ratio of 25:1 is beneficial for *E. coli* growth, and decreasing levels of C/N will lead to early autolysis of the bacteria, while increase will result in imbalance of bacterial metabolism, but Singh et al. [12] believe that C/N ratio of 16:1 supported longer survival for *E. coli*. Soil pH is one of the important factors affecting survival and growth of soil microorganism, pH 6~8 is considered optimal for most bacteria (such as *E. coli*), and the soil pH of 4~10 is viable, which means bacteria is inhibited or inactivating if pH is lower than the minimum value or higher than the maximum value. Williams et al. [13] reported that the survival rate of *E. coli* is highest in pH 7.0. Wang et al. [14] finds that in the acidic soils t_d values decreased as the pH

decreased or become more acidic resulting shorter t_d values, while in the neutral or alkaline soils ($\text{pH} \geq 6.45$) they observed longer t_d values. It is also reported that varying levels of ammonium and nitrate also affect the population dynamics of *E. coli* in soil, an average level of 5.5 mg ammonia/kg of dry chicken droppings results in a 4.5-log decline typhimurium cells to the point of no recovery within 46 h [15], but Taabodi et al. [16] finds that *E. coli* can absorb ammonium as nutrient when concentrations of ammonium and nitrate are both 1.0 mg L⁻¹, in which utilization of ammonium is 3.35 times higher than nitrate.

Hence, there are different perceptions among studies on the effect of varying physicochemical properties of soil on the survival rate and population dynamic change of *E. coli* in soil and conclusions are mostly based on a single environmental factor in soil. However, soil environment is based on multiple factors which create the soil-microbes-atmosphere-continuum, composed of multiple interconnected factors affecting the population dynamics of microbes. There are limited reports available about the synergistic influence of various physicochemical factors of soil on *E. coli* growth. Therefore, a comprehensive study is designed to evaluate the synergistic influence of various soil environments created by varying levels of temperature, relative water content, pH value and ammonium-N, nitrate-N and C/N ratios. The objective of this study is to investigate the influence of various types of soil environments on population dynamics of *E. coli* in soil, which can provide the theoretical basis for application of biogas slurry for soil fertility and crop cultivation in eco-friendly manner.

Experimental

Slurry and Soil

The experiment was conducted in Yantai Institute of China Agricultural University, Yantai city, Shandong Province ($119^{\circ}34' \sim 121^{\circ}57' \text{ E}$, $36^{\circ}16' \sim 38^{\circ}23' \text{ N}$), from November 5th, 2020 to January 30th, 2021. Slurry (pig slurry) was obtained from a biogas plant in Muping district, Yantai, Shandong Province, and soil used in experiment was brown soil (sandy loam soil). Basic properties of soil and slurry used in this study are listed in Table 1.

Table 1. Characteristics of soil and slurry used in this study.

Material	Characteristics				
	Number of <i>E. coli</i> (cfu/L)	Organic Matter Content (%)	Ammonium-N Content (mg/kg)	Nitrate-N Content (mg/kg)	TS (%)
Soil	0.00±0.00	1.201±0.05	21.15±1.02	2.95±0.54	\
Slurry	$3.05 \times 10^7 \pm 2000$	0.46±0.04	453.09±2.53	172.72±2.55	1.57±0.08

Methods

Soil column experiment was conducted in PVC tube (7.5 cm diameter, 25 cm depth). In this study five different factors (soil temperature, relative water content, pH value, ammonium-N content, C/N ratio) were set up to explore the effect of each factor, respectively. Soil was air-dried, ground and sieved (<2 mm), then taken into the column, each column was filled with 1000 g soil. After that every column was filled with 140 g slurry and water content was adjusted with sterile water. The top and bottom of column were sealed with high temperature resistant tissue sealing film.

Experiment Design

Soil Temperature Treatments

Four temperature treatments were set up at 0.6°C, 25°C, 37°C and 45°C, temperature was kept through constant temperature incubator.

Soil Water Content Treatments

Three relative soil water content treatments were set up at 60% (soil water content 21.8%), 80% (soil water content 29.1%), 90% (soil water content 34.5%).

Soil pH Treatments

Three soil pH treatments were set up at 5.0, 7.0, 9.0, soil pH was adjusted through 0.1 mol/L H₂SO₄ and NaOH.

Soil Ammonium-N Content Treatments

Three soil ammonium-N treatments were set up at 125 mg/kg, 135 mg/kg, 145 mg/kg (nitrate-N contents were 25 mg/kg, 15 mg/kg, 5 mg/kg, respectively).

Soil C/N Treatments

Three soil C/N treatments were set up at 10:1, 25:1, 40:1, soil C/N was adjusted through organic fertilizers.

All experiments were repeated three times. Adjusted soil temperature at 37°C, relative water content to 60% (soil water content 21.8%), pH to 7.0, ammonium-N content to 135 mg/kg, nitrate-N content to 15 mg/kg and soil C/N to 25:1.

Sampling and Examination

Soil samples were collected by inserting a cylindrical sampler (2 cm diameters, 25 cm high) vertically into soil column 20 cm from the center of soil column surface after 1, 2, 4, 8, 15, 30, 45 and 60 days of incubation, experimental units were separate units for each timepoint. The population of *E. coli* was estimated by the enumeration of typical colonies and were reported as colony-forming units per g of fresh soil (cfu g⁻¹) [17], soil ammonia concentration were determined through Kjeldahl distillation [18], nitrate-N content were measured using a high-resolution continuous-flow analyzer AutoAnalyzer3 [19], soil organic carbon (SOC) and total nitrogen (TN) content were determined by potassium dichromate oxidation method and Kjeldahl method, respectively [20]. Soil oxygen content was determined through gas detector (Qingdao Junyuan environmental protection equipment co. LTD, 20H17041).

Data Analysis

Main text paragraph. The effects of the treatments were tested by one-way analysis of variance (ANOVA). Means were compared between the treatments using the LSD (least significant difference) test at the 0.05 probability level.

Results and Discussion

Influence of Soil Temperature on Survival and Population of *E. coli*

Statistical analysis showed that compared with 0.6°C treatment, counts of *E. coli* in soil at 25°C, 37°C and 45°C were increased significantly after 8 days of incubation ($P < 0.05$), which were increased by 39.92%, 85.02%, 19.96%, respectively (Table 2), after 60 days

Table 2. Influence of soil temperature on counts of *E. coli*

Temperature (°C)	Counts of <i>E. coli</i> in soil (10 ⁶ cfu/kg)							
	1 d	2 d	4 d	8 d	15 d	30 d	45 d	60 d
0.6	3.30ab	3.40bc	3.55c	4.12d	4.57c	3.54c	2.39b	1.12a
25	3.77a	4.05b	4.64b	5.76b	5.87b	4.17b	2.29b	0.46ab
37	3.86a	4.79a	6.48a	7.62a	7.30a	5.42a	3.22a	0.81a
45	3.54b	3.67bc	4.31b	4.94c	4.76c	3.10cd	1.39c	0.21ab

Note: Different letters in the table refer to statistically significant differences ($P < 0.05$) among different treatments, the followed are as the same.

Table 3. Influence of soil water content on counts of *E. coli*.

Soil Water Content	Counts of <i>E. coli</i> in soil (10 ⁶ cfu/kg)							
	1 d	2 d	4 d	8 d	15 d	30 d	45 d	60 d
21.8%	3.86a	4.79a	6.48ab	7.62b	7.30ab	5.42b	3.22a	0.81ab
29.1%	4.03a	4.88a	6.81a	8.20a	7.84a	5.96a	3.74a	1.20a
34.5%	3.69a	4.11ab	5.77b	6.18c	5.57c	3.78c	2.12b	0.33b

of incubation, count of *E. coli* at 0.6°C was 1.44, 0.39, 4.35 times higher than which at 25°C, 37°C, 45°C, respectively, there was no significant difference among treatments ($P>0.05$). In terms of culture time, counts of *E. coli* in soil at 0.6°C, 25°C, 37°C and 45°C were all increased first and then decreased with the increasing time, the treatments of 37°C and 45°C peaked at 8th day of incubation, and 0.6°C, 25°C treatments peaked after 15 days of incubation. The maximum *E. coli* count was observed at 8 days of incubation at 37°C treatment being the highest, followed by 25°C, 45°C, 0.6°C.

Influence of Soil Water Content on Survival and Population of *E. coli*

Statistical analysis showed that compared with soil water content 34.5% treatment, counts of *E. coli* in soil at 21.8% and 29.1% were increased significantly after 8 days of incubation ($P<0.05$), which were increased by 23.26% and 32.68%, respectively (Table 3), after 60 days of incubation, count of *E. coli* at 29.1% treatment was 1.48 and 3.64 times higher at 21.8% as compared to 34.5%, significant difference ($P<0.05$) was found at 60th day. In terms of culture time, counts of *E. coli* in all treatments were increased first and then decreased with the increase in time, which peaked after 8 days of incubation with 29.1% treatment being the highest, followed by 21.8%, 34.5%.

Influence of Soil pH on Survival and Population of *E. coli*

Statistical analysis showed that compared with soil pH 9.0, counts of *E. coli* in soil at pH 5.0 and 7.0 were increased significantly after 8 days of incubation ($P<0.05$), which were increased by 37.98% and 78.16%, respectively. After 60 days of incubation, count of *E. coli* at soil pH 7.0 and 5.0 treatments were 1.72 and 4.26 times higher as compared to pH 9.0, there was no significant difference among treatments ($P>0.05$) (Fig. 1). In terms of culture time, counts of *E. coli* in all treatments were increased first and then decreased as time went by, which peaked after 8 days of incubation with soil pH 7.0 treatment being the highest, followed by soil pH 5.0, 9.0. Population of *E. coli* in all treatments decreased to less than 10⁵ cfu/kg after 60 days of incubation.

Influence of Soil Ammonium-N and Nitrate-N on Survival of *E. coli*

Fig. 2 showed that compared with 125 mg-ammonium/kg, counts of *E. coli* in soil at 135 mg-ammonium/kg and 145 mg-ammonium/kg were increased significantly after 8 days of incubation ($P<0.05$), which were increased by 22.08% and 13.14%, respectively. After 60 days of incubation, count of *E. coli* in soil at 135 mg-ammonium/kg was 3.00, 2.03 times than which at 125 mg-ammonium/kg and 145 mg-ammonium/kg in turn, there was no significant difference among treatments ($P>0.05$). In terms of culture time, counts of *E. coli* in all treatments were increased first and then decreased as time went by, which peaked after 8 days of incubation with 135 mg-ammonium/kg treatment being the highest, followed by 145 mg-ammonium/kg, 125 mg-ammonium/kg. The population of *E. coli* in all treatments decreased to less than 10⁵ cfu/kg after 60 days of incubation.

Influence of Soil C/N on Survival and Population of *E. coli*

Statistical analysis showed that compared with low C/N (10:1), counts of *E. coli* in soil at C/N 25:1 and 40:1 treatments were increased significantly after 8 days of incubation ($P<0.05$), which were increased by 11.88% and 5.12%, respectively. After 60 days of incubation, count of *E. coli* in soil at C/N of 25:1 was 3.49, 1.81 times than which at C/N 10:1 and 40:1 treatments in turn, there was no significant difference among treatments ($P>0.05$) (Fig. 3). In terms of culture time, counts of *E. coli* in all treatments were increased first and then decreased as time went by, which peaked after 8 days of incubation with C/N of 25:1 treatment being the highest, followed by C/N 40:1, 10:1. The population of *E. coli* in all treatments decreased to less than 10⁵ cfu/kg after 60 days of incubation.

Discussion

The results showed that *E. coli* growth was stimulated in soil at 37°C and inhibited when soil temperature was higher or lower than 37°C (0.6°C, 25°C, 45°C). It means 37°C is optimum temperature for growth of *E. coli*, it might be suitable for enzyme activities in *E. coli*, which ultimately promoted




Fig. 1. Influence of soil pH on counts of *E. coli*.

Note: Different letters in the figure refer to statistically significant differences ($P < 0.05$) among different treatments, the followed are the same.




Fig. 2. Influence of soil ammonium-N and nitrate-N on counts of *E. coli*.




Fig. 3. Influence of soil C/N on counts of *E. coli*.

E. coli growth, and decrease of temperature resulted in *E. coli* transient growth inhibition and a cold shock response consisting of transient induction of several proteins, repression of heat shock proteins [21-22]. There was a dramatic increase in protein degradation at high temperature (>37°C), which resulted in inhibited *E. coli* metabolism [23]. Our results showed that counts of *E. coli* in the soil at 0.6°C was higher than which at other temperatures after 60 days of incubation, it might be attributed to greater amount of protective enzymes and other proteins and also to a lower degree of permeability of the envelopes to toxic [24]. Other temperature treatment (25°C) was conducive to soil nitrification, which led to increased soil nitrate-N content and inhibited *E. coli* growth [25-26], or stimulated formation of protein aggregates at elevated temperatures (45°C) which led to complete inactivation of the protein [27].

Water content is one of the important factors of soil physical properties and has important effect on reproduction of soil microorganisms. The results showed that regardless of length of culture time, counts of *E. coli* in soil at relative water content of 80% treatment was always higher than which at 60% and 90% treatments. It illustrated that 80% relative soil water content improved *E. coli* metabolic activity, *E. coli* community-level activity decreases with decreasing soil moisture (60%) [28], soil oxygen content decreased as water content increased (90% of relative soil water content), which led to inhibited *E. coli* growth [29-30].

The results showed that counts of *E. coli* in soil was highest with soil pH 7.0 treatment (Fig. 1). It illustrated that soil with pH 7.0 provided a suitable acid-base environment for the propagation and growth of *E. coli* and improved its metabolic activity [31]. Counts of *E. coli* in soil decreased as soil pH increased (9.0) or decreased (5.0). That was because high adsorption capacities of Fe/Al (hydro) oxides with positive charges in acidic soils led to high inactivation for *E. coli* [32], and it was likely to be more negatively charged with increasing pH levels and could have led to the destruction *E. coli* at pH 9.0, meanwhile greater antimicrobial potential of humic acid contributed to inactivation for *E. coli* as it is more soluble at higher pH levels [33]. According to Fig. 1, alkaline soil (pH 9.0) have a stronger inhibitory effect on *E. coli* growth compared with acidic soil (pH 5.0). It might be because soil with lower pH provided less substrate for soil nitrification and dampens the soil microbial activity of nitrification, which led to a decrease of soil nitrate-N content and the toxicity of nitrate-N to *E. coli* was weaken, in soil with high pH (9.0) nitrate-N content increased because of promoted nitrification, which strengthened inhibition on *E. coli* growth [34].

The results showed that there were different positive or negative effects of contents of ammonium-N and nitrate-N in soil on *E. coli* growth (Fig. 2). Counts

of *E. coli* in soil showed a trend of increasing first and then decreasing after the application of slurry, it might be because at initial stage of application of slurry ammonium-N content in soil was enough as the nitrogen source for *E. coli* growth, but contents of ammonium-N and nitrate-N was decreased and increased, respectively with extension of culture time caused by mineralization, immobilization, nitrification, which inhibited *E. coli* growth [25, 35-36]. Compared with 135 mg-ammonium/kg (25 mg-nitrate-N/kg), the decrease of counts of *E. coli* in soil at treatment of 125 mg-ammonium/kg (35 mg-nitrate-N/kg) was caused by the inhibitory effect of nitrate-N (35 mg/kg) on *E. coli*, and the decrease at treatment of 145 mg-ammonium/kg (15 mg-nitrate-N/kg) showed that high content of ammonium-N also led to inhibition of *E. coli* growth [37].

Statistical analysis showed that soil environment with C/N of 25:1 was beneficial for *E. coli* growth, soil environments with C/N 10:1 and 40:1 inhibited *E. coli* growth, and inhibitory effect of soil C/N of 10:1 was more significant than which of 40:1, which was the same as conclusions in previous study [38]. The reason resulted in such phenomenon was that environment with C/N of 25:1 was more suitable for level of microorganism metabolic activity [39], too high C/N (40:1) led to mineral N immobilization, which limited *E. coli* to utilize nitrogen as nutrients source [40], and low C/N increased the net nitrification rates, which increased soil nitrate-N content and led to inhibition on *E. coli* survival [40-42].

Conclusions

The influence of different soil environments on population dynamics of *E. coli* were determined in the study. The results showed that, after 60 days of incubation, the soil treated with 0.6°C, 80% of relative soil water content, pH 7.0, 135 mg-ammonium/kg and C/N of 25:1 had the maximum of *E. coli*, and 45°C, 90% of relative water content, pH 9.0, 125 mg-ammonium/kg, C/N of 10:1 inhibited *E. coli* growth. It also indicated that 37°C, 80% of relative soil water content, pH 7.0, 135 mg-ammonium/kg and C/N of 25:1 stimulated *E. coli* growth, which led to the peak of coliforms occurring during cultivation. Also, interactions among soil environment factors on *E. coli* should be taken into comprehensive account.

Acknowledgments

This work was supported by Yantai Education Bureau Subject Development Project (project No. 2021XDRHXMQT19). We are equally indebted to Yantai Institute of China Agricultural University for providing research facilities.

Conflict of Interest

The authors declare no conflict of interest.

References

- SHIBIN L., JINYANG W., SHENGYAN P., EVGENIA B., YAKOV K., S. R.B. Impact of manure on soil biochemical properties: A global synthesis. *Science of the Total Environment*, **745**, 141003, **2020**.
- YONGHUA L., NAN B., ZHENKUI T., XIAOTIAN M., GANG H., ZHAOHUI W. Rethinking application of animal manure for wheat production in China. *Journal of Cleaner Production*, **318** (1), 128473, **2021**.
- LIU W.R., ZENG D., SHE L., SU W.X., HE D.C., WU G.Y., MA X.R., JIANG S., JIANG C.H., YING G.G. Comparisons of pollution characteristics, emission situations, and mass loads for heavy metals in the manures of different livestock and poultry in China. *Science of the Total Environment*, **734**, 139023, **2020**.
- VENGLOVSKY J., SASAKOVA N., GREGOVA G., PAPAJOVA I., TOTH F., SZABOOVA T. Devitalisation of pathogens in stored pig slurry and potential risk related to its application to agricultural soil. *Environmental Science and Pollution Research*, **25** (22), 21412, **2018**.
- MARSZALEK M., KOWALSKI Z., MAKARA A. The possibility of contamination of water-soil environment as a result of the use of pig slurry. *Ecological Chemistry and Engineering S-Chemia I Inzynieria Ekologiczna S*, **26** (2), 313, **2019**.
- JEONG H., KIM H., JANG T. Irrigation Water Quality Standards for Indirect Wastewater Reuse in Agriculture: A Contribution toward Sustainable Wastewater Reuse in South Korea. *Water*, **8** (4), 169, **2016**.
- TACK I.L.M.M., NIMMEGEERS P., AKKERMANS S., LOGIST F., VAN IMPE J.F.M. A low-complexity metabolic network model for the respiratory and fermentative metabolism of *Escherichia coli*. *Plos One*, **13** (8), e0202565, **2018**.
- SOARES A.S., MIRANDA C., TEIXEIRA C.A., COUTINHO J., TRINDADE H., COELHO A.C. Impact of different treatments on *Escherichia coli* during storage of cattle slurry. *Journal of Environmental Management*, **236** 323, **2019**.
- XUAN THANH B., WOLFF A., MADSEN M., BANG D.D. Fate and survival of *Campylobacter coli* in swine manure at various temperatures. *Frontiers in Microbiology*, **2** 262, **2011**.
- COOLS D., MERCKX R., VLASSAK K., VERHAEGEN J. Survival of *E. coli* and *Enterococcus* spp. derived from pig slurry in soils of different texture. *Applied Soil Ecology*, **17** (1), 53-62, **2001**.
- ERICKSON M.C., HABTESELASSIE M.Y., LIAO J., WEBB C.C., MANTRIPRAGADA V., DAVEY L.E., DOYLE M.P. Examination of factors for use as potential predictors of human enteric pathogen survival in soil. *Journal of Applied Microbiology*, **116** (2), 335, **2014**.
- SINGH R., KIM J., SHEPHERD M.W., JR., LUO F., JIANG X. Determining Thermal Inactivation of *Escherichia coli* O157:H7 in Fresh Compost by Simulating Early Phases of the Composting Process. *Applied and Environmental Microbiology*, **77** (12), 4126, **2011**.
- WILLIAMS M.L., LEJEUNE J.T., MCSPADDEN GARDENER B. Soil Conditions That Can Alter Natural Suppression of *Escherichia coli* O157:H7 in Ohio Specialty Crop Soils. *Applied and Environmental Microbiology*, **81** (14), 4634, **2015**.
- WANG H.Z., WEI G., YAO Z.Y., LOU J., XIAO K.C., WU L.S., WU J.J., XU J.M. Response of *Escherichia coli* O157:H7 survival to pH of cultivated soils. *Journal of Soils and Sediments*, **14** (11), 1841, **2014**.
- TURNBULL P.C., SNOEYENBOS G.H. Roles of ammonia, water activity, and pH in salmonellacidal effect of long-used poultry litter. *Avian Diseases*, **17** (1), 72, **1973**.
- TAABODI M., HASHEM F.M., OSCAR T.P., PARVEEN S., MAY E.B. The Possible Roles of *Escherichia coli* in the Nitrogen Cycle. *International Journal of Environmental Research*, **13** (3), 597, **2019**.
- NL L., SR S., DM B.T., EB P., CL R. Decay of *Escherichia coli* in soil following the application of biosolids to agricultural land. *Water and Environment Journal*, **17** (1), 23, **2003**.
- MASSIMO Z., SILVIA M., FULVIA T., BARBARA S., GABRIELE B., ANDREA S., FABRIZIO A. The distribution of functional N-cycle related genes and ammonia and nitrate nitrogen in soil profiles fertilized with mineral and organic N fertilizer. *Plos One*, **15** (6), e0228364, **2020**.
- CHEN Y., ZHANG P., WANG L., MA G., LI Z., WANG C. Interaction of nitrogen and phosphorus on wheat yield, N use efficiency and soil nitrate nitrogen distribution in the North China Plain. *International Journal of Plant Production*, **14** (3), 415, **2020**.
- PATHAK P., REDDY A.S. Vertical distribution analysis of soil organic carbon and total nitrogen in different land use patterns of an agro-organic farm. *Tropical Ecology*, **62** (3), 386, **2021**.
- PINSACH J., DE MAS C., LOPEZ-SANTIN J., STRIEDNER G., BAYER K. Influence of process temperature on recombinant enzyme activity in *Escherichia coli* fed-batch Cultures. *Enzyme and Microbial Technology*, **43** (7), 507, **2008**.
- JONES P.G., CASHEL M., GLASER G., NEIDHARDT F.C. FUNCTION OF A RELAXED-LIKE STATE FOLLOWING TEMPERATURE DOWNSHIFTS IN *ESCHERICHIA-COLI*. *Journal of Bacteriology*, **174** (12), 3903, **1992**.
- FAREWELL A., NEIDHARDT F.C. Effect of temperature on in vivo protein synthetic capacity in *Escherichia coli*. *Journal of Bacteriology*, **180** (17), 4704, **1998**.
- CEBRIAN G., SAGARZAZU N., PAGAN R., CONDON S., MANAS P. Resistance of *Escherichia coli* grown at different temperatures to various environmental stresses. *Journal of Applied Microbiology*, **105** (1), 271, **2008**.
- TAABODI M., MAY E.B., BRYANT R.B., SAPORITO L.S., SKEEN O.K., HASHEM F.M., ALLEN A.L. *Aeromonas hydrophila*, *Bacillus thuringiensis*, *Escherichia coli* and *Pseudomonas aeruginosa* utilization of Ammonium-N, Nitrate-N and Urea-N in culture. *Heliyon*, **6** (4), e03711, **2020**.
- THIAGALINGAM K., KANEHIRO Y. Effect of temperature on nitrogen transformation in Hawaiian soils. *Plant and Soil*, **38** (1), 177, **1973**.
- GUR E., BIRAN D., GAZIT E., RON E.Z. In vivo aggregation of a single enzyme limits growth of *Escherichia coli* at elevated temperatures. *Molecular Microbiology*, **46** (5), 1391, **2002**.
- MANZONI S., SCHIMEL J.P., PORPORATO A. Responses of soil microbial communities to water stress:

- results from a meta-analysis. *Ecology*, **93** (4), 930, **2012**.
29. BALOGH J., PINTER K., FOTI S., CSERHALMI D., PAPP M., NAGY Z. Dependence of soil respiration on soil moisture, clay content, soil organic matter, and CO₂ uptake in dry grasslands. *Soil Biology & Biochemistry*, **43** (5), 1006, **2011**.
30. WAGAI R., BRYE K.R., GOWER S.T., NORMAN J.M., BUNDY L.G. Land use and environmental factors influencing soil surface CO₂ flux and microbial biomass in natural and managed ecosystems in southern Wisconsin. *Soil Biology & Biochemistry*, **30** (12), 1501, **1998**.
31. LAWFORD H.G., ROUSSEAU J.D. Effects of pH and acetic-acid on glucose and xylose metabolism by a genetically-engineered ethanologenic *Escherichia-coli*. *Applied Biochemistry and Biotechnology*, **39** 301, **1993**.
32. WANG H., ZHANG T., WEI G., WU L., WU J., XU J. Survival of *Escherichia coli* O157:H7 in soils under different land use types. *Environmental Science and Pollution Research*, **21** (1), 518, **2014**.
33. BAKER C.A., DE J., SCHNEIDER K.R. *Escherichia coli*O157 survival in liquid culture and artificial soil microcosms with variable pH, humic acid and clay content. *Journal of Applied Microbiology*, **130** (2), 416, **2021**.
34. ZHAOLEI L., ZHAOQI Z., DASHUAN T., JINSONG W., ZHENG F., FANGYUE Z., RUIYANG Z., WEINAN C., YIQI L., SHULI N. Global patterns and controlling factors of soil nitrification rate. *Global Change Biology*, **26** (7), 4147, **2020**.
35. WYSOCKA-CZUBASZEK A. Dynamics of nitrogen transformations in soil fertilized with digestate from agricultural biogas plant. *Journal of Ecological Engineering*, **20** (1), 108, **2019**.
36. CHERNIAEVA I.I., LAGUTINA T.M. Effect of ammonium and nitrate nitrogen on the physiological and biochemical properties of *Bacillus brevis* and *Pseudomonas mycophaga*. *Prikladnaia biokhimiia i mikrobiologiia*, **14** (1), 67, **1978**.
37. SINGH R., KIM J., JIANG X. Heat inactivation of *Salmonella* spp. in fresh poultry compost by simulating early phase of composting process. *Journal of Applied Microbiology*, **112** (5), 927, **2012**.
38. THOMAS C., IDLER C., AMMON C., AMON T. Effects of the C/N ratio and moisture content on the survival of ESBL-producing *Escherichia coli* during chicken manure composting. *Waste Management*, **105**, 110, **2020**.
39. WU X., YAO W., ZHU J., MILLER C. Biogas and CH₄ productivity by co-digesting swine manure with three crop residues as an external carbon source. *Bioresource Technology*, **101** (11), 4042, **2010**.
40. BRUUN E.W., AMBUS P., EGSGAARD H., HAUGGAARD-NIELSEN H. Effects of slow and fast pyrolysis biochar on soil C and N turnover dynamics. *Soil Biology & Biochemistry*, **46** 73, **2012**.
41. ZHAO Y., ZHANG J., MUELLER C., CAI Z. Temporal variations of crop residue effects on soil N transformation depend on soil properties as well as residue qualities. *Biology and Fertility of Soils*, **54** (5), 659, **2018**.
42. DAI X., ZHOU W., LIU G., LIANG G., HE P., LIU Z. Soil C/N and pH together as a comprehensive indicator for evaluating the effects of organic substitution management in subtropical paddy fields after application of high-quality amendments. *Geoderma*, **337** 1116, **2019**.