

Seasonal Measurement of Greenhouse Gas Concentrations and Emissions Along the Longitudinal Profile of a Small Stream

Václav Mach^{1*}, Adam Bednařík¹, Lubomír Čáp², Jan Šipoš¹, Martin Rulík¹

¹Department of Ecology and Environmental Sciences, Palacky University, Faculty of Science, Šlechtitelů 11, CZ-783 71 Olomouc, Czech Republic

²Department of Analytical Chemistry, Faculty of Science, Palacky University Tř. 17. Listopadu 1192/12, CZ-771 46 Olomouc, Czech Republic

Received: 29 September 2015

Accepted: 2 February 2016

Abstract

In order to find out whether streams might be a major source of greenhouse gases to the atmosphere, our investigation sought to determine the total emissions of CH₄, CO₂, and N₂O from the surface water of a small stream. Over a period of a year we used floating chambers to measure gas emissions along the longitudinal profile of Sitka Stream (Czech Republic). Additionally, we measured gas concentrations of surface and interstitial waters. We found that interstitial and surface waters were supersaturated by all monitored gases – especially by CH₄ – and that the stream is a significant emitter of these greenhouse gases. The concentrations and the emission rates of all three gases were higher in the downstream part than upstream. In the case of CH₄ the majority of total annual emissions (90%) was released from the most downstream section, representing only 1/5 of the stream's total surface area (0.18 km²). The majority of CH₄ and CO₂ emissions were released during warmer periods of the year and the highest N₂O emissions from Sitka were recorded during summer and winter. The total annual emissions of CH₄, CO₂, and N₂O into the atmosphere from the water's surface were estimated to be 0.6 t, 210 t, and 0.2 t, respectively. After conversion of the greenhouse gas emissions to CO₂ equivalents using a calculation by IPCC, CO₂ accounts for the majority of total annual emissions of greenhouse gases (70.1%), with the second being N₂O (22.7%), and the last CH₄ (7.2%) for a 100-year time horizon. This work brings worthwhile data of greenhouse gas emissions and concentrations from a small water stream based on seasonal measurements along the longitudinal profile.

Keywords: methane, greenhouse gases, stream water, carbon dioxide, nitrous oxide

Introduction

In 2011 the global mean concentration of atmospheric CO₂ was 391 ppm [1], and the average rate of increase

in the concentration of atmospheric CO₂ over the period spanning from 1960 to 2005 of 1.4 ppm yr⁻¹ is therefore considered to be the main driver of climate change [2]. The global mean concentrations of CH₄ and N₂O in 2011 were 1,803 ppb and 324 ppb, respectively [1]. The significant ability of CH₄ and N₂O molecules to absorb

*e-mail: vaclav.mach@riseup.net

infrared radiation makes them more efficient greenhouse gases than CO_2 , thus resulting in a substantial contribution to the heating effect of the atmosphere in particular, and to climate change in general. In the Fifth Assessment Report of the IPCC, Myhre et al. [3] reconsidered the global warming potentials (GWPs) as an instrumental metric for transferring emissions of different gases to a common scale. The recently suggested GWPs of CH_4 rose significantly from 72 to 86 and from 25 to 34 for the 20-year and 100-year time horizons, respectively, which makes CH_4 a more important greenhouse gas. The radiative forcing of a molecule of N_2O is even greater, and the GWPs for N_2O are 268 and 298, respectively, for the 20-year and 100-year time horizons.

With respect to concerns about climate change, numerous studies have suggested possible ways and strategies to mitigate emissions of greenhouse gases, particularly those originating from anthropogenic sources. On the other hand, some natural sources of these gases were neglected owing to the fact that we have only limited information about their role in the production of greenhouse gases [4]. Extensive efforts have been directed toward the identification and quantification of the sources of greenhouse gas emissions into the atmosphere from a variety of aquatic environments. Wetlands are believed to be a major natural source of CO_2 and CH_4 [5-7], however both CO_2 and CH_4 are also supersaturated in many surface waters, with streams and rivers frequently exhibiting partial pressures of gases many times in excess of the atmospheric equilibrium [8-9]. The production of N_2O is linked mainly with excesses of nitrogen in agricultural soils and its runoff in drainage water [10]. However, streams similarly show significant supersaturation of N_2O with respect to the atmosphere [11]. Unquantified potential sources of greenhouse gases are of obvious climatic concern, and while streams and rivers are not usually implicated directly, there is a growing body of evidence to suggest that alongside CO_2 and N_2O , rivers and streams can also emit significant quantities of CH_4 [12-13]. Saarnio et al. [14] estimated CH_4 release from wetlands and watercourses in Europe, but data for rivers and streams still have a high error rate. If we consider that rivers and streams are likely to cover 0.30-0.56% of the land surface (with a significant partition of third-order streams) and make contributions to global processes and greenhouse gas emissions that may be 20-200% greater than those implied by previous estimates [15], explorations of greenhouse gases productions and emissions in river ecosystems are required.

Long-term study on Sitka documented the production of greenhouse gases in hyporheic sediments [16]. Hlaváčová et al. [17] reported that respiration of oxygen, nitrate, sulphate, and methanogenesis may simultaneously coexist within the hyporheic zone, and that an anaerobic metabolism and methanogenesis appear to be an important pathway in organic carbon cycling in the Sitka stream sediments. Additionally, Cupalová and Rulík [18] found that members of the domain *Archaea* may commonly occur within well-oxygenated hyporheic sediments.

Buriánková et al. [19] found relatively well-developed populations of methanogenic *Archaea*, and significant methanogenic potential in hyporheic Sitka sediments. The methanogenesis and other degradation processes result in the production of CO_2 , CH_4 , and N_2O , which then supersaturate the interstitial water and eventually emit them into the atmosphere [20]. Based on previous studies, we suppose that stream sediments are a source of gases that are important in global warming. In particular, CH_4 production in hyporheic sediments seems to be novel as published global CH_4 emission estimates from streams are also sparse [21]. Moreover, recent discoveries of metabolic pathways in both the nitrogen and carbon cycle across a spectrum of aquatic ecosystems highlights the need for new directions and a multidisciplinary approach to quantify the flux of carbon and nitrogen through rivers [22].

Past studies have focused mainly on the downstream section of Sitka, but we were also curious to know if other parts along the stream's profile will also produce greenhouse gases. In contrast to previous studies concerned with greenhouse gas emissions, we extended our exploration to five study sites along a longitudinal profile of Sitka. The main objectives of our study were to measure the concentrations of CH_4 , CO_2 , and N_2O both in surface and interstitial waters, and their emissions at different localities along a stream continuum, and to estimate the total annual emissions of greenhouse gases from surface water to the atmosphere. Our study presents at once total annual emissions of all three biogenic greenhouse gases from a small water stream, and their relative contribution to total annual emissions expressed in standard metric units (CO_2 equivalent emission). This study is worthwhile mainly because there is a paucity of studies estimating greenhouse gas emissions from the whole river ecosystem during a year.

Material and Methods

Study Site

The sampling sites were located on Sitka Stream in the Czech Republic (Fig. 1). Sitka is an undisturbed, third-order, 35-km stream originating in the Hrubý Jeseník




Fig. 1. Localities (black circles) on Sitka Stream.

Table 1. Longitudinal physicochemical patterns of Sitka Stream (means \pm standard error).

Variable/ Locality	I.	II.	III.	IV.	V.
Elevation above sea-level [m]	535	330	240	225	215
Distance from the spring [km]	6.9	18.2	25.6	30.9	34.9
Average flow velocity [m.s ⁻¹]	0.46 \pm 0.09	0.19 \pm 0.04	0.45 \pm 0.08	0.39 \pm 0.04	0.20 \pm 0.04
Stretch surface area [km ²]	0.043	0.060	0.043	0.024	0.012
Grain median size [mm]	12.4	12.9	13.2	0.2	5.4
Surface dissolved oxygen saturation [%]	101 \pm 22.6	110 \pm 13.4	104 \pm 3.7	114 \pm 14.6	103 \pm 3.8
Hyporheic dissolved oxygen saturation [%]	63 \pm 23.6	89 \pm 13.8	79 \pm 4.4	61 \pm 9.2	58 \pm 12.7
Surface water temperature [°C]	9.1 \pm 3.59	9.7 \pm 3.56	12.0 \pm 4.39	12.3 \pm 4.93	11.0 \pm 5.27
Interstitial water temperature [°C]	13.9 \pm 1.17	14.8 \pm 1.40	17.5 \pm 1.72	18.3 \pm 2.04	14.2 \pm 3.44
Surface water DOC* [mg/l]	2.47 \pm 0.21	0.81 \pm 0.19	2.62 \pm 0.15	2.69 \pm 0.19	3.74 \pm 0.39
Interstitial water DOC* [mg/l]	2.05 \pm 0.17	1.31 \pm 0.29	2.71 \pm 0.32	5.76 \pm 0.91	2.62 \pm 0.24

* dissolved organic carbon

Mountains at 650 m above sea level. With the exception of short sections, the Sitka is unregulated with well-established riparian vegetation. A detailed description of the stream can be found in Hlaváčová et al. [17]. In order to assess the emissions released from a total stream area, the stream was divided into five stretches according to changing conditions along the river's course (Table 1). For each stretch we chose one representative sampling site (localities I-V) where samples of both surface water and air were taken repeatedly. The numbering of each locality corresponds to the numbering of each stretch. The stream flows in its upper section until it reaches the town of Šternberk through a forested area with low intensity of anthropogenic effects (localities I-II), while the lower course of the stream is naturally meandering through a more intensively managed agricultural landscape (localities III-V). The areas of the sections of stream were calculated from known lengths and mean channel widths (measured by a metal measuring type at different sites). The longitudinal distances of the sections were worked out using ArcGIS software with digitized maps of the stream and GPS coordinates that were obtained during field measurements. The total area of Sitka Stream was estimated at 0.18 km².

Water Sampling

Surface water samples were collected from running water at a depth of about 10 cm below the surface level at all localities four times from autumn 2005 to autumn 2006 (25 October, 28 February, 25 May, and 18 July). Surface water samples were taken by submersing glass vials under the water level. Glass vials were washed by surface water and sealed under the water level. Three replicates of surface water were collected on each sampling date. All water samples were taken between 09:00 and 14:00. Interstitial water samples were collected using a set of

two mini-piezometers randomly placed at a depth of 30 cm in sediments at each study site twice in 2006 (25 May and 18 July). The initial 100 ml of water from each mini-piezometer was used as a rinse and was discarded. Three replicates of interstitial water from each mini-piezometer were collected from a continuous column of water using a 100 ml polypropylene syringe connected to a hard PVC tube, and drawn from a mini-piezometer and overflowed and sealed into vials. All water samples were collected into sterile, clear vials (40 ml) with screw-tops, covered by a polypropylene cap with PTFE silicone septa, and stored in ice on the way to the laboratory. On each sampling date, temperatures of interstitial and surface waters were measured for calculating saturation ratios. Saturation ratio R was calculated as the measured concentration of gas divided by the concentration in equilibrium with the atmosphere at the temperature of the sampled water using the solubility data of Wiesenburg and Guinasso [23], Weiss and Price [24], and Weiss [25].

Measurement of Emissions

Gas fluxes across the air-water interface were determined by the floating chamber method [26-27] four times from autumn 2005 to summer 2006 (25 October, 28 February, 25 May, and 18 July). On each sampling date, 4-5 replicated measurements were performed at all the localities between 09:00 and 14:00. The open-bottom floating polyethylene chambers (internal diameter 20 cm, total volume 5.5 L) were held in place on the water surface by a floating body (polystyrene) attached to the outside. The chambers were attached to a metal pole anchored to the stream bed in order to hold the chambers in flowing conditions. The chambers were allowed to float on the water surface for three hours. Preincubations were performed to assess linearity of gas concentrations in

headspace of the chambers and establish the incubation time required for reliable flux measurements. Due to trees on the banks, the chambers at all the study sites were continuously in the shade. On each sampling occasion, ambient air samples were collected for determining initial background concentrations. Samples of headspace gas were collected through a rubber stopper inserted at the top of each chamber, and stored in 100 ml polyethylene gas-tight syringes until analysis. Emissions were calculated as the differences between initial background and comparing air concentration in the chamber headspace, and expressed in one square metre of the surface level per one day according to the formula:

$$F = \frac{(c_I \times c_R) \times V \times 24 / t \times 1000}{p} \quad (1)$$

...where F is a gas flux in $\text{mg m}^{-2}\text{day}^{-1}$, c_I is a concentration of a particular gas in the chamber headspace in $\mu\text{g l}^{-1}$, c_R is a concentration of a particular gas in the background air, V is volume of the chamber in L, t is time of incubation in hr, and p is an area of the chamber expressed in m^2 .

Analysis of Gases, Calculations, and Statistics

Concentrations of dissolved gases in the water and air samples were measured using gas chromatography as described in Hlaváčová et al. [20]. Estimations of total annual emissions into the atmosphere from the five segments of the stream were calculated according to the following formula that we developed:

$$E_a = \frac{\sum p_i \times F_i \times 365}{1000000} \quad (2)$$

...where E_a is average estimation of emission of a particular gas from the total stream area in t yr^{-1} for CO_2 and kg yr^{-1} for CH_4 and N_2O , p_i is an area of stretch in m^2 representing the given locality, F_i is the average emission of a particular gas from the given locality (expressed in $\text{g m}^{-2}\text{day}^{-1}$ for CO_2 and in $\text{mg m}^{-2}\text{day}^{-1}$ for CH_4 and N_2O).

Data analysis was performed using statistical software R (version 2.6.0). Data for regression analyses embodies non-parametric distributions, and normality was not achieved by data transformation. Shapiro-Wilk normality test was used ($P < 0.05$). Therefore, robust regressions were used and linear models were tested by robust F-test for the relationships between gas concentrations and gas emissions, and for the relationships between surface and interstitial water concentrations.

Results

Patterns of Gas Concentrations

Highly supersaturated concentrations of all observed gases were found in both surface and interstitial water

at all the localities (Table 2), and the saturation ratio R usually reflects those concentrations (Table 3). Measured surface water concentrations were between $0.19\text{--}35.47 \mu\text{g l}^{-1}$, $0.64\text{--}4.5 \text{ mg l}^{-1}$, $0\text{--}5.8 \mu\text{g l}^{-1}$ for CH_4 , CO_2 , and N_2O . A significant enhancement of CH_4 concentration was found at localities IV and V compared to upstream localities (I-III). Similarly a remarkable increase in the surface N_2O concentration was found at localities III-V compared to the upstream ones (I-II).

Generally, greenhouse gas concentrations measured in interstitial water were higher as compared to those from surface water, and ranged between $0.19\text{--}11,968.9 \mu\text{g l}^{-1}$, $0.85\text{--}10.82 \text{ mg l}^{-1}$, and $0.1\text{--}26.93 \mu\text{g l}^{-1}$ for CH_4 , CO_2 , and N_2O , respectively. Interstitial CH_4 concentrations at downstream localities III-V were much higher compared to those at the upstream reach (localities I-II). Significant positive correlation was found between interstitial water concentrations and surface water concentrations of CH_4 ($y = 1.5 + 0.004 x$, $R^2 = 0.61$, $F = 284.5$, $P < 0.001$) and CO_2 ($y = 0.08 + 1.77 x$, $R^2 = 0.54$, $F = 15.5$, $P < 0.01$). No relationship was observed between N_2O concentrations in the interstitial and surface water ($F = 0.5$, $P = 0.48$). Saturation ratios of CO_2 were found at levels slightly in excess of the atmospheric equilibrium leading to the supersaturation. CO_2 saturation ratios of surface and interstitial water showed only slight variation compared to the CH_4 saturation ratios.

Patterns of Emissions

The highest emissions were observed for CO_2 ($0\text{--}17.33 \text{ g m}^{-2} \text{ day}^{-1}$) followed by CH_4 ($0\text{--}167.35 \text{ mg m}^{-2} \text{ day}^{-1}$), and the lowest values reached N_2O ($0\text{--}19.69 \text{ mg m}^{-2} \text{ day}^{-1}$). Emissions of CO_2 were usually three orders of magnitude greater than CH_4 and N_2O . We found substantial spatial variations of greenhouse gas emissions along the longitudinal stream profile, when a significant enhancement of emissions for all three greenhouse gases were observed in downstream localities (IV-V) compared to upstream ones (I-III) (Table 4). Moreover, significant positive correlation was found between surface water concentrations and measured emissions of all the gases with coefficient of determination $R^2 = 0.57$ ($y = -3.83 + 1.62 x$, $F = 664.5$, $P < 0.001$), $R^2 = 0.77$ ($y = -6.06 + 3.66 x$, $F = 81.3$, $P < 0.001$), and $R^2 = 0.61$ ($y = -3.36 + 1.89 x$, $F = 24.7$, $P < 0.001$) for CH_4 , CO_2 , and N_2O , respectively (Fig. 2). Multiple linear regression with gas concentrations and flow velocity as predictor variables and gas emissions as a response variable was not significant for all monitored gases.

The seasonal contribution by CH_4 emissions to the total annual emissions was the highest during the spring and summer periods (81.9 %), while a relatively high proportion of CO_2 (65.7 %) was emitted during summer and autumn months. The contribution of N_2O emissions was the highest during summer (32.3 %) and winter (31.9 %) (Fig. 3).

On the basis of annual mean emissions of particular gasses and areas of the defined segments, approximately

Table 2. Annual means of gas concentrations in surface and interstitial waters at Sitka Stream localities (mg l⁻¹ for CO₂ and µg l⁻¹ for CH₄ and N₂O) ±standard errors.

Locality	CH ₄	CO ₂
I		
Surface	1.25±0.43	1.76±0.29
Interstitial	1.34±0.55	3.00±0.49
II		
Surface	0.46±0.20	1.13±0.18
Interstitial	0.65±0.26	1.46±0.35
III		
Surface	1.37±0.49	1.81±0.30
Interstitial	19.20±16.93	2.28±0.38
IV		
Surface	15.00 ± 6.63	2.01±0.39
Interstitial	5,196.01±2,322.28	5.80±2.16
V		
Surface	12.12±3.77	3.36±0.45
Interstitial	15.38±7.23	4.34±1.30

211 tons of greenhouse gases were emitted annually to the atmosphere from Sitka water surface with a dominant proportion being CO₂ (Table 5). If total annual emissions of each gas are converted to standard metric units (CO₂

Table 3. Annual means of saturation ratio of CH₄, CO₂, and N₂O in surface and interstitial waters at all localities ±standard errors.

Locality	CH ₄	CO ₂
I		
Surface	25.2±10.22	1.9±0.12
Interstitial	29.5±5.14	3.9±0.299
II		
Surface	9.9±5.54	1.3±0.20
Interstitial	15.4±8.94	2.1±0.61
III		
Surface	31.5±13.49	2.3±0.51
Interstitial	487.9±382.69	3.4±0.98
IV		
Surface	379.5±216.86	2.5±0.70
Interstitial	130,437.4±83,544.46	9.8±4.49
V		
Surface	246.4±81.98	4.1±0.82
Interstitial	364.9±190.73	6.2±3.14

Table 4. Mean gas emissions to the atmosphere (g m⁻²d⁻¹ for CO₂ and mg m⁻²d⁻¹ for CH₄ and N₂O) ±standard errors.

Locality	CH ₄	CO ₂	N ₂ O
I	2.39±0.96	3.68±0.71	2.83±2.01
II	0.25±0.08	1.15±0.67	0.80±0.92
III	1.41±0.91	3.28±1.87	6.63±2.48
IV	41.31±20.25	4.15±1.99	3.34±1.69
V	49.33±26.25	9.33±1.69	8.01±3.74

equivalent emissions) using GWP described in IPCC methodology [3], then CO₂ accounts for most of total annual emissions of greenhouse gases from the stream, with the second being N₂O, and the last CH₄ – for both 20- and 100-year time horizons.

Discussion

Gas Concentrations in Surface and Interstitial Waters

We found both surface and interstitial waters to be supersaturated with CH₄, CO₂, and N₂O compared to the atmosphere at all five localities, except for the N₂O concentration in the surface water at locality II. Supersaturated concentrations of CO₂ and N₂O in interstitial and surface waters are commonly reported [11, 28-35] because CO₂ is produced mainly by decomposition of organic matter, carbonate equilibrium reaction, and respiration; and N₂O by nitrification, denitrification, and coupled nitrification-denitrification as a byproduct. Dissolved CH₄ concentrations in the surface water of the stream are in agreement with earlier studies that focused on the measurement of CH₄ concentrations in rivers [36-37]. Some authors have suggested that increased concentrations of CH₄ in rivers result from lateral diffusion of stream banks and drainage of forest and agricultural soils, or from outflow of urbanized areas and sewage treatments [9, 36, 38]. Striegl et al. [8] mentioned that river CH₄ could be derived from multiple sources, including groundwater, surface water runoff, and benthic and water column microbial processing of organic carbon. The exact role of each of these processes is not yet quantified in the overall river carbon budget. Other authors [30, 39] have found, as we did, a significant increase of CH₄ in the interstitial water.

In spite of a commonly perceived view of streams as well-oxygenated habitats, we assume internal CH₄ production in hyporheic sediments as some other authors have suggested [12, 19, 39-40]. Moreover, we can exclude the possibility that CH₄ in Sitka sediments is leaked from natural gas extraction or from a geothermal and volcanic source due to the absence of these processes in the watershed of the stream. This supposition of biogenic origin of CH₄ is supported by carbon isotope analysis




Fig. 2. The relationship between atmospheric emissions and surface water concentrations for CH_4 a), CO_2 b), and N_2O c).

of CH_4 in bubbled gas from hyporheic sediments of the stream [41]. We also excluded significant contributions to drainage by wetlands and peatlands (due to absence of wetland ecosystems in the watershed of the stream), as has been reported in some cases [42-43]. Internal CH_4 production could easily be explained by anaerobic conditions occurring in hyporheic biofilm in micro-scale dimensions. Physical and chemical gradients in hyporheic sediments enable a broad spectrum of metabolic pathways to occur within small spatial scales and imply important biological consequences.

The seemingly well-oxygenated hyporheic zone contains anoxic and hypoxic pockets associated with irregularities in sediment surfaces, small pore spaces, or local deposits of organic matter, thus creating a ‘mosaic’ structure of various environments where different microbial populations can live and different microbial processes can occur simultaneously [44]. As the interstitial water in comparison to surface waters were markedly more supersaturated by all monitored gases, we suggest that they are produced in hyporheic sediments of the stream.

Levels of the gas concentrations in the surface water do not depend only on gas production in hyporheic

sediments, but they are affected by many other effects like photosynthesis, methanotrophy, respiration, or downstream transport of gases. Moreover, a part of the gases can result from drainage of soils in the watershed [43]. Since aquatic macrophytes and macroscopic algal communities were not observed in the streambed of the studied localities, we do not expect marked daily variation of CO_2 concentration in surface water due to photosynthesis (although small fluctuations could be caused by daily temperature fluctuations and microbial biofilms containing primary producers on the streambed).

Downstream localities (III-V) had higher interstitial and surface water concentrations and emissions of all the gases in comparison to upstream localities (I-II). For example, the majority of the total annual CH_4 emissions (90%) was released from the two most downstream stretches (IV-V) of the stream, representing only 1/5 of the total stream area. An ideal water stream embodies a continuum of changing physical conditions downstream with decreasing flow velocity and increasing mass of finer particles and organic matter in hyporheic sediments [45]. Therefore, production of biogenic gases increases downstream together with an amount of organic-rich sediments that enable microbial processes. High concentrations of the gases found in interstitial water of site IV (Table 2) were caused probably by very fine sediment and substantial amounts of organic matter in this locality. Such a place may be considered a ‘hot spot’ of gas production and subsequent release to surface water. Dissolved gases are repeatedly degassed from river reaches containing riffles and replenished in slower moving reaches within pools. High CH_4 concentration in the upstream part, with subsequent decline further downstream, has been reported from the United States. [36]. Therefore, greenhouse gas concentrations and emissions are more likely related to local conditions than to an accumulation of transported gases from upstream parts of a stream. The most important factor associated with river N_2O production is nitrate concentration [30, 46]. In the case of N_2O we suggest that high surface and interstitial concentrations on the downstream localities (III-V), and consequently also emissions, are due to input of nitrate from sewage water




Fig. 3. Contributions of gas emissions to the atmosphere from different seasons to total annual emissions of each gas.

Table 5. Estimation of total annual emissions of particular greenhouse gases (t yr^{-1} for CO_2 and kg yr^{-1} for CH_4 and N_2O) from Sitka Stream to the atmosphere, and relative metric units to 1 kg of CO_2 calculated using GWPs for 20- and 100-year time horizons. The percentage contribution of particular greenhouse gases to relative metric units of the total annual emission is in parenthesis.

	CH_4	CO_2	N_2O
Total annual emission	637.2	210.2	228.6
100-year GWP	21,665 (7.2%)	210,200 (70.1%)	68,123 (22.7%)
20-year GWP	54,799 (16.8%)	210,200 (64.4%)	61,265 (18.8%)

outflows from the town of Šternberk (located between localities II and III), and due to drainage of agricultural land on downstream parts of the watershed. Generally, N_2O concentrations are related to the amount of flooding and leaching of N from a catchment, especially if the catchment contains agricultural areas.

Gas Emissions to the Atmosphere

The range of measured greenhouse gas emissions presented in this study was in accordance with most other studies from rivers. Average values of CH_4 emissions on Sitka Stream are similar to values reported for small streams in Arizona [39] and for rivers in Oregon [36]. The annual mean CO_2 emissions to the atmosphere from Sitka were also similar to those found on temperate rivers of the United States [47]. Wilcock and Sorrell [13] found, as we did, that rivers could emit significant amounts of N_2O .

Generally, gas emissions comprise a flux mediated by submerged plants, an ebullition from sediments, and a diffusive flux from the water surface [48]. Since we did not record any primary producers in the streambed, contributions by plant-mediated flux to overall emissions is most probably not a case or is negligible in Sitka. However, we observed gas bubbles released from the sediment, specifically at locality IV. Although an ebullition is rarely captured by the floating chamber technique [49], the preference to use this method was accurate. The methods based on calculation models account for only diffusive flux, and we observed ebullition at locality IV. The rate of gas diffusion into the atmosphere depends on several factors: partial pressure of a particular gas in the atmosphere and its concentration in water, water temperature, and further on molecular diffusion coefficient of a particular gas, water depth, and flow velocity [38, 50]. Accordingly, a temperature increase that reduces gas solubility could lead to higher emission rates during the summer. Furthermore, streams with higher flow velocity have higher gas exchange coefficients compared to wetlands or large rivers. Nevertheless, multiple linear regression with gas concentrations and flow velocity as predictor variables and gas emissions as a response variable was not significant for all monitored gases, but simple linear regression between gas concentrations and emissions was significant for all monitored gases (Fig. 2), as was found on three rivers in China [9]. This contradiction could be caused by highly turbulent flows influencing gas exchange on the air-water interface, which

make flow velocity less predictive of gas emissions from the stream.

The CH_4 emissions from rivers usually reach the highest maximum at the end of summer [51-52]. The positive effect of temperature on CH_4 production was also observed for river sediments in laboratory conditions [53]. Besides the direct effect of temperature, the positive correlation of CH_4 emissions with temperature could be related to the prolonged growth season of macrophytes, which provides the organic substrates by primary production [54]. The CH_4 production rate of sediment grows with increasing temperature up to 40°C , while it is inhibited by temperatures below 12°C [55]. The same acceleration of CH_4 production inferred from positive correlation of dissolved CH_4 with temperature was recorded by Yang et al. [9]. Therefore, the increase of dissolved CH_4 concentrations in surface and interstitial water corresponds with the hypothesis that river sediment is probably an important CH_4 source for surface water in warmer months. Indeed, it has been shown that summer months are characterized by increased CH_4 surface concentrations in many European rivers [56], and a significant positive linear correlation between dissolved CH_4 concentrations and CH_4 emissions was observed by Yang et al. [9]. If we consider the seasonal distribution of gas emissions, it is clear, in concordance with the above-mentioned relations, that the majority of CH_4 and CO_2 emissions were released during warmer periods of the year. The highest N_2O emissions from Sitka were recorded during summer and winter months, and an equivocal seasonal pattern was obtained. But the 31.9% contribution of N_2O emissions during winter to the total annual budget of the stream is not surprising. Nitrate concentrations (the main predictor of N_2O emissions) usually increase during winter and decline during growing seasons because the export of nitrate from the watershed is controlled by plant uptake of inorganic N, which is lower during the non-growing season [57].

Based on our results, we suggest that CH_4 and N_2O emissions contribute considerably to total greenhouse gas emissions from the stream. We found that CH_4 emissions represent 0.8% of the mass of carbon released to the atmosphere from Sitka. Although the total CH_4 mass emitting from the stream is three orders of magnitude less than total CO_2 mass, CH_4 emissions play a considerable role in the climate impact of the stream due to its GWP. The total annual CH_4 emissions contribute 7.2% to the total annual emission of all gases (converted to CO_2

equivalents) from the stream for the 100-year time-horizon, but as much as 16.8% for the 20-year horizon. The higher CH₄ contribution in the 20-year horizon was expected due to the short residence time of CH₄ in the atmosphere. Total annual N₂O emissions contribute 22.7% to the total annual emissions of all gases, converted to CO₂ equivalents, from the stream for the 100-year time-horizon, and 18.8% for the 20-year horizon. These rates indicate that CH₄ and N₂O emissions should not be omitted in greenhouse gas budgets of rivers and streams, especially in a short time scale. Moreover, Campeau and Del Giorgio [58] estimated that greenhouse gas emissions (CO₂ equivalent) from boreal rivers and streams may increase markedly, driven mostly by a steep increase in the contribution of CH₄ under plausible scenarios of climate change over the next 50 years. Nevertheless, our results should be interpreted carefully with the up-scaling of emissions data for the whole riverine systems—especially because our dataset is spatially and temporally restricted. Moreover, we did not include greenhouse gas emissions from parafluvial zones, and the riverine (landscape) level fluxes can be rather different from those gathered from limited local points.

Conclusions

CH₄, CO₂, and N₂O are produced in hyporheic sediments of Sitka Stream. The longitudinal pattern of greenhouse gas concentrations and emissions along streams is heterogeneous and reflects a spatial distribution of physicochemical characteristics. Although concentrations and emission rates of the greenhouse gases did not show increasing trends along the stream, sampling sites located downstream indicated higher values of concentrations and emission rates than upstream localities, and such longitudinally located sampling sites are necessary for an emission estimate. Therefore, we suggest that our approach of the direct measurement on a set of sampling sites along a water stream during a season is proper for an estimation of total annual emissions of greenhouse gases from river ecosystems.

The total annual emissions of CO₂ to the atmosphere was the highest for all the monitored greenhouse gases, and the total annual emissions of CH₄, CO₂, and N₂O were estimated to be 637 kg, 210 t, and 229 kg, respectively. However, counting the global warming potential, the total annual contributions of CH₄ and N₂O to total annual greenhouse gas emissions from Sitka amount to nearly 30% for the 100-year horizon. Therefore, CH₄ and N₂O should not be omitted in an evaluation of greenhouse gas emissions from running waters. Our results utilize directly measured gas concentration and emission data to contribute to recent assessments [59–62] of the importance of inland waters in regional and global carbon cycles, and quantify the importance of CH₄, CO₂, and N₂O emissions in the carbon and nitrogen balance of a small temperate water stream system.

Acknowledgements

Our research was supported by a grant No. NPV II 2B06101.

References

1. IPCC Summary for Policymakers. In: STOCKER T.F., QIN D., PLATTNER G.-K., TIGNOR M., ALLEN S.K., BOSCHUNG J., NAUELS A., XIA Y., BEX V., MIDGLEY P.M. (eds.) *Climate Change 2013: The Physical Science Basis. Contribution of WGI to the 5AR of the IPCC*. Cambridge University Press, **2013**.
2. FORSTER P., RAMASWAMY V., ARTAXO P., BERNTSEN T., BETTS R., FAHEY D.W., HAYWOOD J., LEAN J., LOWE D.C., MYHRE G., NGANGA J., PRINN R., RAGA G., SCHULZ M., VAN DORLAND R. Changes in Atmospheric Constituents and in Radiative Forcing. In: SOLOMON S., QIN D., MANNING M., CHEN Z., MARQUIS M., AVERYT K.B., TIGNOR M., MILLER H.L. (eds.) *Climate Change 2007: The Physical Science Basis*. Cambridge University Press, **2007**.
3. MYHRE G., SHINDELL D., BRÉON F.-M., COLLINS W., FUGLESTVEDT J., HUANG J., KOCH D., LAMARQUE J.-F., LEE D., MENDOZA B., NAKAJIMAT., ROBOCK A., STEPHENS G., TAKEMURA T., ZHANG H. Anthropogenic and Natural Radiative Forcing. In: *Climate Change 2013: The Physical Science Basis. Contribution of WGI to the 5AR of the IPCC* [STOCKER T.F., QIN D., PLATTNER G.-K., TIGNOR M., ALLEN S.K., BOSCHUNG J., NAUELS A., XIA Y., BEX V., MIDGLEY P.M. (eds.)]. Cambridge University Press, **2013**.
4. FRANKIGNOULLE M., MIDDELBURG J.J. Biogases in tidal European estuaries: the BIOGEST project. *Biogeochemistry* **59**, 1, **2002**.
5. PULLIAM W.M. Carbon dioxide and methane exports from a southeastern floodplain swamp. *Ecol. Monogr.* **63**, 29, **1993**.
6. WHALEN S.C. Biogeochemistry of methane exchange between natural wetlands and the atmosphere. *Environ. Eng. Sci.* **22** (1), 73, **2005**.
7. NYAMADZAWO G., WUTA M., NYAMANGARA J., REES R.M., SMITH J.L. The effects of catena positions on greenhouse gas emissions along a seasonal wetland (dambo) transect in tropical Zimbabwe. *Arch. Acker. Pfl. Boden.* **61**, 203, **2014**.
8. STRIEGL R.G., DORNBLASER M.M., MCDONALD C.P., ROVER J.R., STETS E.G. Carbon dioxide and methane emissions from the Yukon River system. *Global. Biogeochem. Cycles.* **26**, 1, **2012**.
9. YANG L.B., LI X.J., YAN W.J., MA P., WANG J.N. CH₄ concentrations and emissions from three rivers in the Chao-hu Lake Watershed in Southeast China. *J. Integr. Agric.* **11**, 665, **2012**.
10. GROFFMAN P.M., BOULWARE N.J., ZIPPERER W.C., POUYAT R.V., BAND L.E., COLOSIMO M.F. Soil nitrogen cycle processes in urban riparian zones. *Environ. Sci. Tech.* **36**, 4547, **2002**.
11. COLE J.J., CARACO N.F. Emissions of Nitrous Oxide (N₂O) from a Tidal, Freshwater River, the Hudson River, New York. *Environ. Sci. Technol.* **35**, 991, **2001**.
12. SANDERS I.A., HEPPELL C.M., COTTON J.A., WHARTON G., HILDREW A.G., FLOWERS E.J., TRIMMER M. Emissions of methane from chalk streams has potential

- implications for agricultural practices. *Freshw. Biol.* **52**, 1176, **2007**.
13. WILCOCK R.J., SORRELL B.K. Emissions of greenhouse gases CH₄ and N₂O from low-gradient streams in agriculturally developed catchments. *Water. Air. Soil. Pollut.* **188**, 155, **2008**.
 14. SAARNIO S., WINIWARTER W., LEITA J. Methane release from wetlands and watercourses in Europe. *Atmos. Environ.* **43**, 1421, **2009**.
 15. DOWNING J.A., COLE J.J., DUARTE C.A., MIDDELBURG J.J., MELACK J.M., PRAIRIE Y.T., KORTELAINEN P., STRIEGL R.G., MCDOWELL W.H., TRANVIK L.J. Global abundance and size distribution of streams and rivers. *Inland Waters* **2**, 229, **2012**.
 16. RULÍK M., ČÁP L., HLAVÁČOVÁ E. Methane in the hyporheic zone of a small lowland stream (Sitka, Czech Republic). *Limnologica* **30**, 359, **2000**.
 17. HLAVÁČOVÁ E., RULÍK M., ČÁP L. Anaerobic microbial metabolism in hyporheic sediment of a gravel bar in a small lowland stream. *River. Res. Appl.* **21**, 1003, **2005**.
 18. CUPALOVÁ J., RULÍK M. Bacterial community analysis in river hyporheic sediments – the influence of depth and particle size. *Acta Universitatis Carolinae Environmentalica* **21**, 47, **2007**.
 19. BURIÁNKOVÁ I., BRABLCOVÁ L., MACH V., HÝBLOVÁ A., BADUROVÁ P., CUPALOVÁ J., ČÁP L., RULÍK M. Methanogens and methanotrophs distribution in the hyporheic sediments of a small lowland stream. *Fundam. Appl. Limnol.* **181**, 87, **2012**.
 20. HLAVÁČOVÁ E., RULÍK M., ČÁP L., MACH V. Greenhouse gases (CO₂, CH₄, N₂O) emissions to the atmosphere from a small lowland stream. *Arch. Hydrobiol.* **165**, 339, **2006**.
 21. ANDERSON B., BARTLETT K., FROLKING S., HAYHOE K., JENKINS J., SALAS W. Methane and Nitrous Oxide Emissions From Natural Sources. Office of Atmospheric Programs US EPA, **2010**.
 22. TRIMMER M., GREY J., HEPPELL C.M., HILDREW A.G., LANSDOWN K., STAHL H., YVON-DUROCHER G. River bed carbon and nitrogen cycling: State of play and some new directions. *Sci. Total. Environ.* **434**, 143, **2012**.
 23. WIESENBURG D.A., GUINASSO N.L. Equilibrium solubilities of methane, carbon monoxide, and hydrogen in water and sea water. *J. Chem. Eng. Data.* **24**, 356, **1979**.
 24. WEISS R.F., PRICE B.A. Nitrous oxide solubility in water and seawater. *Mar. Chem.* **8**, 347, **1980**.
 25. WEISS R.F. Carbon dioxide in water and seawater: the solubility of a non-ideal gas. *Mar. Chem.* **2**, 203, **1974**.
 26. CRILL P., BARTLETT K.B., HARRISS R.C., GORHAM E., VERRY E.S., SEBACHER D.I., MADZAR L., SANNER W. Methane flux from Minnesota peatlands. *Glob. Biogeochem. Cyc.* **2**, 371, **1988**.
 27. PIEROBON E., BOLPAGNI R., BARTOLI M., VIAROLI P. Net primary production and seasonal CO₂ and CH₄ fluxes in a *Trapa natans* L. meadow. *J. Limnol.* **69** (2), 225, **2010**.
 28. COLE J.J., CARACO N.F., KLING G.W., KRATZ T.K. Carbon Dioxide Supersaturation in the Surface Waters of Lakes. *Science* **265**, 1568, **1994**.
 29. RIVETT M.O., BUSS S.R., MORGAN P., SMITH J.W.N., BEMMENT C.D. Nitrate attenuation in groundwater: A review of biogeochemical controlling processes. *Water Research* **42**, 4215, **2008**.
 30. WANG D.Q., CHEN Z.L., SUN W.W., HU B.B., XU S.Y. Methane and nitrous oxide concentration and emission flux of Yangtze Delta plain river net. *Sci. China. Ser. B.* **52**, 652, **2009**.
 31. BUTMAN D., RAYMOND P.A. Significant efflux of carbon dioxide from streams and rivers in the United States. *Nat. Geosci.* **4**, 839, **2011**.
 32. ROSAMOND M.S., THUSS S.J., SCHIFF S.L. Dependence of riverine nitrous oxide emissions on dissolved oxygen levels. *Nat. Geosci.* **5**, 715, **2012**.
 33. HINSHAW S.E., DAHLGREN R.A. Dissolved nitrous oxide concentrations and fluxes from the eutrophic San Joaquin River, California. *Environ. Sci. Technol.* **47** (3), 1313, **2013**.
 34. RAYMOND P.A., HARTMANN J., LAUERWALD R., SOBEK S., MCDONALD C., HOOVER M., BUTMAN D., STRIEGL R., MAYORGA E., HUMBORG C., KORTELAINEN P., DÜRR H., MEYBECK M., CIAIS P., GUTH P. Global carbon dioxide emissions from inland waters. *Nature* **503**, 355, **2013**.
 35. RASERA M.F.F.L., KRUSCHE A.V., RICHEY J.E., BALLESTER M.V.R., VICTÓRIA R.L. Spatial and temporal variability of pCO₂ and CO₂ efflux in seven Amazonian Rivers. *Biogeochemistry* **116**, 241, **2013**.
 36. LILLEY M.D., DE ANGELIS M.A., OLSON J.E. Methane concentrations and estimated fluxes from Pacific Northwest rivers. *Mitt. Internat. Verein. Limnol.* **25**, 187, **1996**.
 37. ZHANG G., ZHANG J., LIU S., REN J., XU J., ZHANG F. Methane in the Changjiang (Yangtze River) Estuary and its adjacent marine area: riverine input, sediments release and atmospheric fluxes. *Biogeochemistry* **91**, 71, **2008**.
 38. DE ANGELIS M.A., LILLEY M.D. Methane in surface waters of Oregon estuaries and rivers. *Limnol. Oceanogr.* **33**, 716, **1987**.
 39. JONES J.B., HOLMES R.M., FISCHER S.G., GRIMM N.B., GREENE D.M. Methanogenesis in Arizona, USA dryland streams. *Biochemistry* **31**, 155, **1995**.
 40. STOREY R.G., FULTHORPE R.R., WILLIAMS D.D. Perspectives and predictions on the microbial ecology of the hyporheic zone. *Freshw. Biol.* **41**, 119, **1999**.
 41. MACH V., BLASER M.B., CLAUS P., CHAUDHARY P.P., RULIK M. Methane production potentials, pathways, and communities of methanogens in vertical sediment profiles of river Sitka. *Front. Microbiol.* **6**, 506, **2015**.
 42. HOPE D., PALMER S.M., BILLET M.F., DAWSON J.J.C. Variations in dissolved CO₂ and CH₄ in a first-order stream and catchment: an investigation of soil-stream linkages. *Hydrol. Process.* **18**, 3255, **2004**.
 43. SILVENNOINEN H., LIIKANEN A., RINTALA J., MARTIKAINEN P.J. Greenhouse gas fluxes from the eutrophic Temmesjoki River and its Estuary in the Liminganlahti Bay (the Baltic Sea). *Biogeochemistry* **90**, 193, **2008**.
 44. BAKER M.A., DAHM C.N., VALLET H.M. Acetate retention and metabolism in the hyporheic zone of a mountain stream. *Limnol. Oceanogr.* **44**, 1530, **1999**.
 45. BOUWMAN A.F., BIERKENS M.F.P., GRIFFIOEN J., HEFTING M.M., MIDDELBURG, J.J., MIDDELKOOP H., SLOMP C.P. Biogeosciences Nutrient dynamics, transfer and retention along the aquatic continuum from land to ocean: towards integration of ecological and biogeochemical models. *Biogeosciences* **10**, 1, **2013**.
 46. BEAULIEU J.J., ARANGO C.P., TANK J.L. The effects of season and agriculture on nitrous oxide production in headwater streams. *J. Environ. Qual.* **38**, 637, **2009**.
 47. BUTMAN D., RAYMOND P.A. Significant efflux of carbon dioxide from streams and rivers in the United States. *Nature Geoscience*, NGE01294, **2011**.
 48. BASTVIKEND., COLE J., PACE M., TRAVNIK L. Methane emissions from lakes: Dependence on lake characteristics, two regional assessments, and a global estimate. *Global Biogeochem. Cycles* **18**, 1, **2004**.

49. MAECK A., HOFMANN H., LORKE A. Pumping methane out of aquatic sediments – ebullition forcing mechanisms in an impounded river. *Biogeosciences* **11**, 2925, **2014**.
50. SEGERS R. Methane production and methane consumption: a review of processes underlying wetland methane fluxes. *Biogeochemistry* **41**, 23, **1998**.
51. YVON-DUROCHER G., MONTOYA J.M., WOODWARD G., JONES J.I., TRIMMER M. Warming increases the proportion of primary production emitted as methane from freshwater mesocosms. *Glob. Chang. Biol.* **17**, 1225, **2011**.
52. ORTIZ LLORENTE M. J., ALVAREZ COBELAS M. Comparison of biogenic methane emissions from unmanaged estuaries, lakes, oceans, rivers and wetlands. *Atmos. Environ.* **59**, 328, **2012**.
53. SHELLEY F., ABDULLAHI F., GREY J., TRIMMER M. Microbial methane cycling in the bed of a chalk river: oxidation has the potential to match methanogenesis enhanced by warming. *Freshw. Biol.* **59**, fwb.12480, **2014**.
54. NATCHIMUTHU S., SELVAM B.P., BASTVIKEN D. Influence of weather variables on methane and carbon dioxide flux from a shallow pond. *Biogeochemistry*. **119**, 403, **2014**.
55. YANG S.S. Methane production in river and lake sediments in Taiwan. *Environ Geochem. Health* **20**, 245, **1998**.
56. MIDDELBURG J.J., NIEUWENHUIZE J., IVERSEN N., HØGH N., DE WILDE H., HELDER W., SEIFERT R., CHRISTOF O. Methane distribution in tidal estuaries. *Biogeochemistry* **59**, 95, **2002**.
57. LEE K.E., LORENZ D.L., PETERSEN J.C., GREENE J.B. Seasonal patterns in nutrients, carbon, and algal responses in wadeable streams within three geographically distinct areas of the United States, 2007–08: U.S. Geological Survey Scientific Investigations Report **2012**.
58. CAMPEAU A., DEL GIORGIO P.A. Patterns in CH₄ and CO₂ concentrations across boreal rivers: Major drivers and implications for fluvial greenhouse emissions under climate change scenarios. *Glob. Change. Biol.* **20**, 1075, **2014**.
59. COLE J.J., PRAIRIE Y.T., CARACO N.F., MCDOWELL W.H., TRANVIK L.J., STRIEGL R.G., DUARTE C.M., KORTELAJAINEN P., DOWNING J.A., MIDDELBURG J.J., MELACK J. Plumbing the global carbon cycle: integrating inland waters into the terrestrial carbon budget. *Ecosystems* **10**, 172, **2007**.
60. BATTIN T.J., LUYSSAERT S., KAPLAN L.A., AUFDENKAMPE A.K., RICHTER A., TRANVIK L.J. The boundless carbon cycle. *Nat. Geosci.* **2**, 598, **2009**.
61. TRANVIK L.J., DOWNING J.A., COTNER J.B., LOISELLE S.A., STRIEGL R.G., BALLATORE T.J., DILLON P., FINLAY K., FORTINO K., KNOLL L.B., KORTELAJAINEN P.L., KUTSER T., LARSEN S., LAURION I., LEECH D.M., MCCALLISTER S.L., MCKNIGHT D.M., MELACK J.M., OVERHOLT E., PORTER J.A., PRAIRIE Y., RENWICK W.H., ROLAND F., SHERMAN B.S., SCHINDLER D.W., SOBEK S., TREMBLAY A., VANNI M.J., VERSCHOOR A.M., WACHENFELDT E., WEYHENMEYER G.A. Lakes and reservoirs as regulators of carbon cycling and climate. *Limnol. Oceanogr.* **54**, 2298, **2009**.
62. ALIN S.R., RASERA M.F.F.L., SALIMON C.I., RICHEY J.E., HOLTGRIEVE G.W., KRUSCHE A.V., SNIDVONGS A. Physical controls on carbon dioxide transfer velocity and flux in low-gradient river systems and implications for regional carbon budgets. *J. Geophys. Res.-Biogeo.* **116**, G01009, **2011**.