

Original Research

# The Effect of Some Indoor Ornamental Plants on CO<sub>2</sub> Levels During the Day

Hakan Sevik<sup>1</sup>, Mehmet Cetin<sup>2\*</sup>, Kerim Guney<sup>3</sup>, Nur Belkayali<sup>2</sup>

<sup>1</sup>Kastamonu University, Faculty of Engineering and Architecture, Department of Environmental Engineering, 37150 Kastamonu/Turkey

<sup>2</sup>Kastamonu University, Faculty of Engineering and Architecture, Department of Landscape Architecture, 37150 Kastamonu/Turkey

<sup>3</sup>Kastamonu University, Faculty of Forestry, Department of Forest Engineering, 37150 Kastamonu/Turkey

Received: 18 June 2017

Accepted: 2 August 2017

## Abstract

The aim of the present study is to determine the effect of yucca (*Yucca elephantipes* Regel), dieffenbachia (*Dieffenbachia amoena* Gentil), and spathiphyllum (*Spathiphyllum floribundum* Schott) as common types of indoor plants on CO<sub>2</sub> levels in the environment. The study was conducted in a closed environment where air inlet/outlet was absent. As a result, the plants were found to affect the level of CO<sub>2</sub> in the environment to different extents; and while Dieffenbachia began to respire at around 13:00 hrs, yucca and spathiphyllum continued to photosynthesize until 19:00. While dieffenbachia and spathiphyllum could not lower the CO<sub>2</sub> level in the environment to below 500 ppm, yucca could decrease it to 475 ppm in a day. The results of the study demonstrated that CO<sub>2</sub> levels in photosynthesis during the day were 8.3, 5.8, and 1.4 times more in yucca, spathiphyllum, and dieffenbachia, respectively, in comparison with CO<sub>2</sub> levels released through respiration.

**Keywords:** indoor plants, CO<sub>2</sub>, leaf area, ornamental plants, photosynthesis, respiration

## Introduction

Due to increased construction and changes in recent living conditions, people spend at least 80% of their lives indoors [1-3]. This situation has led to many studies on indoor air pollution and increased of CO<sub>2</sub> levels in line with the metabolic activities of living things. As a result of the increased carbon dioxide level in the environment, fatigue, perception difficulty, and sleepiness are experienced. Carbon dioxide also causes

various problems that lead to poor performance – the reason for which cannot be determined. When the CO<sub>2</sub> level in an environment increases up to 1,000 ppm, headache, vertigo, fatigue, concentration problems, and smell disorders are experienced, while itchy nose and throat, nasal discharge, cough, and eye discharge appear when it exceeds 1,500 ppm [4-6].

According to the U.S. Environmental Protection Agency (EPA), the maximum carbon dioxide level indoors should be 1,000 ppm – even in crowded environments such as schools and conference halls [5]. However, relevant studies determined that the carbon dioxide level indoors was higher than this maximum level in many

\*e-mail: mcerin@kastamonu.edu.tr

environments and even exceeded 1,900 ppm in student residence halls [7]. It reached 3,700 ppm in exam rooms [5] and 5,400 ppm [8] in classrooms.

Indoor plants are living things and have several functions. For example, they reduce air pollution [9-13], increase work productivity [14], relax people psychologically, and reduce stress and negative feelings [5-6, 15-16]. Another reason why people want to grow plants indoors is that they have an effect on carbon dioxide levels. Plants are known to photosynthesize and thus reduce the level of CO<sub>2</sub> in an environment [5-6]. However, plants are living things and respire when the conditions are not suitable for them to photosynthesize, thus increasing the CO<sub>2</sub> level in an environment. However, the number of studies on how plants affect CO<sub>2</sub> levels depending on the conditions in the environment is limited.

This present study aims to determine how some indoor plants affect the CO<sub>2</sub> level in an environment at different times of the day.

### Material and Methods

The present study aims to determine the effect of some indoor ornamental plants on the CO<sub>2</sub> level in a closed environment. *Spathiphyllum floribundum* Schott), *yucca (Yucca elephantipes* Regel), and *dieffenbachia (Dieffenbachia amoena* Gentil) were used in this study. These plants have different ecological needs and physical characteristics (leaf area, type of stem, etc.) and are the most common indoor plants around the world.

The plants were placed in a glass wall (0.7 × 0.7 × 1 m) with a volume of about 0.5 m<sup>3</sup>, which was not air-permeable, and the measurements were taken using the Extech Desktop Indoor Air Quality CO<sub>2</sub> Datalogger. The glass wall was placed in the south of the building so that it received plenty of daylight. It received direct sunlight between 07:00 and 11:00 and was illuminated until around 17:00. In the area where the study was conducted, the sun rose at around 05:05 and went down at around 20:30.

After the plants were placed in the glass wall, a CO<sub>2</sub> measuring device was set to measure the level of CO<sub>2</sub> every five minutes. The plants were placed in the glass wall between 13:00 and 14:00, and the CO<sub>2</sub> level within the glass was increased. Data were obtained from this time on; however, only those measurements taken after 04:00 were considered.

The plants were kept inside the glass wall for at least five days. This study was conducted June to July, when daylight is longer. The results obtained were evaluated using graphs.

### Results and Discussion

To better evaluate the obtained data we used hourly values, and graphs showing the effect of *dieffenbachia* on the CO<sub>2</sub> level during the day are presented in Fig. 1.

Fig. 1 shows that the 1,868 ppm CO<sub>2</sub> level at 05:02 quickly decreased during the day, and this decline continued until 13:02. It went down from 1,868 to 766 ppm at 13:02. After this point, the CO<sub>2</sub> level kept increasing until 05:02 the next day and reached 1,506 ppm. The 1,506 ppm CO<sub>2</sub> level measured at 05:02 the second day declined to 670 ppm at 12:02 the same day. This situation indicates that *dieffenbachia* decreased the CO<sub>2</sub> level very quickly. The 1,868 ppm CO<sub>2</sub> level at 05:02 on the first day declined to 766 ppm at 13:02. In other words, it decreased by 1,102 ppm within eight hours. However, the increase in CO<sub>2</sub> level resulting from respiration during hours with insufficient daylight was also fairly high. The 766 ppm CO<sub>2</sub> level measured at 13:02 reached 1,506 ppm at 05:02 the next day. In other words, it increased by 740 ppm when the daylight was insufficient for photosynthesis.

When the results were evaluated, *dieffenbachia* was found to decrease CO<sub>2</sub> from 1,868 to 1,506 ppm within 24 hours. So the CO<sub>2</sub> level declined by only 362 ppm in total during the day. Out of the 1,102 ppm carbon dioxide used during the day, 766 ppm (nearly 69% of it) was used when daylight was insufficient for photosynthesis.

A graph showing the effect of *spathiphyllum* on CO<sub>2</sub> level during the day is presented in Fig. 2, which shows that *spathiphyllum* began to decrease the CO<sub>2</sub> level in the environment at 04:45, which continued until 18:45. During this time, the CO<sub>2</sub> level declined from 2,748 to 1,564 ppm; in other words, it declined by 1,184 ppm. After that, the CO<sub>2</sub> level initially took a horizontal course and then increased, which continued until 05:45 the next




Fig. 1. Effect of *dieffenbachia* on CO<sub>2</sub> levels during the day.




Fig. 2. Effect of spathiphyllum on CO<sub>2</sub> levels during the day.




Fig. 3. Effect of yucca on CO<sub>2</sub> levels during the day.

day, reaching 1,769 ppm. A graph showing the effect of yucca on CO<sub>2</sub> level during the day is presented in Fig. 3.

The starting CO<sub>2</sub> level for Yucca was measured at 3,322 ppm at 04:10, which declined quickly and hit 732 ppm at 19:10. In other words, a decline of approximately 2,590 ppm was observed. The CO<sub>2</sub> level began to increase after 19:10 and continued until 06:10 the next day, reaching 1,036 ppm. Thus, the total increase was 304 ppm. Considering the 24-hour performance of the plant starting at 05:10, a decline from 3,310 to 732 ppm was seen and an increase from 732 to 1,042 ppm was observed within a day. The CO<sub>2</sub> level that declined by 2,578 ppm during the day because of photosynthesis increased by 310 ppm with respiration during hours when sunlight was insufficient.

Based on these results, all the plants were observed to photosynthesize between 06:00 and 12:00, when the daylight was sufficient, and thus they decreased the CO<sub>2</sub> level. However, each plant had a different effect on the decline of CO<sub>2</sub>. Analysis of the relevant graphs could be misleading because CO<sub>2</sub> level is one of the factors affecting the rate of photosynthesis, and ensuring that all the plants were at the same CO<sub>2</sub> level when the experiment started was impossible. However, the graphs provide important results about certain aspects.

Results from the study show that the plants began to increase CO<sub>2</sub> levels at different hours during the day. Dieffenbachia began to increase the CO<sub>2</sub> level, or respire, at 13:00, while spathiphyllum and yucca did so at 19:00. All the plants were taken to the same place in the experiment, and all the experiments were conducted in a completely cloudless sunny weather. Thus, the duration of illumination and daylight in the environment can be considered the same (a difference of a few minutes was noted as days became shorter or longer). However, the durations of photosynthesis and respiration were fairly different. Under uniform light conditions, dieffenbachia respired while spathiphyllum and yucca photosynthesized.

Another striking point was the ratio of the CO<sub>2</sub> level used in photosynthesis during the day to the CO<sub>2</sub> level released during hours when daylight was insufficient. In the evaluations conducted on one particular day, dieffenbachia was found to cause a 1,102 ppm decrease in CO<sub>2</sub> level during the day but released 766 ppm when daylight was insufficient. Spathiphyllum caused a decrease of 1,184 ppm but released 205 ppm during hours when daylight was insufficient. Yucca caused a decrease of 2,578 ppm through photosynthesis during the day but released 310 ppm during hours when daylight was insufficient.

The ratio of the CO<sub>2</sub> level used in photosynthesis during the day to the CO<sub>2</sub> level released through respiration during hours when daylight was insufficient was 8.3 in yucca, 5.8 in spathiphyllum, and 1.4 in dieffenbachia. These figures demonstrate the importance of choosing plants according to the conditions in the environment.

One of the most important factors affecting the rate of photosynthesis is CO<sub>2</sub> level. In this regard, several studies have focused on agricultural plants [17]. In the present

study, the CO<sub>2</sub> level was first kept at 2,000 ppm and not considered a determining factor. However, long-term measurements were performed on the plants to determine their behavior when CO<sub>2</sub> level declined.

As we made a general evaluation of the plant types used in the study, the CO<sub>2</sub> level for dieffenbachia, which was 1,868 ppm at 05:02, declined very quickly on the first day; however, it remained stable between 15:00 and 650 ppm during the last days. This indicates that the CO<sub>2</sub> level in the environment affects the rate of photosynthesis considerably. In spathiphyllum, the CO<sub>2</sub> level was 2,977 ppm at 15:45 on the first day and 1,699 ppm the next day at the same hour. The CO<sub>2</sub> level that continued to decline in a regular course remained between 600 and 1,100 ppm during the last three days. With Yucca, the 2,782 ppm CO<sub>2</sub> level at 10:30 on the first day declined to 502 ppm at 10:30 the next day, within 24 hours, and fell to its lowest level at 457 ppm. Then it remained at between 470 and 1,200 ppm. These levels indicate that yucca can make the CO<sub>2</sub> level much lower than other types of plants can.

Fig. 2 illustrates that the CO<sub>2</sub> level generally fluctuated within a certain range during the day. A striking point was the difference in the lowest CO<sub>2</sub> levels. Although they differed, the highest levels could be proportional to the amount of leaves in plants. However, the lowest CO<sub>2</sub> level indicates that plants do not photosynthesize under a certain level, depending on the type of plant. The graph shows that the lowest CO<sub>2</sub> level was that in yucca, followed by spathiphyllum and dieffenbachia.

The results from the study demonstrate that all the plants decreased the CO<sub>2</sub> level in the environment (depending on daylight conditions during the day) and increased it when light was absent. However, dieffenbachia was found to start increasing the CO<sub>2</sub> level, or photosynthesize, at 13:00, while spathiphyllum and yucca started at 19:00. This indicates that dieffenbachia respire while spathiphyllum and yucca photosynthesize under the same light conditions, and the latter two could photosynthesize even in environments that have less daylight. Besides, the ratio of the CO<sub>2</sub> level used through photosynthesis during the day to the CO<sub>2</sub> level released through respiration differed, which was 8.3 in yucca, 5.8 in spathiphyllum, and 1.4 in dieffenbachia.

Similar results were obtained from studies that aimed to determine the effect of plants on the CO<sub>2</sub> level in an environment. Cetin and Sevik [6] indicated that *Ficus elastica* caused a decrease of 2,216 ppm, *Yucca massengena* 2,579 ppm, *Ocimum basilicum* 401 ppm, *Sinningia speciosa* 725 ppm, and *Codiaeum variegatum* 401 ppm. Another study found that *Schefflera arboricola* caused a decrease of 1,252 ppm, *Fuchsia magellanica* 252 ppm, and *Ficus benjamina* 657 ppm [18].

Cetin and Sevik [6] pointed out that *Codiaeum variegatum*, *Ficus elastica*, and *Yucca massengena* decreased the CO<sub>2</sub> level even with little light in the environment, while *Sinningia speciosa* and *Ocimum basilicum* increased the CO<sub>2</sub> level during the same time. Similarly, Sevik et al. [19] stated that *Fuchsia*

*magellanica* started to respire at 15:28, while *Ficus elastica* and *Yucca massengena* continued to photosynthesize until around 17:00. This phenomenon could be associated with the anatomy of the plants [20-21]. Kacar et al. [22] indicated that the amount of light needed for photosynthesis changes according to type of plant. For example the rate of photosynthesis in *Asarum caudatum* reached its highest level when the light condition was 200 μmol m<sup>-2</sup>s<sup>-1</sup>, while the highest photosynthesis rate in *Atriplex triangularis* was reached with a light condition of 1,700 μmol m<sup>-2</sup>s<sup>-1</sup>.

The results from the study demonstrate that plants kept in the same environment react to conditions in the environment in different ways. While they cannot decrease the CO<sub>2</sub> level in the environment to below 500 ppm during the day, yucca can decrease it to lower degrees (up to 475 ppm) compared with other plant types. However, this number is above the CO<sub>2</sub> level in the atmosphere [6, 20, 23-24]. Nevertheless, a study found that a CO<sub>2</sub> level of around 391 ppm during the day and 422 ppm at night in the winter months remained at 148 ppm during the day and 229 ppm at night in the summer months [21, 23-25].

That plants affect the CO<sub>2</sub> level in an environment, besides being used for aesthetic purposes – especially in landscaping – is a known fact [25]. Relevant studies indicate that a beech tree with a leaf area of 1,600 m<sup>2</sup> can meet the oxygen needs of 10 people [18, 26]. Tarran et al. [27] stated that the presence of plants in the environment decreases the CO<sub>2</sub> level in offices with air conditioners by 10%, and in environments with natural ventilation by 25%. However, determining the plants to be utilized with respect to the conditions of the environment is necessary for more effective use. Studies on this subject are still very limited.

The present study aimed to find out the effects of certain plants on the CO<sub>2</sub> level in an environment. It is known that green plants photosynthesize with sufficient light and that the CO<sub>2</sub> level in an environment declines as a result of photosynthesis [27]. However, the effect of plants on the quality of air in an environment is not limited to regulation of CO<sub>2</sub> levels. Several studies have demonstrated that plants decrease the sulfur level in indoor air [28] and increase the quality of air by filtering pollutants that are harmful to living things, such as dust, ash, pollen, smoke, particles, and the like [29-30]. Nevertheless, increasing the number of studies on this subject and repeating the existing ones elaborately is necessary so that plants can be used effectively in decreasing indoor CO<sub>2</sub> levels.

## Conclusions

The results from our study can provide significant clues on choosing plants according to the conditions of the environment and the amount of time these plants spend in the environment. For example, dieffenbachia starts to respire at a slight decline in the amount of light

in the environment, while yucca and spathiphyllum can continue to photosynthesize even under poor light conditions. Considering that these plants are used indoors and that the amount of light is limited indoors for the majority of a day, dieffenbachia should not be chosen as an indoor plant. The results from the study also indicate that CO<sub>2</sub> levels during the day were 8.3, 5.8, and 1.4 times more in yucca, spathiphyllum, and dieffenbachia, respectively, compared with the CO<sub>2</sub> levels released through respiration. Accordingly, the most suitable plant for indoor use to reduce the CO<sub>2</sub> level is yucca. However, as mentioned before, studies on this subject are still very limited, and increasing the number of studies on this topic or elaborately replicating the existing ones is required so that plants can be used effectively to reduce the CO<sub>2</sub> level indoors. In this regard, studies on different plants should continue, and plants that can photosynthesize at a faster rate indoors should be searched. Another point that should be studied is the opportunity to increase the photosynthesis rate of plants by changing indoor conditions. Here, light intensity and type of light should be studied because the effect of plants on the quality of indoor air depends on photosynthesis, and light that comes in is one of the conditions that affect photosynthesis.

### Acknowledgements

This study is funded by the Scientific and Technological Research Council of Turkey (TUBITAK) as project number 114Y033. We all thank TUBITAK for the support.

### References

- CETIN M. Determining the bioclimatic comfort in Kastamonu city. *Environ. Monit. Assess.* **187**(10), 640, **2015**.
- CETIN M. Using GIS analysis to assess urban green space in terms of accessibility: case study in Kutahya. *Int. J. Sust. Dev. World.* **22**(5), 420, **2015**.
- CETIN M. Evaluation of the sustainable tourism potential of a protected area for landscape planning: a case study of the ancient city of Pompeii in Kastamonu. *Int. J. Sust. Dev. World.* **22**(6), 490, **2015** doi: 10.1080/13504509.2015.1081651,2015.
- CETIN M., SEVIK H., ISINKARALAR K. Changes in the particulate matter and CO<sub>2</sub> concentrations based on the time and weather conditions: the case of Kastamonu, *Oxid. Commun.* **40** (1-II), 477, **2017**.
- CETIN M. A Change in the Amount of CO<sub>2</sub> at the Center of the Examination Halls: Case Study of Turkey. *Studies on Ethno-Medicine*, **10** (2), 146, **2016**. Retrieved from [http://krepublishers.com/02-Journals/S-EM/EM-10-0-000-16-Web/S-EM-10-2-16-Abst-PDF/S-EM-10-2-146-16-444-Cetin-M/S-EM-10-2-146-16-444-Cetin-M-Tx\[7\].pdf](http://krepublishers.com/02-Journals/S-EM/EM-10-0-000-16-Web/S-EM-10-2-16-Abst-PDF/S-EM-10-2-146-16-444-Cetin-M/S-EM-10-2-146-16-444-Cetin-M-Tx[7].pdf)
- CETIN M., SEVIK H. Measuring the impact of selected plants on indoor CO<sub>2</sub> concentrations. *Pol J Environ Stud.* **25** (3), 973, **2016**.
- ISINKARALAR K., CETIN M., ICEN H.B., SEVIK H. Indoor quality analysis of CO<sub>2</sub> for student living areas. *The International Conference on Science, Ecology and Technology I (Iconsete'2015 – Vienna) Abstract Book*, , 123, **2015**. August 25-28, 2015, Vienna, Austria.
- BULGURCU H., ILTEN N., COSGUN A. Indoor air quality problems and solutions in schools. *Journal of Installation Engineering*, **96**, 59, **2006**, [In Turkish].
- CETIN M., SEVIK H. Evaluating the recreation potential of Ilgaz Mountain National Park in Turkey. *Environ. Monit. Assess.* **188** (1), 52, **2016**.
- AYDOGAN A., MONTOYA L.D. Formaldehyde removal by common indoor plant species and various growing media. *Atmos. Environ.* **45** (16), 2675, **2007**.
- PAPINCHAK H., HOLCOMB E.J., OREDOVICI B.T., DECOTEAU D.R. Effectiveness of houseplants in reducing the indoor air pollutant ozone. *HortTechnol.* **19** (2), 286, **2009**.
- TANI A., HEWITT C.N. Uptake of aldehydes and ketones at typical indoor concentrations by houseplants. *Environ. Sci. Technol.* **43** (21), 8338, **2009**.
- YANG D.S., PENNISI S.V., Son K.-C., Kays S.J. Screening indoor plants for volatile organic pollutant removal efficiency. *HortScience*, **44** (5), 1377, **2009**.
- SEVIK H., AHMAIDA E.A., CETIN M. Chapter 31: Change of the Air Quality in the Urban Open and Green Spaces: Kastamonu Sample. Eds: Irina Koleva, Ulku Duman Yuksel, Lahcen Benaabidate. *Ecology, Planning and Design*. St. Kliment Ohridski University Press, ISBN: 978-954-07-4270-0, 409, **2017**.
- CETIN M., SEVIK H. Chapter 5: Assessing Potential Areas of Ecotourism through a Case Study in Ilgaz Mountain National Park. Eds: Leszek Butowski. *Tourism-From Empirical Research Towards Practical Application*. InTech, ISBN:978-953-51-2281-4, 81, **2016**.
- CETIN M., MOSSI M.M.M., AKBUDAK K.Y. Chapter 35: Visual Examination of Natural and Cultural Landscape Values in Kastamonu City Center for Sustainable Spatial Development. Eds: Irina Koleva, Ulku Duman Yuksel, Lahcen Benaabidate. *Ecology, Planning and Design*. St. Kliment Ohridski University Press, ISBN: 978-954-07-4270-0, 465, **2017**.
- CETIN M., ADIGUZEL F., KAYA O., SAHAP A. Mapping of bioclimatic comfort for potential planning using GIS in Aydin. *Environ Dev Sustain.* **1**, **2016**, doi:10.1007/s10668-016-9885-5
- SEVIK H., CETIN M., ISINKARALAR K. Effects of some indoor ornamental plants on the amount of indoor Carbondioxide. *Duzce University. The Journal of Science and Technology.* **4** (2), 493, **2015**. [In Turkish].
- SEVIK H., CETIN M., BELKAYALI N. Determination of characteristics factors of grafted natural varieties in landscaping: a case study of Black Pine (*Pinus nigra*) clone. *Oxid. Commun.* **39** (3), 2820, **2016**.
- SEVIK H., CETIN M. Effects of water stress on seed germination for select landscape plants. *Pol J Environ Stud.* **24** (2), 689, **2015**.
- SEVIK H., CETIN M., GUNEY K., BELKAYALI N. The influence of house plants on indoor CO<sub>2</sub>, *Pol. J. Environ. Stud.* **26** (4), 1-9, **2017**. DOI: 10.15244/pjoes/68875
- KACAR B., KATKAT V., ÖZTÜRK Ş. *Plant Physiology* (4. Edition). Ankara: Nobel Publication distribution **2010** [In Turkish].
- SEVIK H., CETIN M., BELKAYALI N. Effects of forests on amounts of CO<sub>2</sub>: case study of Kastamonu and Ilgaz Mountain National Parks. *Pol. J. Environ. Stud.* **24** (1), 253, **2015**.

24. CETIN M. 2016. Sustainability of urban coastal area management: a case study on Cide. *J Sustainable For.* **35** (7), 527, **2016**.
25. CETIN M. Changes in the amount of chlorophyll in some plants of landscape studies. *Kastamonu University Journal of Forestry Faculty.* **16** (1), 239, **2016**.
26. CETIN M. Consideration of permeable pavement in landscape architecture. *J Environ Prot Ecol.* **16** (1), 385, **2015**.
27. TARRAN J., TORPY F., BURCHETT M. Use of living pot-plants to cleanse indoor air – research review. *Proceedings of the Sixth International Conference on Indoor Air Quality, Ventilation and Energy Conservation in Buildings – Sustainable Built Environment*, **3**, 249, **2017**.
28. CETIN M., SEVIK H. Indoor quality analysis of CO<sub>2</sub> for Kastamonu University, Conference of the International Journal of Arts & Sciences, **9** (3), 71, **2016**.
29. CETIN M. Determination of bioclimatic comfort areas in landscape planning: A case study of Cide Coastline. *Turkish Journal of Agriculture-Food Science and Technology.* **4** (9), 800, **2016**.
30. CETIN M. Change in amount of chlorophyll in some interior ornamental plants. *Kastamonu University Journal of Engineering and Sciences* **3** (1), 11, **2017**.