

Snow Cover Studies: a Review on the Intensity of Human Pressure

Katarzyna Cichała-Kamrowska¹, Marek Błaś^{2**}, Mieczysław Sobik²,
Żaneta Polkowska¹, Jacek Namieśnik^{1*}

¹Department of Analytical Chemistry, Chemical Faculty, Gdańsk University of Technology,
G. Narutowicza 11/12, 80-233 Gdańsk, Poland

²Department of Climatology and Atmospheric Protection, Institute of Geography and Regional Development,
University of Wrocław, Kosiby 8, 51-670, Wrocław, Poland

Received: 27 May 2010

Accepted: 28 January 2011

Abstract

Snow cover should be treated as a highly dynamic medium. A variety of physical and chemical processes interact to alter its chemical composition and affect the metamorphism of its individual crystals and layers; this, in turn, has a direct influence on their mutual adhesion, and ultimately on the stability of the entire snow cover.

The literature data here reported provide basic information on the forms of solid precipitation, the types of snow and their generation, as well as the formation, stability, and chemical composition of snow cover in terms of a broad spectrum of inorganic compounds and of the levels of certain organic compounds in snow-pit sampled in mountain regions in Poland and elsewhere in the world.

Keywords: snow, snowpit, snow cover, deposition of contaminants, snow cover stability

Introduction

Snow on the ground is a veritable storehouse of atmospheric contaminants. The chemical composition of subsequent meltwater is not merely the sum of all chemical components accumulating in snow cover after each successive precipitation and deposition. As long as snow cover persists, it is subject to numerous processes: deposition of contaminants and also rime on its surface, wind erosion and accumulation from different areas, partial top-down or bottom-up thawing, and related possible seepage of meltwater into the ground. As a result, the chemistry of snow on the ground is continually changing. If the snow does not melt, the overall contaminant load builds up with each succeeding snowfall.

The reason for studying the chemistry of snow is the considerable influence of snow cover on the deposition rate of contaminants. Assays of the chemical composition of atmospheric precipitation within 24 h do not reflect the real rate of pollutant deposition. When conditions for evaporation are unfavourable, precipitation increases the amount of water in the snow cover and the contaminant load 'stored' in.

This condition is particularly enhanced in mountain regions where, given favourable weather, the snow on the ground can persist for several months. Only once the thaw sets in are pollutants released from the snow cover, from where they pass into the soil and surface/ground waters [1]. The result is that in the space of a very short time (a few days), a large amount of the accumulated chemical load of contaminants is released from the snow cover, leading to intense environmental stress.

Research programmes and projects for evaluating the state/composition of snow cover and the processes occur-

*e-mail: chemanal@pg.gda.pl

**e-mail: blasnm@meteo.uni.wroc.pl

ring in it are crucial not only from meteorological and climatic points of view, but also from hydrological and applied ones. The role played by snow in nature is tremendously important, particularly in relation to the balance between incoming solar radiation and outgoing from the Earth's surface, to the heat balance between the atmosphere and the lithosphere, and also in conditioning weather, climate, and human health [2].

Solid Atmospheric Precipitation

Atmospheric precipitation, both liquid and solid, are formed above the condensation level, that is the altitude at which water vapour becomes saturated. Clouds (from which precipitation falls) can be divided into those containing water, ice, and a mixture of water and ice. Water clouds form at temperatures above 0°C, mixed ones between 0°C and 40°C. Ice clouds form only at very low temperatures. Snow occurs from the following types of clouds: *Altostratus*, *Nimbostratus*, *Stratocumulus*, *Stratus* (granular snow only), *Cumulus*, and *Cumulonimbus*.

The microphysical processes taking place within clouds and leading to the formation of water droplets and ice crystals are complex. Most or nearly all the incipient crystals are formed around ice nuclei, which are outgoing agents, e.g. mineral, artificial, or smoke particles carried upwards to considerable heights from the Earth's surface by convection. In the upper part of the cloud, the film of water adhering to the nucleus forms droplets or incipient ice crystals. A snowflake is formed as a result of sublimation (the direct transition from ice to water vapour driven by surrounding heat) or coagulation (the coalescing of water droplets into larger drops as a result of collisions between them during gravitational freefall or turbulent air movements). Overcoming the force of the ascending air currents, the largest flakes fall to the base of the cloud, where supercooled water droplets are prevalent. The snowflake becomes larger by sublimation. Its convex parts most distant from the centre grow most quickly, and a six-beam star grows out of a hexagonal flake [3].

Once snow flakes have escaped from the cloud, they begin their journey toward the Earth's surface. Depending on the altitude of the cloud from which the snow is falling, the distance to be covered may range from 500 to 5,000 m. The forces of gravitation, floating, and air resistance together act on a falling snowflake. The rate of fall of a snowflake is known as its hydraulic velocity and depends mainly on its mass, shape and size, and air density. In frosty weather, snow in the shape of small needles and columns usually falls with speeds up to 0.5 m/s, flakes and stars 0.5-1.0 m/s, graupel (snow pellets and ice pellets) at 1.0-2.7 m/s. The velocity of a snowflake and the distance it travels depend mainly on the speed of the wind during the fall [4].

Snow is solid precipitation in the form of fine ice crystals, often joined together in various shapes (snowflakes). On the ground these crystals form a porous cover of low density. The shapes, sizes and concentrations of snow crys-

tals vary considerably, depending on the temperature at which they were created and the conditions prevailing during their development. Snow is not the only kind of solid precipitation. On their way down through a cloud, snowflakes collide with supercooled droplets of water and become granular, the large drops enveloping and freezing on them transform them into ice/freezing rain or hail [4-9].

According to the classification of the International Commission on Snow and Ice (ICSI), the following forms of precipitation particles can be distinguished (Table 1) [9].

The Formation of Snow

During a single fall of snow a great number of different types of crystal are usually identifiable; in nature, however, these crystal types may be considered infinite in number. International classifications have been developed in an attempt to categorize snow crystal forms. One of the earliest, compiled by Professor U. Nakaya from the University of Hokkaido, was the "Classification of snow crystals" [7]. This classification describes 41 different morphological types divided into several groups (Table 1):

- needle-shaped crystals
- columnar crystals
- planar crystals
- aggregates (combinations) of columnar and flat crystals
- columnar crystals with extensive facets
- crystals with surface hoar, and graupel (depending on cloud temperatures and physical processes in the cloud)
- irregular snow particles

An extension of Nakaya's classification is the one by Magono and Lee [8]; in fact, it is the most widely ramified classification of snow crystals in existence, covering 80 separate morphological types.

The structure of a snowflake results from the sum of the process of its formation: this begins when ice condenses on a nucleus. Initially, a hexagonal or planar crystal is formed, a fraction of a millimetre in size. At temperatures between -3°C and -8°C and below -30°C conditions favour the deposition of ice on the crystal's edges, from which six branches (arms) grow out. At temperatures between -3°C and -5°C and below -60°C, ice is deposited on the facets, and the crystal grows vertically, forming needles. At random places along the needle the crystallization of a new needle begins at an initial angle of 60°. Other shapes occur when, during the crystal growth, the ambient conditions change (Table 1) [4, 9].

Most snow crystals are planar and have six, more or less identical, branches (arms). Depending on the temperature, humidity and pressure of the air, other forms like columns, needles, plates and pellets may appear.

The Formation of a Snow Cover

Snow cover is a natural body with specific properties that is subject to continual change under the influence of external and internal factors (Fig. 1).

Table 1. The main forms of precipitation particles (ICSI) – basic information [9].

Form of precipitation particles	Temperature of formation (physical process)	Brief description (shape)
Columns	Growth from water vapour at -3 to -8°C and below -30°C	Prismatic crystal, solid or hollow
Needles	Growth from water vapour at high super-saturation at -3 to -5°C and below -60°C	Very thin snow particles, approximately cylindrical in shape
Plates	Growth from water vapour at 0 to -3°C and -8 to -70°C	Plate-like, mostly hexagonal structure
Stellars, Dendrites	Growth from water vapour at high super-saturation at 0 to -3°C and at -12 to -16°C	Planar or spatial crystals resembling patterned stars, usually with 6 branches.
Irregular crystals	Polycrystals growing in varying environmental conditions	Clusters of very small crystals.
Graupel	Heavy riming of particles by accretion of supercooled water droplets	Heavily rimed particles, spherical, conical, hexagonal, or irregular in shape (size: ≤ 5 mm)
Hail	Growth by accretion of supercooled water	Laminar internal structure, translucent or milky glazed surface (size: > 5 mm)
Ice pellets	Freezing of raindrops or refreezing of largely melted snow crystals or snowflakes. Graupel or snow pellets encased in thin ice layer (small hail)	Transparent surface, mostly small spheroids, white core (size: ≤ 5 mm)
Rime	Formed onto surfaces as well as on freely exposed objects	Accretion of small, supercooled fog droplets frozen in place. Thin breakable crust forms on snow surface if process continues long enough.

Table 2. Classification of the main types of snow [10, 11].

Type of snow	Conditions of formation
Fresh	Powdery snow crystals falling at temperatures $< -10^{\circ}\text{C}$ do not coalesce, and form a light, fluffy layer not bound to the substrate.
Loose, graupel-like	Snow formed from falls of graupel and fine granular snow
Settled, wind-blown	Dry snow, at temperatures from -10°C to -3°C , displays only a slight tendency to coalesce. The cover of powdery snow begins to settle. An important feature of such snow is the ease with which it is blown by the wind.
Compressed, dry	Packed powder is patchy in occurrence and consists of heavy but not wet snow, well compressed due to the action of the wind and its own weight.
Wet snow	At temperatures at and above -3°C falling snow crystals are wet, which facilitates their coalescence. A layer of such snow easily attaches to the substrate. This snow has quite a high specific gravity (up to 200 kg/m^3) and considerable plasticity. It has a fairly high liquid water content. The greater the plasticity of the snow, the more resistant the snow cover is to forces tending to detach it from the substrate. With their high coherence and weight, damp or wet snows are wind-resistant.
Crust	Snow with an icy surface, breakable under the icy surface, the snow is mostly dry
Ice crust	Snow with a supportable icy surface
Granular snow	A cover of granular snow consists of large, hard crystals formed as a result of recrystallization.
Firm	Consists of large, irregular grains, not coalesced, always wet. The interstices between the grains are air-filled. This is the final phase in the metamorphosis of snow

Snow derived from precipitation and lying for some time on the ground as a layer no thinner than 0.5 cm constitutes snow cover. Once on the ground, snow crystals retain their original form for only a short time, after which they are transformed into different shapes, depending on temperature, humidity, wind, time on the ground, crystal form, and the thickness of the snow cover. As the snow cover thickens, the contaminants contained in the particles of precipitation gradually accumulate within it, but when it

melts, the contaminants are released along with the meltwater, which seeps into the soil and surface/ground waters.

Depending on the duration of the interval between two falls of snow and on the weather during such intervals, the layers of snow are transformed both on the surface and within. Different types of snow begin to form. The snow cover changes under the influence of melting, wind, evaporation, recrystallization, and pressure. In the Polish mountains nine types of snow occur during winter (Table 2) [10].

Three states of matter can be distinguished in snow forming a cover:

- solid – in which ice takes the form of crystals or grains;
- liquid – when at temperatures around the melting point water forms a coating around the grains of snow;
- gas – formed by water vapour or air present in the interstices between the snow crystals.

The variations in the proportions of each of these phases and the metamorphism taking place within the snow directly affect the properties of a snow cover.

The Processes Taking Place within a Snow Cover

Both on the surface and within, a cover of snow is subject to continual change as a result of the physical processes taking place in it. Both external and internal factors affect these processes and the time when they occur. External factors include temperature, wind, radiation, and air humidity, while internal factors include the pressure of successive layers of snow, and the temperature within the snow cover.




Fig. 1. Processes causing changes in the physical/chemical texture of snow cover.




Fig. 2. The changes in height of snow cover (HS) and air temperature (T) in winter 2005-06 at Szrenica in the Giant Mountains (Poland).

An important aspect, affecting the way individual snow crystals join together is their morphology (shape) and the processes of sublimation and recrystallization taking place within the snow cover. The interaction of these factors leads to changes in the properties of the snow as a result of:

- a) the constant-temperature metamorphism of snow – i.e. **destructive metamorphism** – that occurs in all crystals throughout the snow cover at temperatures $<0^{\circ}\text{C}$, most rapidly just below 0°C , and which declines at -15°C and ceases altogether at -40°C ; and
- b) the metamorphism of snow under the influence of a temperature gradient – i.e. **constructive metamorphism** – that takes place when there is a temperature difference between the bottom layer of snow (usually 0°C) and the snow surface; highly unstable, cup-shaped crystals then appear, a process that takes place rapidly when there is a big difference in temperature [4].

The metamorphosis of snow oscillates between melting and refreezing when temperatures change from above-zero to sub-zero, and vice versa. The effect of 24 h changes in temperature and the melting and refreezing give rise to the fact that the melting snow grains become larger and more spherical; the end result is a stable snow cover layer. The degree of metamorphosis of the various layers has a direct influence on mutual adhesion, and by extension, on the stability of the snow cover as a whole.

The potential causes of deviations of height and duration of a snow cover from the hypsometric profile include:

- exposure to the sun (insolation)
- total winter precipitation
- air temperature
- the extent to which snow is wind-blown
- surface vegetation

Both temperatures, of air and within snow cover, are main factors affecting the metamorphosis of snow. Snow has low thermal conductivity, so temperature within snow cover changes slowly, far more than the corresponding temperature changes in the atmosphere. Temperatures are usually highest in the deepest layers, so water vapour diffuses from the lower layers to the surface. The consequence of this is a loss of crystals in the lower part and a loose texture of the lower layers, which eventually disturbs the equilibrium of the snow cover. When above-zero temperatures act on the snow cover for a longer period of time, the water penetrating down into the bulk of the snow gives rise to wet snow. When air temperature is low, snow mass contraction takes place, leading to the appearance of cracks and crevices. The surface layers are also subject to metamorphosis, changing the type of snow from which they were




Fig. 3. Snowpit hardness from: 1. Greenland Ice Core (2005) [20], 2. The Summit of Nevado Illimani, Bolivia (1999) [21], 3. Pika Glacier – Alaska Range (2001) [22], 4. The Swiss Alps [23].

Table 3. Primary physical characteristics of deposited snow (ICSI-2009) [9].

Snow cover measurements		Method of measurement			
Grain shape	1.	Precipitation particles		(Fig. 5)	
	2.	Machine-made snow			
	3.	Decomposing and fragmented precipitation particles			
	4.	Rounded grains			
	5.	Faceted crystals			
	6.	Depth hoar			
	7.	Surface hoar			
	8.	Melt forms			
	9.	Ice formations			
Grain size	very fine	<0.2 mm		(Fig. 6)	
	fine	0.2-0.5 mm			
	medium	0.5-1.0 mm			
	coarse	1.0-2.0 mm			
	very coarse	2.0-5.0 mm			
	extreme	>5 mm			
Snow density [kg/m ³]	Determined by weighing snow of a known volume				(Fig. 7)
Snow hardness (Figs. 3, 4)	HH index	Hand-hardness test		Ram resistance [N]	
				Range	Mean
	1.	very soft	Fist	0-50	20
	2.	soft	Four fingers	50-175	100
	3.	medium	One finger	175-390	250
	4.	hard	Pencil	390-715	500
	5.	very hard	Knife blade	715-1,200	1,000
6.	ice formations	Ice	>1200	>1,200	
Liquid water content	Wetness index		Range		
	1.	dry	0% water		
	2.	moist	< 3% water		
	3.	wet	3-8% water		
	4.	very wet	8-15% water		
	5.	slush	> 15% water		
Temperature [°C]	Measuring air temperature, temperature at the snow surface, ground temperature, snow profile temperature: 5 cm above the ground, and 5 cm below the surface				(Fig. 8)
Impurities	It cover those cases where the type and amount of an impurity influence physical characteristics of the snow				
Thickness of layers [cm]	An essential parameter when characterizing the current state of snowpack. Usually measured vertically				

formed. When water vapour rises from the warmer grains of snow near the ground to the upper layers, where it condenses in the colder air, a crust is formed. Firn is formed when heat entering the snow pack from above initiates the melting and refreezing of snow crystals. In contrast, an ice crust is formed when after a period of prolonged insolation, during which the surface layer of the snow cover melts,

there is a sudden drop in temperature at the surface (Table 2) [1, 11]. The wind also affects the nature of the surface layer of snow. Wind breaks up and comminutes the snow crystals as they are blown about by the wind. On windward slopes the snow is blown away and compacted, it is deposited on ridges forming cornices, and on leeward slopes, it forms wind slabs.

Table 4. Additional measurements of snow cover [9].

Snow cover measurements	Units	Method of measurement
Height	cm	Height is the coordinate measured vertically (line of plumb) from the base. Ground surface is usually taken as the base, but on firm fields and glaciers it refers to the level of either the firm surface or glacier ice.
Thickness	cm	Used when measurements are taken perpendicularly, i.e., at a right angle to the slope on inclined snow covers. It is measured from the base and the same comments apply as for height.
Height of snowpack	cm	Total height of the snowpack, i.e., the vertical distance from base to snow surface.
Height of new snow	cm	The thickness of freshly fallen snow that accumulated on a snow board during a standard observation period of 24 hours.
Snow water equivalent	mm w.e.* kg/m ²	Snow water equivalent is the depth of water that would result if the mass of snow melted completely. It is the product of the snow height in metres and the vertically-integrated density in kilograms per cubic metre. It can be describe by: snow water equivalent of snow cover (SWE, HSW), water equivalent from the base up to the height H (HW), water equivalent of a single layer of thickness L (LW), and water equivalent of snowfall (HNW).
Water equivalent of snowfall	mm w.e.* kg/m ²	Typically measured for a standard observation period of 24 hours.
Snow strength	Pa	Snow strength can be regarded as the maximum or failure stress on a stress-strain curve. It is the maximum stress snow can withstand without failing or fracturing.
Penetrability of snow surface	cm	The depth that an object penetrates into the snow from the surface.
Surface features	cm	Described more generally in terms of roughness elements that are not related to snow microstructure: smooth, wavy, concave furrows, convex furrows, random furrows.
Snow-covered area	1%	The areal extent of snow-covered ground, usually expressed as a fraction (%) of the total area investigated.
Slope angle	°	The acute angle measured from the horizontal to the plane of a slope. Slope angle is measured with a clinometer.
Aspect of slope	°	Aspect is the compass direction toward which a slope faces. The direction is taken downslope and normal to the contours of elevation, i.e., along the fall line.
Time	s, min, h, d, week, month, year	Usually given in seconds

Fig. 2 shows a continuous record of snow cover height and air temperature.

Progressive deforestation in the last 20 years has also had a significant influence on snow cover formation. In the Jizera Mountains (west part of Western Sudetes), where the rate of deforestation has been even faster than in the Giant Mts. (a central part of the Western Sudetes), the snow cover remains on the ground for a good three weeks less in clear-felled areas than in the adjoining forest. During the first half of winter the height of the snow cover and the supply of water it contains are similar both within the forest and beyond it. But when the thaw sets in, the snow cover in the open ground melts far more rapidly, and the pH and contaminant load of the resulting meltwaters are correspondingly high [12]. The same scenario was repeated in the Polish part of the Giant Mts and in the Jizera Mts in 1998, when the thawing period in the catchment area of the Kamienna River was 25 days shorter than normal and the largest masses of meltwater flowed downstream already in April rather than in May [13].

Measurement of Snow Cover

The standard observations of snow cover do not take into account its internal structure and properties to any great extent. Obtaining full information on the current structure of a snow layer and the processes occurring within it enable its stability to be analyzed, which is a crucial factor in assessing and forecasting avalanche hazards. For this purpose meteorologists carry out features and additional measurements of the snow cover (Tables 3 and 4) [9]. Figs. 3-8 illustrate measurements of the hardness, density and temperature of snow, and crystal sizes and grain forms.

If a pit is dug in the snow cover and one of its sides cut vertically, its stratified structure is revealed. The sediment accumulated in the surface layers of snow is known as *snowpit*. The layers making up a cover of snow differ in their physical and mechanical properties, depending on the weather conditions and time when they were formed, and also on how long they have been in existence. With a com-




Fig. 4. Ram penetrometer profile of southern and northern aspects in March 2000 (Jackson Hole Mountain Resort and Grand Teton National Park, Wyoming) [24].




Fig. 5. Grain profile measured at sites A and B at Niseko, Japan, in January 1998 [24], and February and March 2003 [26].

Table 5. Classification of ram hardness profiles [14, 17, 25, 27].

Type	Profile	Base	Stability
1		weak	unstable
2			depends on other conditions
3			
4			
5			unstable
6		strong	stable
7			unstable
8			depends on other conditions
9			unstable
10			stable

Table 6. Relative importance of parameters for interpretation of profiles and the RB (Rutschblock test) [14-19, 30, 31].

Profile interpretation	Importance	RB interpretation	Importance
Ram hardness profile	1	Slab thickness	0.9
Weak layers	1	Type of fracture plane	0.9
Rutschblock	0.9	RB score	0.8
Grain type and size	0.7	Type of failure	0.8
Snow temperature	0.5	Slope angle	0.7
Hand hardness	0.4	Snow depth	0.6
Liquid water content	0.1	Slab hardness	0.4

plete set of measurements for a given profile at hand, and having entered these data into the relevant computer program, we obtain a graphic picture of the snow cover and the vertical distribution of its parameters; this is known as a description of the stratigraphic profile.

Weak layers are mostly soft, usually 'fist' or '4-fingers' in the hand-hardness scale. It is important to search for differences in hardness, as they are often associated with weaker layers. Increasing hardness with decreasing depth




Fig. 6. Snowpit grain size from: 1. Pika Glacier – Alaska Range (2001) [22], 2. The Summit of Nevado Illimani, Bolivia (1999) [21], 3. the Swiss Alps [23], 4. Site A – Niseko, Japan – January 1998 [24].

may indicate that the snowpack is unstable. Weak layers responsible for snowpack instability are often found between two hard layers. Frequent small changes in hardness in the snowpack profile are not as critical as a large difference between two layers [5, 14, 17, 27].

The vertical distribution of a snowpack's penetration resistance can be illustrated by a ram profile. The main disadvantage, however, is that because of low resolution, hard or soft thin layers cannot be detected. The ram hardness profile can be presented in different ways (see Table 5 – the wider the darkened area, the harder the snowpack layer). The fact that the base may be weak does not necessarily mean that the snowpack is unstable [5, 14].

Surface hoar, faceted grains, or depth hoar are usually present in the weak layers of skier-triggered avalanches. These grains are coarse and have plane surfaces, so the bonding is weaker and hence the grain types are weaker, too.

If the grains are buried they become rounded in shape and thus are less critical. The snowpack is stabilized by the ice crust created by the melting and refreezing of surface snow layer. On the other hand, they may provide surfaces over which new, upper snow layers can slip. Wetting the crust layer may reduce the friction and instability of a snowpack [5, 14].

The strength of a snow layer depends on grain size. There is negative correlation between grain size and number of bonds per unit volume. The dependence of grain size on snow depth is presented in Fig. 6 [5, 14].

The density of a snow layer does not directly indicate its instability, but critical layers are often less dense than the surrounding ones. Generally, a dense snow layer overlying a low-density layer may be dangerous. Obtaining more information about stability, strength, hardness, and grain size measurements is necessary, not just density measurements [5, 14].




Fig. 7. Snowpit density from 1. The High Arctic (1981) [28], 2., 5. Greenland Ice Core (2004) [29], 3. The Summit of Nevado Illimani, Bolivia (1999) [21], 4. Greenland Ice Core (2005) [20].

For establishing the stability of a snow layer, temperature seems to be a parameter of little value. There are no clear rules about how snow temperature should be evaluated [5, 14-16].

Table 6 provides information on the importance of the various parameters determined in snow samples taken from a snowpit profile (1: very important, 0: not important).

A snow cover is a highly dynamic medium. Both physical and chemical processes can cause its chemical composition to change, so comprehensive tests should be carried out to ascertain the contaminant content in snow. The chemical composition of snow layers reflects the composition of native atmospheric aerosols, dry precipitation, and the nature of the adsorbent surface. The accumulation of components in snow cover present in the atmosphere depends on the weather and on the properties of the surface covered by the snow. The chemical composition of snow can provide a great deal of useful information on the levels of contaminants in the atmosphere, the changes in these

levels over time, and the mechanism of their transport and deposition. Changes in the chemical composition of the atmosphere are reflected in changes in the chemical composition of snow and should be associated with various sources and transport pathways (Fig. 1).

The Chemistry of Snow Cover

Reliable information on the chemistry of snow cover is provided by the assay of representative samples collected using the appropriate equipment and correctly prepared for their analysis for target components and parameters characterizing the chemical composition and the processes taking place within the snow cover.

A number of problems and challenges crop up with this type of analysis, which the analysts involved in this type of assay have to solve. The most important of these include:




Fig. 8. Temperature profile of southern and northern aspects in March 2000 (Jackson Hole Mountain Resort and Grand Teton National Park, Wyoming) [27], and the Swiss Alps (1937, 1950) [23].

Table 7. Results of snowpit sample analyses: a review of literature data

Sampling area	Sampling period	Features of the sampling site	Analytes	Concentration ranges	References
Himalayas – Mt. Everest	1998-2002	The East Rongbuk glacier on the northern slopes of Mt. Everest (27°59'N, 86°55'E). Three snowpit samples were taken at altitudes of 6,400 m and 6,500 m.	NO ₃ ⁻	0.18-1.37 [μeq/l]	[33]
			Cl ⁻	0.55-1.29	
			SO ₄ ²⁻	0.69-0.80	
			NH ₄ ⁺	nd – 1.41	
			Ca ²⁺	2.79-8.26	
			Na ⁺	0.54-0.75	
			K ⁺	0.24-0.46	
			Mg ²⁺	0.21-0.49	
China	1996	Glacier at the head of the River Ürümqi, Tien Shan, altitude 4,230 m	NO ₃ ⁻	0-0.85 [10 ⁻⁶ g/g]	[31]
			Cl ⁻	0.06-1.11	
			SO ₄ ²⁻	0.1-2.3	
Greenland	1988-89	A sampler was placed at the summit of Dye 3 in the southern part of the ice cover at latitude 65°11N, longitude 43°50W, and altitude 2,479 m	Al ³⁺	1.82-6.82 [μg/l]	[34]
			Fe ²⁺	1.10-3.02	
	1991-92	Summit of the Greenland Ice Sheet – samples taken at latitude 72°20N, longitude 38°45W and altitude 3,240 m	Na ⁺	0.1-3.8	[35]
			Mg ²⁺	0.1-8.1	
			K ⁺	0.05-6.9	
	1996	Summit of the Greenland Ice Sheet – the snowpit was located in the SW part of the Greenland Ice Sheet. The camp was sited at latitude 72°58N, longitude 38°48W and altitude 3,207 m	PHE	1295-4,536 [pg/kg H ₂ O]	[36]
			FLU	685-5718	
			PYR	449-4,069	
			BaA	nd – 1081	
			CHR	nd – 1697	
BeP			nd – 1310		
BaP			nd – 2540		
InP			nd – 781		
BgP			nd – 919		
RET	57-399				
Alps	1991	Summit of the Sonnblick (Austria)	NO ₃ ⁻	0-0.13 [ppm]	[37]
			Cl ⁻	0-0.09	
			SO ₄ ²⁻	0-0.31	
			NH ₄ ⁺	0-0.08	
			Na ⁺	0-0.94	
			K ⁺	0-0.15	
	1992	Summit of the Sonnblick (Austria)	NO ₃ ⁻	0.461-0.618	
			Cl ⁻	0.047-0.111	
			SO ₄ ²⁻	0.298-0.372	
			NH ₄ ⁺	0.048-0.08	
			Na ⁺	0.027-0.074	
			Mg ²⁺	0.021-0.03	
			Ca ²⁺	0.056-0.149	

Table 7. Continued.

Sampling area	Sampling period	Features of the sampling site	Analytes	Concentration ranges	References
Alps	1994	Jungfrauoch (Switzerland) – latitude 46.55°N, longitude 7.98°E, altitude 3,400 m	NO ₃ ⁻	2.8-7.1 [μeq/l]	[38]
			Cl ⁻	0.7-5.6	
			SO ₄ ²⁻	2.5-8.1	
	1992-95	The Stubai glacier in the Tyrolean Alps, altitude 2,950 m, latitude 46°59'N, longitude 11°6'E	average value [μeq/l]		[30]
			NO ₃ ⁻	5.3	
			Cl ⁻	1.1	
			SO ₄ ²⁻	3.4	
			NH ₄ ⁺	2.3	
			Na ⁺	1.0	
			Mg ²⁺	0.5	
Ca ²⁺	2.1				
Poland	1993	Jakuszyce	NO ₃	0.339-18.301	[46]
			SO ₄	2.550-8.212	
			Mg	0.105-0.224	
			Na	0.405-0.577	
			Pb	0.008-0.028	
			Zn	0.035-0.063	
		Łabski Szczyt	NO ₃	0.281-9.390	
			SO ₄	0.838-7.510	
			Mg	0.049-0.198	
			Na	0.257-0.625	
			Pb	0.001-0.047	
			Zn	0.048-0.385	
		Szrenica	NO ₃	0.266-2.730	
			SO ₄	1.390-2.301	
			Mg	0.058-0.165	
			Na	0.324-0.495	
			Pb	0.004-0.019	
			Zn	0.037-0.061	
		Odrodzenie	NO ₃	0.595-1.880	
			SO ₄	0.970-3.270	
			Mg	0.035-0.221	
			Na	0.201-0.527	
			Pb	0.001-0.024	
			Zn	0.009-0.058	
Śnieżka	NO ₃	0.319-1.220			
	SO ₄	1.320-2.190			
	Mg	0.039-0.151			
	Na	0.129-0.578			
	Pb	0.008-0.130			
	Zn	0.019-0.066			

- limited access to the appropriate equipment for collecting samples of the different types of solid precipitation and snow cover layers
- the requirement to apply appropriate procedures during the transport and sample preparation stages, prior to the final determination stage
- the lack of reference materials essential for validating the analytical procedures applied, ensuring the appropriate level of control and assuring the quality of the analytical results

It is clear from an analysis of literature data that more and more results are becoming available from analyses of snowpit samples carried out to determine the levels of different components. Such research has been carried out using snowpit samples collected in very different parts of

the world (Greenland, the Alps, the Andes, the Himalayas, Antarctica, the Sudeten Mountains) [21, 32-40]. Levels of the following components were assayed in these samples:

- anions: Cl^- , NO_3^- , SO_4^{2-} ;
- cations: H^+ , Na^+ , NH_4^+ , K^+ , Mg^{2+} , Ca^{2+} ;
- PAHs: phenanthrene, fluoroanthene, pyrene, benz[a]anthracene, chrysene, benzo[e]pyrene, benzo[a]pyrene, indeno[1,2,3-cd]pyrene, benzo[ghi]perylene, retene (1-methyl-7-isopropylphenanthrene);
- metals.

Table 7 presents information on the results of studies performed so far. It gives the features of the sites from which the snow/snowpit samples were taken, the maximum and minimum concentrations determined, and in some cases the mean concentrations of the analytes.




Fig. 9. Mean concentrations of nitrate, sulphate, and ammonium ions in samples of snow collected at a) Śnieżnik, b) Orle, c) Szrenica in January 2008 (author's own results), and d) the Alps (April 1994) [32].

Attention has been drawn to the part played by snow cover in the pollution of mountain ecosystems in Poland (the Sudeten and Carpathian Mountains) since the mid-1990s [41, 42]. This problem takes on particular urgency in the Giant and Jizera Mountains (western Sudetes), where the buffering ability of the bedrock is limited.

Numerous papers have been written on the spatial conditions governing the decomposition of the snow cover in the Western Sudetes. They all emphasise that the snow cover is thicker and lies on the ground longer, the greater the absolute altitude [43-47]. They also draw attention to the influence of the local relief and vegetation on the thickness and duration of the snow cover.

Fig. 9 shows the mean concentrations of ammonium, nitrate and sulphate ions determined in samples of snow collected at Śnieżnik, Orle, and Szrenica in 2008 (author's own results) and in the Alps [32].

Comparison of the diagrams in Fig. 9 indicates that levels of nitrate, sulphate, and ammonium ions are the highest in the surface layers of snow cover.

In contrast, Fig. 10 presents the mean concentrations of sodium, potassium and magnesium ions determined in snow samples collected at Śnieżnik, Orle, and Szrenica in 2008 (author's own results) and from the East Rongbuk glacier, on the northern slopes of Mt. Everest [35].




Fig. 10. Mean concentrations of sodium, potassium, and magnesium ions determined in snow samples collected at a) Śnieżnik, b) Orle, c) Szrenica in January 2008 (author's own results), and d) from the East Rongbuk glacier, on the northern slopes of Mt. Everest (May 2000) [35].

Levels of sodium, potassium, and magnesium ions were the highest in the surface layers of the snow cover. The levels of these ions were much higher in the snow samples collected in Poland.

Fig. 11 shows the mean concentrations of selected metals (cadmium, lead, zinc, copper) determined in snow samples collected at Śnieżnik, Orle, and Szrenica in 2008 (author's own results) and in central Greenland (1992) [49].

There is considerable similarity between the highest concentrations of zinc and copper determined in surface layers of snow in both Poland and Greenland. The concentrations of Zn and Cu in the snow samples from Poland were much higher.




Fig. 11. Mean concentrations of cadmium, lead, zinc, and copper ions in snow samples collected at a) Śnieżnik, b) Orle, c) Szrenica in January 2008 (author's own results – concentrations of Zn^{2+} and Cu^{2+} x 10), d) the Greenland ice sheet (July 1992) [49].

Fig. 12 shows histograms of the mean levels of nitrate, sulphate and ammonium ions in snow samples taken from snowpits at different stations in the Polish Sudeten Mountains (author's own results), the Alps [32, 39, 40], and the Himalayas [35].

The highest levels of nitrate, sulphate and ammonium ions were recorded in the snow samples from Orle (Poland). The other Polish samples did not differ much in their ionic composition. The concentrations of these ions in the Alps and the Himalayas were 6-7 times smaller.

Fig. 13 illustrates the mean levels of the polyaromatic hydrocarbons fluoroanthene, pyrene, and retene in snow samples taken from snowpits in Greenland [50].




Fig. 12. Mean concentrations of nitrate, sulphate, and ammonium ions in snow samples collected from snowpits at different stations in the Polish Sudeten Mountains, the Alps, and the Himalayas.

These studies have shown that PAH concentrations are seasonally dependent: they are highest in winter and early spring. Identical results were obtained from snowpit samples collected in earlier years [51].

Anthropogenic PAH discharges are greater in winter in the industrialized countries, which are most often the source areas (Eastern Europe, Russia, and Northern America). Photochemical degradation of PAHs is minimal in winter because the low luminosity leads to a reduced production of OH radicals, the main photochemical aggressors of PAHs [52].

Conclusions

Analysis of snow cover supplies a complete set of information on its current structure and the processes taking place within it; this, in turn, enables its stability to be assessed, a key factor in assessing and forecasting avalanche hazards [53].

A crucial aspect of the assessment of snow cover stability is the correct interpretation of the snow profile, which requires a great deal of experience on the part of the person performing the profile. No precisely defined methods are in use, so the following parameters are normally assessed: the depth of the snow cover and the thickness of its constituent layers, the type and size of snow crystals, the hardness of the snow cover, and the water content, temperature, and

density of the snow. The vertical hardness profile and the stability test of the snow cover (for the presence of slip layers) are additional crucial factors [14]. This means of assessing the stability of a snow cover is used all over the world, as testified by the numerous publications and other information on research being carried out in different parts of the world (Greenland, Austria, Italy, Switzerland, South America, the Himalayas, Antarctica, Poland) [21, 32-40].

Studies of snow cover, including specially-dug snowpits, can supply valuable information not only about the chemistry of snow, but above all about the state of the atmosphere and the processes taking place in it. Measurements of the chemical composition of 24 h portions of winter precipitation do not reflect the actual rate of contaminant deposition. This applies to mountain regions in particular, where contaminants are not released from the snow cover to seep into the soil, and surface and ground waters until the spring thaw sets in. The 'frontal wave' that then appears removes a large proportion of the accumulated contaminant load in a very short time.

By defining the composition of contaminants present in snowpit samples we can learn much about the parts played by the most important factors governing the chemical composition of snow cover in a given area and also about the balance in the total load of contaminants present in snow cover that will eventually find its way into the environment.

The contaminant composition of the various snow layers depends on the duration of the snow cover. This, in turn, depends not only on the amounts of contaminants discharged into the atmosphere, but also on meteorological conditions such as the direction of flow of air masses, temperature, wind, humidity, pressure, and the frequency of precipitation. Such investigations enable us to monitor environmental degradation, and their results can provide a starting point for taking measures to counteract it.

Acknowledgements

This work was financially supported by Grant No. N N305 373438 and N N523 616439 from the Committee on Scientific Research.




Fig. 13. Dating the strata of superficial snow (correlation of PAHs, fluoranthene, pyrene, and retene) [50].

References

1. BŁAŚ M., CICHALA-KAMROWSKA K., SOBIK M., POLKOWSKA Ż., NAMIEŚNIK J. Conditions controlling atmospheric pollutant deposition via snowpack, *Environ. Rev.*, **18**, 87, **2010**.
2. Snow and snow cover: symposium materials, Zakopane, 15-17.03.1973, IMiGW, Warszawa, **1977** [In Polish].
3. BARTLETT J. T. Physics Cloud formation, *Phys. Educ.*, **6**, 266, **1971**.
4. LIBBRECHT K. G. The physics of snow crystals, *Reg. Prog. Phys.*, **68**, 855, **2005**.
5. COLBECK S., AKITAYA E., ARMOSTRONG R., GUBLER H., LAFEUILLE J., LIED K., MCCLUNG D., MORRIS E. The International Classification for Seasonal Snow on the Ground, Int. Comm. Snow and Ice (IAHS), World Data Center for Glaciology, Univ. of Colorado, Boulder, CO, USA, **1990**.
6. MASON B. The Physics of Cloud, Oxford University Press, **1971**.
7. NAKAYA U. Snow Crystals: Natural and Artificial, Harvard University Press, **1954**.
8. MAGONO C., LEE C. W. Meteorological Classification of Natural Snow Crystals, *Journal of the Faculty of Science, Hokkaido University*, **1966**.
9. FIERZ C., ARMSTRONG R. L., DURAND Y., ETCHEVERS P., GREENE E., MCCLUNG D. M., NISHIMURA K., SATYAWALI P. K., SOKRATOV S. A. The International Classification for Seasonal Snow on the Ground, IHP-VII Technical Documents in Hydrology N°83, IACS Contribution N°1, UNESCO-IHP, Paris, **2009**.
10. KOSIBA A. Snow, glaciers, ice sheets. WSiP, Warszawa, **1978** [In Polish].
11. LIBBRECHT'S K. Field Guide to Snowflakes, **2006**.
12. HOTTELET CH., BLAŽKOVÁ Š., BIČÍK M. Application of the ETH snow model to three basins of different character in Central Europe, *Nordic Hydrology*, **25**, 113, **1994**.
13. DUBICKI A., WOŹNIAK Z. Hydrology changes, water management in mountain regions. [In:] Reforesting in the Sudeten Mountains, scientific-technical seminar, Szklarska Poreba 1-2 October 1998, Engineers and Technicians Association of Forestry and Wood, Directorate General of State Forests, **12**, **1998** [In Polish].
14. SCHWEIZER J., WIESINGER T. Snow profile interpretation for stability evaluation, *Cold Reg. Sci. and Tech.*, **33**, 179, **2001**.
15. MCCLUNG D. M., SCHWEIZER J. Skier triggering, snow temperatures and the stability index for dry-slab avalanche initiation, *J. Glaciol.*, **45**, 190, **1999**.
16. ARMSTRONG R. Temperature and snow stability, *The Avalanche Review*, **1**, (4), 5, **1983**.
17. FÖHN P. M. B. Characteristics of weak snow layers or interfaces. Proceedings International Snow Science Workshop, Breckenridge, CO, USA, 4-8 October 1992, pp. 160-170, **1993**.
18. SCHWEIZER J., CAMPONOVO C. The skier's zone of influence in triggering slab avalanches, *Annals of Glaciology*, **32**, 314, **2001**.
19. JAMIESON B., JOHNSTON C. Snowpack characteristics for skier triggering, *Avalanche News*, **55**, 31, **1998**.
20. MARSHALL H. P., SCHNEEBELI M., KOH G. Snow stratigraphy measurements with high-frequency FMCW radar: Comparison with snow micro-penetrator, *Cold Reg. Sci. and Tech.*, **47**, 108, **2007**.
21. HARDY D., WILLIAMS M. W., ESCOBAR C. Near-surface faceted crystal, avalanches and climate in high-elevation, tropical mountains of Bolivia, *Cold Reg. Sci. and Tech.*, **33**, 291, **2001**.
22. HARPER J. T., BRADFORD J. H. Snow stratigraphy over a uniform depositional surface: spatial variability and measurement tools, *Cold Reg. Sci. and Tech.*, **37**, 289, **2003**.
23. PIELMEIER CH., SCHNEEBELI M. Developments in the stratigraphy of snow, *Surv. Geophys.*, **24**, 389, **2003**.
24. MCELWAIN J., HACHIKUBO A., NEMOTO M., KAIHARA T., YAMADA T., NISHIMURA K. Observations and simulations of the formation of the faceted snow crystals in the weak-layer of the 1998 Niseko Haru no Taki avalanche, *Cold Reg. Sci. and Tech.*, **31**, 235, **2000**.
25. DE QUERVAIN M., MEISTER R. 50 years of snow profiles on the surrounding avalanche activity, Davos Symposium, IAHS, **162**, 161, **1987**.
26. NISHIMURA K., BABA E., HIRASHIMA H., LEHNING M. Application of the snow cover model SNOWPACK to snow avalanche warning in Niseko, Japan, *Cold Reg. Sci. and Tech.*, **43**, 62, **2005**.
27. KOZAK M. C., ELDER K., BIRKELAND K., CHAPMAN P. Variability of snow layer hardness by aspect and prediction using meteorological factors, *Cold Reg. Sci. and Tech.*, **37**, 357, **2003**.
28. WOO M. A guide for ground-based measurement of the Arctic snow cover, **1997**.
29. SCOTT J. B. T., MAIR D., NIENOW P., PARRY V., MORRIS E. A ground-based radar backscatter investigation in the percolation zone of the Greenland ice sheet, *Remote Sens. Environ.*, **104**, 361, **2006**.
30. PERLA R. I., MARTINELLI M. Jr., *Avalanche handbook*, USDA Forest Service, Fort Collins, Colorado, Agriculture Handbook, **489**, 238, **1976**.
31. BIRKELAND K., JOHNSON R., HERZBERG D. The stuffblock snow stability test, U.S. Forest Service Missoula Technology and Development Center, 9623-2836-MTDC, **1996**.
32. KUHN M., HASLHOFER J., NICKUS U., SCHELLANDER H. Seasonal development of ion concentration in a high alpine snow pack, *Atmos. Environ.*, **32**, 4041, **1998**.
33. HOU S., QUIN D. The effect of postdepositional process on the chemical profiles of snow pits in the percolation zone, *Cold Reg. Sci. and Tech.*, **34**, 111, **2002**.
34. STOTTLEMYER R., TROENDLE C. A. Effect of canopy removal on snowpack quantity and quality, Fraser experimental forest, Colorado, *J. Hydrol.*, **245**, 165, **2001**.
35. KANG S., MAYEWSKI P. A., QIN D., SNEED S. A., REN J., ZHANG D. Seasonal differences in snow chemistry from the vicinity of Mt. Everest, central Himalayas, *Atmos. Environ.*, **38**, 2819, **2004**.
36. COLIN J. L., LIM B., HERMS E., GENET F., DRAB E., JAFFREZO J.L., DAVIDSON I. Air-to-snow mineral transfer-crustal elements in aerosols, fresh snow and snowpits on the Greenland ice sheet, *Atmos. Environ.*, **20**, 3395, **1997**.
37. DRAB E., GAUDICHED A., JAFFREZO J. L., COLI J. L. Mineral particles content in recent snow at Summit (Greenland), *Atmos. Environ.*, **36**, 5365, **2002**.
38. SLATER J. F., CURRIE L. A., DIBB J. E., BENNER B. A. Jr., Distinguishing the relative contribution of fossil fuel and biomass combustion aerosols deposited at Summit, Greenland through isotopic and molecular characterization of insoluble carbon, *Atmos. Environ.*, **36**, 4463, **2002**.

39. SCHÖNER W., PUXBAUM H., STAUDINGER M., MAUPETIT F., WAGENBACH D. Spatial variability in the Chemical Composition of the Snowcover at High Alpine Sites, *Theor. Appl. Climatol.*, **56**, 25, **1997**.
40. PICHLMAYER F., SCHÖNER W., SEIBERT P., STICHLER W., WAGENBACH D. Stable isotope analysis for characterization of pollutants at high elevation Alpine sites, *Atmos. Environ.*, **23**, 4075, **1998**.
41. KULICKI A., PIASECKI J., SOBIK M. Acidification of snow cover in the Klesnica valley, Snieznik Massif, The work of the Institute of Geography, Series C. Meteorology and Climatology II, *Acta Univ. Wratislaviensis*, **1705**, 107, **1995** [In Polish].
42. KOT M. Acid precipitation in the Tatra National Park, Chemistry and the impact of acid rain on the environment, edited by B. Walna, Adam Mickiewicz University, Poznań, **1996** [In Polish].
43. REUNIER H. Amount and persistence of snow cover in the Giant Mountains, *Meteorology Magazine*, **52/3**, **1935** [In German].
44. KOSIBA A. Frequency of snow cover on the lands of Silesia, *Wroclaw Scientific Society*, **21**, **1949** [In Polish].
45. KWIATKOWSKI J. Snow cover of Sudeten Mountains, density and water reserve, *Geographical Magazine*, **49**, 419, **1978** [In Polish].
46. KWIATKOWSKI J. Snow cover, frost and avalanches. Conference materials: Giant Mountains, edited by A. Jahn, Ossolineum, **1985** [In Polish].
47. HLADNÝ J., SÝKORA B. Climatology, hydrology, snow cover. Conference materials: Karkonosze National Park, Administration of Karkonosze National Park in the State of Agricultural Publishing House in Prague, Praga, pp. 33-44, **1983** [In Czech].
48. KMIEĆ G., KASPERCZYK K., ZWOŹDZIAK J., ZWOŹDZIAK A., Evaluation of the concentration and type of pollutants in atmospheric precipitation in the region of the Giant Mountains, Institute of Environmental Engineering, Wroclaw University of Technology **1994** [In Polish].
49. CANDELONE J. P., JAFFREZO J. L., HONG S., DAVIDSON C. I., BOURTON C. F. Seasonal variations in heavy metals concentrations in present day Greenland snow, *Sci. Total. Environ.*, **193**, 101, **1996**.
50. MASCLLET P., HOYA V., JAFFREZO J. L., CACHIER H. Polycyclic aromatic hydrocarbon deposition on the ice sheet of Greenland. Part I: superficial snow, *Atmos. Environ.*, **34**, 3195, **2000**.
51. MASCLLET P., HOYAU V. Evidence for the presence of PAH in the polar atmosphere and in polar ice, *Analisis*, **22**, (7), 31, **1994**.
52. PITTS J. N., VAN CAUWENBERGHE K. A., GROSJEAN D., SCHMIDT J. P., FITZ D. R., BELSER W. L., KNUDSON G. B., HYND S. P. M. Atmospheric reactions of PAH: facile formation of mutagenic nitro-derivatives, *Sci.*, **202**, 515, **1978**.
53. JAMIESON J. B., JOHNSTON C. D. Evaluation of the shear frame test for weak snowpack layers, *Ann. Glaciol.*, **32**, 59, **2001**.

