

Original Research

# ***In vivo* Antioxidant Potential of *Raphanus sativus* Seeds in Rat Kidney Against CCl<sub>4</sub>-Induced Toxicity**

**Irum Shehzadi<sup>1</sup>, Naseer Ali Shah<sup>2\*</sup>, Muhammad Rashid Khan<sup>1</sup>, Muhammad Shuaib<sup>3\*</sup>, Muzammil Shah<sup>4</sup>, Aimal Khan<sup>5</sup>, Hani S.H. Mohammed Ali<sup>4</sup>, Yasir Anwar<sup>4</sup>, Firasat Hussain<sup>6,7</sup>, Hayat Khan<sup>6</sup>, Imtiaz Ali Khan<sup>8</sup>, Imdad Kaleem<sup>2</sup>, Asghar Shabbir<sup>2</sup>, Naveeda Riaz<sup>9</sup>, Asad ul-Haq<sup>10</sup>, Yaseer Rasheed<sup>2</sup>, Mahnoor Khan<sup>2</sup>, Adeel Sher<sup>2</sup>, Ijaz Ali<sup>2</sup>, Ghazala Shaheen<sup>11</sup>**

<sup>1</sup>Department of Biochemistry, Faculty of Biological Sciences, Quaid-i-Azam University Islamabad, 45320, Pakistan

<sup>2</sup>Department of Biosciences, COMSATS University Islamabad, 45550 Pakistan

<sup>3</sup>School of Ecology and Environmental Science, Yunnan University, No. 2 North Cuihu Road, Kunming, Yunnan, 650091, P.R. China

<sup>4</sup>Department of Biological Sciences King Abdulaziz University, Jeddah 21589, Saudi Arabia

<sup>5</sup>State Key Laboratory of Plant Genomics, Institute of Genetics and Developmental Biology, Chinese Academy of Science, Beijing 100101, China

<sup>6</sup>Department of Microbiology, University of Swabi, Ambar 23561, Pakistan.

<sup>7</sup>Department of Microbiology, Cholistan University of Veterinary & Animal Sciences Bahawalpur 63100, Pakistan.

<sup>8</sup>Department of Entomology, the University of Agriculture Peshawar, Pakistan

<sup>9</sup>Department of Bioinformatics and Biotechnology, International Islamic University Islamabad Pakistan

<sup>10</sup>Department of Radiology, Samsung Medical Center Irwon Dong Gangnam gu Seoul Korea

<sup>11</sup>Department of Botany, University of Balochistan, Quetta, Pakistan

Received: 26 April 2018

Accepted: 20 August 2018

## **Abstract**

In this study, the methanolic extract of *Raphanus sativus* (RSME) seeds was evaluated for its protective effect against CCl<sub>4</sub>-induced nephrotoxicity. The treatment of Swiss albino rats with intraperitoneal injections of CCl<sub>4</sub> (1 ml/kg body weight) on alternate days for 30 days decreased the antioxidant enzymes, while lipid peroxidation and serum toxicity markers were increased. These changes were reversed in the animals receiving an oral dose of RSME (100 and 200 mg/kg body weight) along with CCl<sub>4</sub>, thus increasing the level of antioxidant enzymes and decreasing serum toxicity markers and thus ameliorating the toxic effect of CCl<sub>4</sub>. These results show that seed extract of *R. sativus* can reciprocate the toxic effects of CCl<sub>4</sub> and can be used to make a chemopreventive drug.

**Keywords:** antioxidant potential, *Raphanus sativus*, seeds, rat kidney, CCl<sub>4</sub>-induced toxicity

---

\*e-mail: zeyadz44@yahoo.com

## Introduction

Those reactive chemical species that have only one unpaired electron in their outer shell are known as free radicals [1]. Due to the unstable configuration in their outer orbit, an energy is generated which is then released by reacting with nearby biomolecules, for example carbohydrates, nucleic acids, proteins and lipids. Oxygen-free radicals and nitrogen-free radicals, more commonly known as “reactive oxygen species” (ROS) and reactive nitrogen species (RNS), are the major free radicals causing damage to cells and tissues. Hydroxyl radical, superoxide anion and hydrogen peroxide are included in ROS and peroxy nitrite and nitric oxide are included in RNS. These free radicals can start self-catalyzed reactions in such a way that substances to which they react are also transformed into free radicals in order to spread the chain of reactions causing damage. When the concentration of these reactive radicals climbs up, they cause the destruction of cell structures, including damage to membranes, lipids, nucleic acids and proteins. This type of damage caused by free radicals is called as “oxidative stress” [2].

According to some researchers, the major reductant of oxygen is present in the mitochondrial membrane and it is supposed to be “ubisemiquinone”. Mitochondria is a main physiological source of superoxide radicals in living cells, as the production of this radical is more than 2–3 nmol/min per mg of protein inside mitochondria [3]. Macrophages, neutrophils, xanthine oxidase, peroxisomes, cytochrome P450, microsomes and eosinophils are some sources of free radicals inside the cell [4]. Radiations, metals, xenobiotics, ions, chlorinated compounds and various environmental agents are the exogenous sources of free radicals Valko et al. (2006).

Although an antioxidant defense system is present in the cell, oxidative injuries add up throughout the life cycle, which includes damage to lipids, proteins and DNA. Various age-dependent diseases, for example arthritis, cancer, neurodegenerative disorders and arteriosclerosis, can be caused due to the accumulation of oxidative stress [5].

For thousands of years, nature has been a source of medicinal agents and a very large number of drugs have been derived from these natural sources, among them many drugs that are based on their use in traditional medicine. Natural compounds offer a huge number of antioxidants, especially those originating from dietary sources [6].

*Raphanus sativus* is generally known as radish and it is commonly available in all parts of the world. Various parts of *Raphanus sativus*, such as leaves, seeds and roots, are also used in traditional medicines [7]. Leaves of radish are culturally being used as an appetizer, laxative, digestive aid, stimulant and in various stomach disorders. Old roots, leaves and seeds are also used in the treatment of chest complaints and asthma [8]. The juice extracted from fresh leaves of radish is diuretic. Its

seeds are stomachic, carminative, laxative and diuretic. As the roots are antispasmodic, diuretic, antiscorbutic, digestive and astringent, it is used for the cure of acid regurgitation, indigestion, diarrhea, abdominal bloating and bronchitis [9]. The main objective of this study is to evaluate the preventive effect of *Raphanus sativus* seeds methanolic extract against CCl<sub>4</sub>-induced nephrotoxicity as an antioxidant.

## Materials and Methods

### Plant Collection

The seeds of *Raphanus sativus* (L.) were collected from District Multan, Pakistan. After collection, seeds (2 kg) were washed with distilled water and dried at room temperature in shade for more than two weeks and ground using an electric grinder.

### Preparation of Seed Extract

*Raphanus sativus*'s extract was prepared by the addition of 4 liters of methanol in powdered seeds (2 kg) with occasional shaking. The extract was filtered using Whatmann filter paper No. 45 after a week. A rotary vacuum evaporator was used to evaporate the filtrate and to obtain a methanolic extract of *Raphanus sativus* (RSME).

### Experimental Design

Six-week old albino rats having weight 160-210 g were obtained from the animal house of the National Institute of Health (NIH) in Islamabad. The animals were kept at the primate facility in Quaid-i-Azam University, Islamabad in regular cages. They were kept at room temperature of 25±5°C with a 12 h light/dark cycle. Before dosing, animals were familiarized with the environment for two weeks and fed with standard laboratory food. Before any treatment initial weights were measured.

### Animal Treatment

For the experiment albino rats were distributed into seven groups such that each group contains six animals. The distribution of groups was as follows:

- Group 1 (control group) without any treatment.
- Group 2 (vehicle control) was given (1 ml/kg animal body weight) corn oil only.
- Group 3 received intraperitoneally solution of 20% CCl<sub>4</sub> in corn oil (1 ml/kg animal body weight).
- Group 4 was given intraperitoneally 20% CCl<sub>4</sub> in corn oil (1 ml/kg rat body weight) and silymarin solution (100 mg/ml in DMSO).
- Group 5 was administrated intraperitoneally 20% CCl<sub>4</sub> in corn oil (1 ml/kg rat body weight) and was also given RSME (100 mg/kg rat body weight)



Table 2. Preventive effect of RSME on urinary (pH, specific gravity, WBCs and RBCs) and serum markers (creatinine, urea).

Group	Treatment	Urinary Markers				Serum Markers			
		pH	Specific gravity	WBC/ul	RBC/ul	LDL (mg/dl)	Creatinine (mg/dl)	Urea(mg/dl)	CK-NAC
I	Control	7.1±0.11**	1.3±0.52**	17.6±1.26**	0.0±0.0**	297.24±6.67**	28.85±0.43**	35.896±1.2**	15.946±0.529**
II	DMSO +Olive oil	7.0±0.21**	1.4±0.05**	19.8±1.43**	0.04 ±0.21**	299.31±7.78**	29.56±0.39**	36.537±1.98**	17.139±0.498**
III	1ml/kg CCl <sub>4</sub>	5.9±0.40	3.7±1.26	91.1±1.47	13.2±0.91	987.95±6.72	63.82±0.54	65.71±3.70	31.075±0.756
IV	1ml/kg CCl <sub>4</sub> + silymarin	7.0±0.32*	1.1 ±0.13*	20.3±1.3*	0.09±0.36*	315.28±8.36*	31.85±0.42*	37.491±2.43*	18.437±0.546*
V	100mg/kg RSME+CCl <sub>4</sub>	6.5±0.10*	1.0±0.01*	59.2±1.00	5.2±0.95	810.42±8.67*	37.74±0.47*	48.121±1.4*	25.054±0.395*
VI	200mg/kg RSME+CCl <sub>4</sub>	7.0±0.28**	1.0±0.08**	21.9±1.52**	1.3±0.26*	660.12±14.45	31.97±0.47*	40.23±3.71*	19.97±0.310*
VII	200mg/kg RSME	7.0±0.10**	1.0±0.01**	12.2±1.17*	0.01±0.01**	432.1±24.9*	29.55±0.92**	42.21±8.45*	16.875±0.456**

urobilinogen and urinary protein. Protective effect of RSME against CCl<sub>4</sub> was observed by a reduction in albumin, urobilinogen and protein levels of urine (Table 1).

As compared to control group, CCl<sub>4</sub> treatment causes a significant (P<0.05) decrease in pH of urine while a considerable (P<0.05) increase was found in specific gravity, WBCs and RBCs. The ameliorative effect of RSME against CCl<sub>4</sub> was observed on specific gravity, pH, WBCs and RBCs as pH level was increased (P<0.05) by the treatment of RSME in relation to CCl<sub>4</sub> group, and a decline of specific gravity, WBCs and RBCs count was also found (Table 2).

### Assessing Kidney Serum Toxicity Markers

In serum, the important markers of renal dysfunction are CK-NAC, LDH, creatinine and urea. Significant elevation (p<0.05) was observed when rats were treated with CCl<sub>4</sub>. The preventive effect was shown by RSME as it decreased its level (Table 2).

### In vivo Antioxidant Enzyme Assessment

The ameliorative effect of RSME against CCl<sub>4</sub> was observed by the estimation of TBARS contents and determining antioxidant enzyme activities. As compared to the control group, CCl<sub>4</sub> treatment caused a significant (P<0.05) decrease in CAT, POD and SOD enzyme activity while the increase in TBARS content was found (Table 3). Treatment of RSME caused an increase in activity of these enzymes and decreased the TBARS level. As compared to the control group, CCl<sub>4</sub> treatment caused a significant (P<0.05) decrease in GSR, GST, GSH and GSH-Px (Table 5), while increasing the total protein level. The treatment of RSME in relation to the CCl<sub>4</sub> group increased the enzyme level and decreased protein content.

## Discussion

CCl<sub>4</sub> is experimentally proved to be hepatotoxic and also an active nephrotoxin. The kidney is a multilayered organ comprised of dissimilar machinery that works in an extremely synchronized way. Various chemicals and drugs have been reported to modify the function and structure of kidney [11].

The main objective of the present study was to examine the protective effect of *Raphanus sativus* seed extract against CCl<sub>4</sub>-induced nephrotoxicity in rats, as CCl<sub>4</sub> is a notorious ROS-producing chemical. So, in this study, the antioxidant potential of *Raphanus* methanolic extract was the point of emphasis. After having a look at the results obtained, it is apparent that the seed extract of *Raphanus sativus* is effective in ameliorating the damage caused by CCl<sub>4</sub>. Due to the presence of glucosinolates and antioxidant vitamins, *Raphanus* is



Table 3. Preventive effect of RSME on protein, TBARS content and renal antioxidant enzymes.

Group	Treatment	TBARS ( $\mu\text{g}/\text{mg}$ protein)	CAT (U/min)	POD (units/min)	SOD (U/mg protein)	GSR (nM/min/mg protein)	GST (nM/min/mg protein)	GSH (nM/mg protein)	GSH-Px (nM/mg protein)	Protein ( $\mu\text{g}/\text{mg}$ tissue)
I	Control	44.35 $\pm$ 4.14*	2.73 $\pm$ 0.200**	5.46 $\pm$ 0.28*	11.39 $\pm$ 1.14*	104.31 $\pm$ 2.67**	82.95 $\pm$ 2.02**	32.60 $\pm$ 2.44**	152.16 $\pm$ 1.36**	2.62 $\pm$ 0.14*
II	DMSO + Olive oil	46.43 $\pm$ 3.92**	2.62 $\pm$ 0.2610**	5.21 $\pm$ 0.29**	10.98 $\pm$ 1.23**	103.56 $\pm$ 1.45**	81.34 $\pm$ 1.65**	30.23 $\pm$ 1.78**	151.32 $\pm$ 1.87**	2.67 $\pm$ 0.18*
III	1 ml/kg CCl <sub>4</sub>	106.49 $\pm$ 6.68	1.67 $\pm$ 0.123	2.31 $\pm$ 0.12	6.18 $\pm$ 1.01	76.73 $\pm$ 3.71	42.42 $\pm$ 1.47	12.23 $\pm$ 1.21	82.48 $\pm$ 1.39	4.50 $\pm$ 0.20
IV	1 ml/kg CCl <sub>4</sub> + silymarin	49.52 $\pm$ 4.32*	2.54 $\pm$ 0.349*	4.95 $\pm$ 0.32*	10.27 $\pm$ 1.67*	101.28 $\pm$ 2.45*	78.67 $\pm$ 1.34*	31.52 $\pm$ 1.49*	148.69 $\pm$ 2.05**	2.58 $\pm$ 0.21**
V	100 mg/kg RSME+CCl <sub>4</sub>	66.58 $\pm$ 3.86*	2.18 $\pm$ 0.390*	3.49 $\pm$ 0.40*	8.02 $\pm$ 1.15*	82.38 $\pm$ 1.34	62.60 $\pm$ 1.98*	22.01 $\pm$ 0.97*	124.71 $\pm$ 2.26**	2.38 $\pm$ 0.19*
VI	200 mg/kg RSME+CCl <sub>4</sub>	55.54 $\pm$ 3.31**	2.33 $\pm$ 0.364**	4.79 $\pm$ 0.15*	11.99 $\pm$ 1.82**	101.39 $\pm$ 1.37**	77.37 $\pm$ 2.25*	32.55 $\pm$ 2.37**	143.92 $\pm$ 2.01**	2.57 $\pm$ 0.22**
VII	200 mg/kg RSME	43.22 $\pm$ 2.84**	2.48 $\pm$ 0.300**	5.42 $\pm$ 0.33**	12.16 $\pm$ 1.56*	111.89 $\pm$ 1.74*	83.20 $\pm$ 1.61**	32.25 $\pm$ 1.58**	154.72 $\pm$ 2.60*	2.31 $\pm$ 0.15*

significant in preventing chronic diseases and offering protection against oxidative damage [11, 12].

As described in the results section of this experimental study, the first thing observed after intraperitoneal injection of CCl<sub>4</sub> was the change in body and kidney weights. As compared to untreated animals, a huge decline in body weights had been observed. This might be due to the alteration in metabolic processes inside the body or to the oxidative damage caused to tissues which had turned out to be a difficult for the body weights to increase. The loss in body weight was recovered by the co-administration of seed extract. Whereas, examining kidney weight, an increase in weight was observed in CCl<sub>4</sub>-administrated animals, while RSME showed restorative effects by decreasing the kidney weight toward the normal group.

Analysis of the waste product produced by the kidney, i.e. urine, is the most important factor and a valuable diagnostic tool to monitor the physiological disorders of the kidney [13, 14]. The concentration of various metabolites present in urine was analyzed in this study.

Concerning the function of the liver, kidney status and acid-base balance of body, examination of urine is knowledgeable about the function of the kidney. Under normal circumstances, no protein, albumin, urobilinogen, WBCs and RBCs are excreted in urine, but a high concentration is present under extreme pathological conditions. Urobilinogen is produced by the conjugation of bilirubin and breakdown products of bacteria which, after passing through bile ducts, are processed in the intestine and finally converted to urobilinogen [15-17]. The CCl<sub>4</sub> treatment causes a huge increase in the concentration of creatinine, urobilinogen, WBCs and RBCs in urine, indicating severe fibrosis, renal necrosis, kidney toxicity and glomerular damage [18-21]. RSME treatment showed the protective effect by decreasing the number of WBCs and RBCs passing out in urine, and thus CCl<sub>4</sub> intoxication.

Another group of serum markers comprising of gamma-GT, total bilirubin, creatinine and urea also highly increase due to oxidative atmosphere. Creatinine is a waste product formed by slow degradation of creatine phosphate. It is the creatine kinase that catalyzes the formation of creatine phosphate through the transfer of phosphate bond from a high-energy ATP molecule. Under CCl<sub>4</sub>-induced oxidative stress, ATP declines occur due to the blockage of TCA cycle, so increasing the amount of free creatine, whose degradation results in creatinine formation and therefor increasing its serum level [22] – a condition specific to nephritis [23].

In this study, because of the production of free radicals by the metabolism of CCl<sub>4</sub> and tissue damage, urine of the CCl<sub>4</sub>-treated rats exhibited a minor reduction in pH and a rise in specific gravity. Co-administration of *R. sativus* seed extract increases the pH and decreases specific gravity toward normal, as shown by similar results reported by Khan, Rizvi [24] and Khan, Khan [25].

A brownish yellow pigment called bilirubin is found in bile and is produced during the breakdown of waste RBC; usually found in conjugation with albumin. The bilirubin tests are used for identification of liver damage. Increased levels of total or direct bilirubin may be due to jaundice, liver tube blockage or bile ducts, cirrhosis. Our results of this study show a marked increase in these tested parameters. RSME prevented renal injuries with a subsequent restoration of all these parameters discussed above. This study is in relation to the studies of Khan, Khan [26] and Venkatanarayana, Sudhakar [27] who investigated how the administration of CCl<sub>4</sub> caused marked oxidative stress in kidneys. Serum creatinine, total bilirubin, gamma-GT and urea concentrations were significantly higher in the CCl<sub>4</sub>-treated rats.

LDL and albumin are some other parameters that were checked in this experiment. Lipid peroxidation products have been shown to affect a variety of cellular processes. During early stages of oxidation, the antioxidants act to prevent the formation of oxidation products. Once these antioxidants are used up, the formation of lipid hydroperoxides of polyunsaturated fatty acids (PUFAs) appear, which are rapidly reduced to their hydroxide form [28]. As is evident from previous studies, lipid abnormalities are often linked with renal function deterioration [29, 30], and together with this there exists an independent risk factor for renal injury known as hypercholesterolemia [31, 32]. The probable cause for hypercholesterolemia is an increase in oxidative stress, increased formation of oxidized low-density lipoprotein (Ox-LDL), and renal inflammation [33]. Serum albumin is the predominant serum protein, which reflects the synthetic function of the liver. In the present study an increase in all the parameters of lipid abnormalities occurred by CCl<sub>4</sub> administration for 18 days on alternative basis, except HDL and albumin, which decreased considerably by CCl<sub>4</sub> toxicity as compared to rats of the control group. A decrease in albumin is known as hypoalbuminemia. It can be proposed that high cholesterol levels in serum are associated with low HDL and high LDL. *R. sativus* methanolic extract blunted the elevated levels of these parameters when given along with CCl<sub>4</sub>.

A relationship between oxidative stress and renal damage is reported in various studies. In order to protect the body from oxidative stress and damage caused by radicals, a family of antioxidant enzymes exists, but oxidative stress reaches a certain level and the antioxidant defense system present in the body becomes useless and inadequate for recovering the damaging effects of reactive oxygen species [34-36].

In this experimental work, the protecting effects of *R. sativus* seed extract on damaged antioxidant enzymes were observed, such as POD, CAT, SOD, GSH, GSR, GSH-Px, and GST, and TBARS level was also examined.

Many *in vivo* and *in vitro* studies have supported the fact that CCl<sub>4</sub> enhances peroxidation of lipids in tissues by the production of CCl<sub>3</sub> radical produced

by the metabolic conversion carried out by CYP-450 2E1. Adewole, Salako [37] and Khan and Siddique [19]. In this study, after the administration of CCl<sub>4</sub> in rats, the TBARS contents increased significantly. By keeping in mind the results of previous experiments it can be supposed that lipid peroxidation has occurred, showing tissue damage [20, 38]. Various antioxidant constituents such as vitamins and glucosinolates are present in the methanolic seed extract of *R. sativus*, so it counters the effect of CCl<sub>4</sub>-produced toxicity. Similar results regarding the level of TBARS content in CCl<sub>4</sub>-intoxicated rat tissues after treatment with black tea was observed [39].

A similar study was carried out by Khan, Khan [40] in which methanolic extract of *Launaea procumbens* avoided the renal injuries by stabilizing the membranes, scavenging of free radicals, and ameliorating the toxic effects of CCl<sub>4</sub>.

## Conclusion

Results obtained in this study show that RSME can cure the damage caused in renal tissue of rats with CCl<sub>4</sub>. So it can be concluded that in future *Raphanus sativus* can be used to cure toxicity, as no abnormal signs were observed in animals treated with RSME only.

## Acknowledgements

The authors acknowledge HEC for providing funds for our study.

## Conflict of Interest

The authors declare no conflict of interest.

## References

1. SHAH N.A., KHAN M.R., SATTAR S., AHMAD B., MIRZA B. HPLC-DAD analysis, antioxidant potential and anti-urease activity of *Asparagus gracilis* collected from District Islamabad. *BMC Complement and Altern Med.* **14**,347, 2014.
2. POLJSAK B., ŠUPUT D., MILISAV I. Achieving the balance between ROS and antioxidants: when to use the synthetic antioxidants. *Oxidative medicine and cellular longevity.* 2013, 2013.
3. BIRBEN E., SAHINER U.M., SACKESSEN C., ERZURUM S., KALAYCI O. Oxidative stress and antioxidant defense. *World Allergy Organization Journal.* **5**, 9, 2012.
4. VALKO M., RHODES C., MONCOL J., IZAKOVIC M., MAZUR M. Free radicals, metals and antioxidants in oxidative stress-induced cancer. *Chemico-biological interactions.* **160**, 1, 2006.
5. NAZ K., KHAN M.R., SHAH N.A., SATTAR S., NOUREEN F., AWAN M.L. *Pistacia chinensis*: A potent



- tetra chloride-induced hepatopathy in rats. *Int J Pharma Bio Sciences*. **2**, 622, **2011**.
39. FADHEL Z.A., AMRAN S. Effects of black tea extract on carbon tetrachloride-induced lipid peroxidation in liver, kidneys, and testes of rats. *Phytotherapy Research*. **16**, 28, **2002**.
40. KHAN R.A., KHAN M.R., SAHREEN S., BOKHARI J. Prevention of CCl<sub>4</sub>-induced nephrotoxicity with *Sonchus asper* in rat. *Food and Chemical Toxicology*. **48**, 2469, **2010**.