

Original Research

Biochar Mitigates Greenhouse Gas Emissions from an Acidic Tea Soil

Hong Wang¹, Huaiting Yi¹, Xiao Zhang¹, Wei Su¹, Xinwei Li¹,
Yaojun Zhang^{2*}, Xiang Gao³

¹College of Resource and Environment, Anhui Science and Technology University, Chuzhou, China

²School of Life Sciences, Henan University, Kaifeng, China

³Sinochem Agriculture Holdings, China

Received: 18 October 2018

Accepted: 18 November 2018

Abstract

Acidic tea soil is an important greenhouse gas (GHG) emission source. Few studies have been done to investigate the impact of alkaline biochar addition on acidic soil GHG emissions. We carried out a 40-day aerobic incubation experiment to investigate the alkaline biochar amendment on carbon dioxide (CO₂), methane (CH₄), and nitrous oxide (N₂O) emissions from an acidic tea soil under N application. Soil samples were collected in the 0-15 cm layers from a tea orchard of Purple Mountain in Nanjing, Jiangsu Province, China. The results showed that biochar amendment significantly increased soil pH, dissolved organic carbon (DOC), total dissolved nitrogen (TDN), and the ratio of DOC/TDN at the end of incubation. N fertilization increased all three GHG emissions. In contrast, biochar amendment significantly decreased soil CO₂ and N₂O emissions by 7.2-9.3% and 36.3-44.2%, respectively. Although the interaction of biochar and N fertilizer on soil CO₂ and CH₄ emissions were not obvious, N₂O emissions were significantly affected by their interaction. Consistent with CO₂ and N₂O emissions, the net GWP was significantly decreased by biochar addition. Overall, the present study suggests that biochar amendment could be used as an effective management mitigating soil GHG emissions and the net GWP from the acidic tea field soil.

Keywords: biochar, carbon dioxide, nitrous oxide, methane, net GWP

Introduction

Carbon dioxide (CO₂), methane (CH₄) and nitrous oxide (N₂O) are three major greenhouse gases (GHGs) and all of them have increased sharply since 1750 due to human activities. Over a 100-year time period, the global warming potential (GWP) of CH₄ and N₂O are

34 and 298 times greater than CO₂, respectively [1]. Besides, N₂O in the atmosphere is also playing an important role in damaging the stratospheric ozone layer [2]. Agricultural soil is an important source of anthropogenic GHG emissions due to a mass of nitrogen (N) fertilizer application [3]. Reducing agricultural soil GHG emissions has gained attention worldwide because of the GWP of GHGs and maintains the sustainability of agricultural production [4]. Thus, there is an urgency to look for an effective method that can mitigate agricultural soil GHG emissions.

*e-mail: njayujzhang@163.com

Tea (*Camellia sinensis*), an important cash crop, is planted widely in China [5-6]. In addition, as a leaf-harvested crop, N nutrient is vital for increasing tea yield and quality [7]. Therefore, tea fields in China always receive large amounts of nitrogen fertilizer. For instance, annual N fertilizer application rates have always exceeded 450 kg N ha⁻¹ (or even more than 1200 kg N ha⁻¹) on tea plantations in China [6, 8-9], which obviously exceeds the suggested rate of 250-375 kg N ha⁻¹ yr⁻¹ for high yields of tea plantations [10]. Undoubtedly, such high N fertilizer application could result in environmental problems such as soil acidification and high rates of soil GHGs (especially N₂O) emissions [8, 11]. Long-term soil acidity could suppress tea production while enhancing soil N₂O emissions, thus making a negative impact on tea plantation ecosystems [6, 12]. It was reported that soil GHGs (especially N₂O) emissions from tea fields induced by N fertilization, were much higher than those from other crop fields [6, 13-14].

Biochar amendment to a field, always with high pH and rich carbon content, has been well reported as an effective management strategy to counteract soil acidification for sustainable agriculture while reducing soil GHG emissions [15-21]. Biochar plays an important role in accommodating soil processes (e.g., soil nitrification, denitrification and organic matter mineralization), thus it affects soil C and N cycling [22]. The effect of biochar addition on soil GHG emissions has been investigated deeply, but the results were inconsistent. Generally, previous studies showed that soil N₂O emissions could be reduced significantly with biochar addition, particularly in acidic soils [23-26]. However, the effect of biochar amendment on soil CO₂ and CH₄ emissions is a different controversy [27-29]. Thus resulting in poor understanding of how biochar addition affects soil GHG emissions and the related GWP.

Here we examined the effect of an alkaline biochar addition on GHG emissions from an acidic tea soil with N application. Our aim was to evaluate the biochar effect on CO₂, CH₄ and N₂O emissions and the related GWP from acidic tea soil. We hypothesized that 1) biochar would result in a significant reduction in N₂O emissions through increasing soil pH and 2) biochar could be used as an effective supplement to reduce acidic tea soil GWP.

Materials and Methods

Soil Sampling and Biochar

Soil samples were collected in the 0-15 cm layers from a tea orchard of Purple Mountain in Nanjing, Jiangsu Province, China (32°07'N, 118°86'E). Thirty-six soil cores were collected randomly and mixed homogeneously to be representative soil sample. After being air-dried for 15 days, any visible plant detritus

Table 1. Soil physicochemical properties (mean±SE) before the incubation.

Property	Value
Clay (%)	40.67±1.09
Sand (%)	49.17±1.01
Silt (%)	10.16±0.36
Total C (g kg ⁻¹)	13.91±0.34
Total N (g kg ⁻¹)	1.35±0.02
Soil C/N ratio	10.34±0.37
pH, H ₂ O(1:2.5)	4.69±0.03
Dissolved organic C (mg kg ⁻¹)	98.03±5.12
Dissolved organic N (mg kg ⁻¹)	24.02±1.80
NH ₄ ⁺ -N (mg kg ⁻¹)	15.82±1.96
NO ₃ ⁻ -N (mg kg ⁻¹)	29.97±2.00
Bulk density (g cm ⁻³)	1.27±0.02

and fragments were picked out by hand, and the soil samples were then sieved at 2 mm. Soil physicochemical properties are shown in Table 1. The biochar used in this study was produced from wheat straw through low-temperature pyrolysis (500°C) and also ground to pass through a 2-mm sieve. Biochar was characterized by a pH of 10.9. Total N content was 6.1 g kg⁻¹ and organic C content was 467.2 g kg⁻¹.

Incubation

Four treatments were performed in our experiment: control, biochar amendment (+Biochar), N application (+Nitrogen), and biochar plus N amendment (+B&N). Each treatment included four replicates. For each treatment, 100 g of air-dried soil was added to a conical flask with 250 ml space, receiving 4 g (4% w/w) biochar added to each soil, and biochar were thoroughly mixed with the soil. Then distilled water was used to meet the desired soil water content of 60% water-holding capacity (WHC). Thereafter, the flasks were pre-incubated for one week in order to stabilize the microbial activity and thus avoid the undesired microbial peaks [30]. Pre-incubation and the subsequent incubation were performed without light at room temperature (25±1°C) for 40 days. Flasks were hermetically sealed with para film to prevent water evaporation. Every 2 or 3 days, flasks were weighed and distilled water was used to replenish water losses if necessary. At the same time, another group was set up and incubated for measuring soil mineral N (NH₄⁺-N, NO₃⁻-N) content changes at days 3, 5, 10, 20 and 40.

Greenhouse Gases Measurement

Gas sampling was taken by a gas-tight syringe from the heads pace of the flasks after pre-incubation. Gas

sampling was measured after 1, 2, 3, 4, 5, 6, 7, 9, 10, 12, 15, 18, 20, 22, 25, 30, 35 and 40 days of incubation. For each measurement, the heads pace air in the flasks was thoroughly mixed with ambient air for 1 min at a rate of 200 mL min⁻¹ before gas sampling. The flasks were then capped promptly with silicone rubber stoppers, which gave an airtight seal, then kept with butyl rubber 2 h for gas sampling. The ambient air gas sample was used as the initial concentration for calculating the GHG emission rate. After this period the air in the heads pace of incubation flasks were sampled to determine the gas concentration increase. Then flasks were flushed

with ambient air again and kept open after sampling.

Gas samples were analyzed with a gas chromatograph (Agilent 7890A, USA) equipped with two detectors within 6 hours: a flame ionization detector (FID) and an electron capture detector (ECD). CO₂ and CH₄ were detected using FID, and N₂O was detected using ECD. CO₂ was reduced to CH₄ by hydrogen, which occurred in a nickel catalytic converter at 375°C. Purified gas of nitrogen and a gas mixture of argon-methane (5%) were used as the carrier gases for CO₂ and N₂O, respectively.

Soil Chemical Analysis

After incubation, soil samples were extracted with 2 M KCl solution (soil/water ratio of 1:5) and shaken at 200 rev min⁻¹ for 1h at 25°C. Then the soil extracts, after filtration, were used for analyzing: 1) soil dissolved organic carbon (DOC) and total dissolved nitrogen (TDN) (Shimadzu TOC-V csh, TNM-1, Kyoto, Japan), and 2) soil mineral N (NO₃⁻-N and NH₄⁺-N) contents following the two wavelength ultraviolet spectrometry by an ultraviolet spectrophotometer (HITA-CHI U-2900, Japan) [24]. Soil or biochar pH was analyzed in a volume ratio of 1:2.5 (soil or biochar/water) by a PHS-3 C mv/pH detector (Shanghai, China).

Data Analysis

Production of CO₂, CH₄ and N₂O were calculated assuming constant rates of production. The net GWP of GHGs was calculated by converting the production of CH₄ and N₂O into CO₂ equivalents. The net GWP for a 100-year time horizon with inclusion of climate-carbon feedback was calculated using a radiative forcing potential relative to CO₂ of 34 for CH₄ and 298 for N₂O [1]. Differences in cumulative GHGs emissions and chemical characteristics as affected by biochar, fertilizer N and their interactions were examined with a two-way analysis of variance (ANOVA). Linear or nonlinear regression analyses were conducted to examine the dependence of CO₂, CH₄ and N₂O emissions on soil mineral N contents. Statistical analysis of data was performed using SPSS software version 21 for Windows (SAS, 2013). The data are presented as means±standard error (SE).

Results and Discussion

CO₂ Emissions Influenced by Biochar and N Addition

Soil CO₂ emissions showed a distinct variation with incubation progress (Fig. 1a). The highest CO₂ emission was observed during the primary stage of the incubation and then decreased gently, which was probably due to the availability of soil labile C. Large amounts of CO₂ emissions in the initial phase of incubation are probably due to heterotrophic consumption of soil




Fig. 1. Hourly CO₂ a), CH₄ b) and N₂O c) emissions during the 40-day incubation. The error bars indicate the standard errors of means (±SE).

availability labile C caused by biochar addition, while with exhaustion of labile C resulting in slow rates of CO₂ emissions in the final stages of incubation [24, 28]. A priming effect of biochar addition on CO₂ emissions resulting in an emissions peak within three days was observed in this study. CO₂ showed higher emissions at the beginning incubation period with N and biochar amendment than in the following incubation period. Furthermore, relative to no fertilizer N addition (control and +Biochar treatments), N addition (+Nitrogen and +B and +N treatments) stimulated the initial CO₂ fluxes and the cumulative emissions, likely due to the CO₂ produced from urea hydrolysis [22]. On the contrary, biochar addition decreased soil CO₂ emissions by 7.17% and 9.29% with or without N application, respectively (Table 2). Short-term pulses of CO₂ emissions stimulated by biochar amendment have been reported, and ascribed this to additions of labile C accompanied with biochar amendment [24, 31-34]. Soil CO₂ emissions decreased gradually after the emission peaks, and then slowed gently during the last few days of incubation.

Over the whole incubation period, cumulative CO₂ emissions averaged 664.10, 602.38, 744.14, and 690.74 g kg⁻¹ soil for the control, +Biochar, +Nitrogen, and +B and +N treatments, respectively (Table 2). Biochar addition significantly decreased soil CO₂ emissions while N application reduced its depression effect (-7.17% and -9.29% under the N or no N application treatments, respectively). In the present study, biochar amendment decreasing the cumulative soil CO₂ emissions was probably due to the reduction of soil availability N (e.g., soil NH₄⁺-N decreased by 30.7% and 55.8% with or without N application, respectively; soil NO₃⁻-N decreased by 49.7% and 63.3% with or without N application, respectively) and its stable property and C sequestration [15, 35].

CH₄ Emissions Response to Biochar and N Addition

Different from CO₂ emissions, soil CH₄ emissions were shown highly variable with no distinct pattern (Fig. 1b). During the whole incubation period, N application significantly stimulated soil CH₄ emissions mainly due to the urea hydrolysis for methane bacteria [22]. Although there were several sporadic positive CH₄ emission peaks observed in all treatments, the majority of soil CH₄ emissions were negative, which means the soil showed as a net CH₄ oxidation sink. In addition, relative to CO₂ emissions, soil CH₄ emissions fluctuated strongly and showed an opposite effect following biochar amendment associate with or without N application (Fig. 1, Table 2). Biochar amendment could improve soil aeration and decrease an oxic conditions in soil, thus decreasing CH₄ production and increasing its oxidation. Further more, labile C in soils is also obviously influencing methane oxidation [36].

Cumulative CH₄ emissions were affected by N application but unaffected by biochar addition

Table 2. Average cumulative CO₂, CH₄, and N₂O emission rates and net GWP from soils and their changes as well as results of two-way ANOVA during the 40-day aerobic incubation periods. Different letters within each line indicate significant differences for Fisher LSD test ($P < 0.05$).

	CO ₂		CH ₄		N ₂ O		net GWP	
	g CO ₂ kg ^{-1a}	Percent of change (%) ^b	mg CH ₄ kg ⁻¹	Percent of change (%)	mg N ₂ O kg ⁻¹	Percent of change (%)	g CO _{2(eq)} kg ⁻¹	Percent of change (%)
Control	664.10±19.08 ab		-9.14±1.40 b		1.24±0.22 c		664.16±19.06 ab	
+Biochar	602.38±16.44 b	-9.29	-7.68±0.66 b	+15.97	0.79±0.16 c	-36.29	602.35±16.44 b	-9.31
+Nitrogen	744.14±22.46 a		-4.29±0.61 a		4.37±0.25 a		745.29±22.44 a	
+B&+N	690.75±32.25 a	-7.17	-6.71±0.94 ab	-56.41	2.44±0.12 b	-44.16	691.25±37.93 a	-7.25
Biochar (B)	*		NS		***		*	
Nitrogen (N)	**		*		***		**	
B × N	NS		NS		**		NS	
Model	*		*		***		*	

^a Average CO₂, CH₄ and N₂O emission rates and net GWP presented by mean±SE.

^b Percentage of change with positive (+) or negative (-) values indicate stimulating or depressing effect due to biochar addition, respectively.

*, **, and *** indicate statistically significant at the 0.05, 0.01 and 0.001 probability levels by a two-way ANOVA, respectively, NS, not significant.

(Table 2). Biochar increased CH₄ emissions by 15.97% under the control treatment but decreased by 56.41% with N application. The inconsistent effects of biochar (with or without N addition) on soil CH₄ emissions should give more attention, with a focus on better identification and quantification of the carbon input by biochar.

Effects of Biochar and N Addition on N₂O Emissions

N₂O emissions followed a significant temporal variation during the first 20-day incubation period, and the greatest tN₂O emissions (up to 9.89 µg kg⁻¹ h⁻¹) occurred at day 18 followed by the N addition (Fig. 1c). Afterward, N₂O emissions declined rapidly and were kept steadily low until the end of incubation. N₂O emissions were depressed significantly by biochar addition in the first 20-day incubation (-51.1%), but no obvious effect until the incubation finished (-0.9%) with N application. The short-term N₂O emission pulses induced by N addition indicated that N₂O emission peaks occur rapidly and shortly in response to fertilizer N application [24, 37-38]. The cumulative N₂O emissions averaged 1.24, 0.79, 4.37, and 2.44 mg kg⁻¹ soil for the control, +Biochar, +Nitrogen, and +B and +N treatments, respectively. Biochar addition showed a more significantly inhibiting effect on N₂O emissions with N application (-44.16%) rather than no N addition (-36.29%). Biochar amendment decreasing soil N₂O emissions has also been reported in previous studies [24-26], which are mainly attributed to changes in soil C/N ration and aeration, soil microbial community composition and size structures, and microbial enzymes and processes (e.g., nitrification, denitrification) involved in N cycling in soil [25, 39-40]. In general, N₂O emissions were significantly influenced by N fertilizer, biochar and their interaction from acidic tea soil in the present study (Table 2).

Overall Global Warming Potential

Net GWP (t CO₂ equivalent ha⁻¹) was calculated in our study in order to evaluate the mitigation effects of biochar amendment on the combined climatic impacts of CO₂, CH₄ and N₂O emissions in the acidic tea soil. The net GWP was significantly affected by biochar and N application but not their interaction (Table 2). N addition resulted in the greatest GWP (745.29 g CO_{2(eq)}kg⁻¹), while biochar amendment decreased by 7.25%. The least GWP (602.35 g CO_{2(eq)}kg⁻¹) was found in the treatment where soil was only amended with biochar (+Biochar), which was depressed by 9.31% compared with control. The net GWP was positive for all treatments, suggesting that the acidic tea soil acted as an important GHG source. Here, the net GWP significantly increased with N fertilization while decreasing with biochar amendment throughout the incubation period. The obvious decrease in net GWP with biochar amendment was potentially attributed to its C sequestration [15], and thus indicating that biochar amendment could be used as an effective management tool for mitigating the net GWP from the acidic tea field soil.

Soil Characteristic Changes Regulating GHGs Emissions

Biochar and N application significantly affected soil pH while showing no interactions (Table 3). The addition of biochar increased soil pH by more than 1 unit compared with the control treatment. Soil mineral N contents were significantly influenced by both biochar and N addition (Table 3). Soil NH₄⁺-N and NO₃⁻-N contents increased significantly with N application but decreased with biochar addition, which means that biochar amendment could inhibit soil mineralization and nitrification. NO₃⁻-N increased with incubation time while NH₄⁺-N decreased sharply with N application. NH₄⁺-N contents changed rarely and were

Table 3. Soil characteristics as well as results of two-way ANOVA after the 40-day incubation (mean±SE). Different letters within each line indicate significant differences for Fisher LSD test (*P*<0.05).

	pH	NH ₄ ⁺ -N(mg kg ⁻¹)	NO ₃ ⁻ -N(mg kg ⁻¹)	DOC(mg kg ⁻¹)	TDN(mg kg ⁻¹)	DOC/TDN
Control	4.62±0.03 c	20.94±0.93 c	38.76±5.03 bc	126.97±10.62 c	29.48±2.75 c	4.32±0.07 b
+Biochar	5.72±0.02 a	9.26±0.63 d	14.24±2.10 c	177.90±7.74 ab	36.42±1.42 b	4.90±0.22 a
+Nitrogen	4.04±0.01 d	43.03±1.10 a	135.28±13.41 a	152.65±12.00 bc	39.64±2.38 b	3.84±0.11 c
+B& +N	5.29±0.14 b	29.82±1.80 b	68.01±13.32 b	188.48±12.55 a	47.01±1.76 a	4.00±0.13 bc
Biochar (B)	***	***	**	**	**	*
Nitrogen (N)	***	***	***	NS	***	***
B × N	NS	NS	NS	NS	NS	NS
Model	***	***	***	**	***	***

*, **, and *** indicate statistically significant at the 0.05, 0.01 and 0.001 probability levels by a two-way ANOVA, respectively; NS, not significant.

kept low in the control and biochar treatments compared with the N treatments.

Soil N_2O is primarily produced through soil nitrification and denitrification processes, which are highly dependent on soil characteristics, such as soil mineral N contents and pH [41-42]. Soil pH, which has been considered a central factor influencing N transformations [43], was increased by biochar addition in acidic soils and might be an important factor

decreasing N_2O emissions in the present study (Table 3). High soil pH induced by biochar addition decreased the N_2O emissions, probably due to the high soil pH destroyed activity of the functional N_2O reductase enzyme, thus depressing the denitrification progress [25, 43-44]. N_2O emissions depend significantly on soil mineral N in the present study (Fig. 2c), which is in accordance with previous studies [45]. In addition, Singh et al. [38] proposed that soil N immobilization by the sorption capacity of biochar could also reduce N_2O emissions. We found that soil NH_4^+ and NO_3^- decreased by 30.7-55.8% and 49.7-63.3%, respectively, suggesting that N immobilization is an important factor influencing soil N_2O emissions. Generally, soil CO_2 and N_2O emissions were significantly correlated with soil mineral N content during the incubation time (Figs 2a and c).

Biochar addition significantly affected soil DOC, TDN and the ratio of DOC/TDN over the whole incubation period. The interaction of biochar and N showed no effects on soil DOC, TDN and the ratio of DOC/TDN (Table 3).

Conclusions

This study showed that N addition could increase all three GHG emissions and the net GWP in acidic tea soil. However, biochar amendment significantly decreased soil CO_2 and N_2O emissions, and the related net GWP, while showing inconsistent results on CH_4 emissions. The results suggest that biochar amendment (either alone or combined with N) could be used as an effective method for reducing GHG emissions in acidic tea soil.

Acknowledgements

This work was supported by National Natural Science Foundation of China (41701283), the China Postdoctoral Science Foundation (2018M632760), the Natural Science Foundation of Anhui Province in China (16030701102), and the Key University Science Research Project of Anhui Province (KJ2016A173, KJ2019A0819).

Conflict of Interest

The authors declare no conflict of interest.

References

1. IPCC. In: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change (eds T. F. Stocker et al.,) 5-14 (Cambridge University Press, 2013). **2013**.




Fig. 2. Soil CO_2 a), CH_4 b) and N_2O c) emissions dependent on soil mineral N (NH_4^+ -N+ NO_3^- -N) contents during the 40-day incubation.

2. RAVISHANKARA A., DANIEL J., PORTMANN R. Nitrous oxide (N₂O): The dominant ozone-depleting substance emitted in the 21st century. *Science* **326** (5949), 123, **2009**.
3. ZHANG W., DOU Z., HE P., JU X., POWLSON D., CHADWICK D., NORSE D., LU Y., ZHANG Y., WU L., CHEN X., CASSMAN K., ZHANG F. New technologies reduce greenhouse gas emissions from nitrogenous fertilizer in China. *Proc. Natl. Acad. Sci. USA* **110** (21), 8375, **2013**.
4. JU X., XING G., CHEN X., ZHANG S., ZHANG L., LIU X., CUI Z., YIN B., CHRISTIE P., ZHU Z., ZHANG F. Reducing environmental risk by improving N management in intensive Chinese agricultural systems. *Proc. Natl. Acad. Sci. USA* **106** (9), 3041, **2009**.
5. CHENG Y., WANG J., ZHANG J., MÜLLER C., WANG S. Mechanistic insights into the effects of N fertilizer application on N₂O emission pathways in acidic soil of a tea plantation. *Plant Soil*, **389** (1-2), 45, **2015**.
6. YAO Z., WEI Y., LIU C., ZHENG X., XIE B. Organically fertilized tea plantation stimulates N₂O emissions and lowers NO fluxes in subtropical China. *Biogeosciences* **12** (20), 5915, **2015**.
7. TOKUDA S., HAYATSU M. Nitrous oxide flux from a tea field amended with a large amount of nitrogen fertilizer and soil environmental factors controlling the flux. *Soil Sci. Plant Nutr.* **50** (3), 365, **2004**.
8. FU X., LI Y., SU W., SHEN J., XIAO R., TONG C., WU J. Annual dynamics of N₂O emissions from a tea field in southern subtropical China. *Plant Soil Environ.* **58** (8), 373, **2012**.
9. LI Y., FU X., LIU X., SHEN J., LUO Q., XIAO R., LI Y., TONG C., WU J. Spatial variability and distribution of N₂O emissions from a tea field during the dry season in subtropical central China. *Geoderma* **193-194**, 1, **2013**.
10. ZHU T., ZHANG J., MENG T., ZHANG Y., YANG J., MÜLLER C., CAI Z. Tea plantation destroys soil retention of NO₃⁻ and increase N₂O emissions in subtropical China. *Soil Biol. Biochem.* **73**, 106, **2014**.
11. HIRONO Y., NONAKA K. Nitrous oxide emissions from green tea fields in Japan: contribution of emissions from soil between rows and soil under the canopy of tea plants. *Soil Sci. Plant Nutr.* **58** (3), 384, **2012**.
12. BAGGS E., SMALES C., BATEMAN E. Changing pH shifts the microbial source as well as the magnitude of N₂O emission from soil. *Biol. Fert. Soils*, **46** (8), 793, **2010**.
13. AKIYAMA H., YAN X., YAGI K. Estimations of emission factors for fertilizer-induced direct N₂O emissions from agricultural soils in Japan: summary of available data. *Soil Sci. Plant Nutr.* **52**, 774, **2006**.
14. HAN W., XU J., WEI K., SHI Y., MA L. Estimation of N₂O emission from tea garden soils, their adjacent vegetable garden and forest soils in eastern China. *Environ. Earth Sci.* **70** (6), 2495, **2013**.
15. LEHMANN J. Bio-energy in the black. *Front. Ecol. Environ.* **5**, 381, **2007**.
16. WOOLF D., AMONETTE J., STREET-PERROTT A., LEHMANN J., JOSEPH S. Sustainable biochar to mitigate global climate change. *Nat. Commun.* **1** (5), 56, **2010**.
17. WANG J., PAN X., LIU Y., ZHANG X., XIONG Z. Effects of biochar amendment in two soils on greenhouse gas emissions and crop production. *Plant Soil*, **360** (1-2), 287, **2012**.
18. XIANG J., LIU D., DING W., YUAN J., LIN Y. Effects of biochar on nitrous oxide and nitric oxide emissions from paddy field during the wheat growth season. *J. Clean. Prod.* **104**, 52, **2015**.
19. SINGH R., BABU J., KUMAR R., SRIVASTAVA P., SINGH P., RAGHUBANSHI A. Multifaceted application of crop residue biochar as a tool for sustainable agriculture: An ecological perspective. *Ecol. Eng.* **77** (C), 324, **2015**.
20. ZHANG D., YAN M., NIU Y., LIU X., VAN ZWIETEN L., CHEN D., BIAN R., CHENG K., LI L., JOSEPH S., ZHENG J., ZHANG X., ZHENG J., CROWLEY D., FILLEY T., PAN G. Is current biochar research addressing global soil constraints for sustainable agriculture? *Agr. Ecosyst. Environ.* **226**, 25, **2016a**.
21. SUBEDI R., BERTORA C., ZAVATTARO L., GRIGNANI C. Crop response to soils amended with biochar: expected benefits and unintended risks. *Ital. J. Agron.* **12** (2), 161, **2017**.
22. VAN ZWIETEN L., KIMBER S., MORRIS S., DOWNIE A., BERGER E., RUST J., SCHEER C. Influence of biochars on flux of N₂O and CO₂ from ferrosol. *Aust. J. Soil Res.* **48**, 555, **2010**.
23. YANAI Y., TOYOTA K., OKAZAKI M. Effects of charcoal addition on N₂O emissions from soil resulting from rewetting air-dried soil in short-term laboratory experiments. *Soil Sci. Plant Nutr.* **53** (2), 181, **2007**.
24. WANG J., ZHANG M., XIONG Z., LIU P., PAN G. Effects of biochar addition on N₂O and CO₂ emissions from two paddy soils. *Biol. Fert. Soils* **47** (8), 887, **2011**.
25. CAYUELA M., VAN ZWIETEN L., SINGH B., JEFFERY S., ROIG A., SÁNCHEZ-MONEDERO. Biochar's role in mitigating soil nitrous oxide emissions: A review and meta-analysis. *Agr. Ecosyst. Environ.* **191**, 5, **2014**.
26. NELISSEN V., SAHA B., RUYSSCHAERT G., BOECKX P. Effect of different biochar and fertilizer types on N₂O and NO emissions. *Soil Biol. Biochem.* **70** (2), 244, **2014**.
27. ZIMMERMAN A., GAO B., AHN M. Positive and negative carbon mineralization priming effects among a variety of biochar-amended soils. *Soil Biol. Biochem.* **43**, 1169, **2011**.
28. MAUCIERI C., ZHANG Y., MCDANIEL M., BORIN M., ADMAS M. Short-term effects of biochar and salinity on soil greenhouse gas emissions from a semi-arid Australian soil after re-wetting. *Geoderma* **307**, 267, **2017**.
29. OO A., SUDO S., AKIYAMA H., WIN K., SHIBATA A., YAMAMOTO A., SANO T., HIRONO, Y. Effect of dolomite and biochar addition on N₂O and CO₂ emissions from acidic tea field soil. *PLoS ONE* **13** (2), e0192235, **2018**.
30. HUANG Y., ZOU J., ZHENG X., WANG Y., XU X. Nitrous oxide emissions as influenced by amendment of plant residues with different C:N ratios. *Soil Biol. Biochem.* **36** (6), 973, **2004**.
31. LUO Y., DURENKAMP M., DE NOBILI M., LIN Q., BROOKES P. Short term soil priming effects and the mineral is at ion of biochar following its incorporation to soils of different pH. *Soil Biol. Biochem.* **43** (11), 2304, **2011**.
32. CHENG C., LIN T., LEHMANN J., FANG L., YANG Y., MENYAILO O., CHANG K., LAI J. Sorption properties for black carbon (wood char) after long term exposure in soils. *Org. Geochem.* **70** (5), 53, **2014**.
33. FANG Y., SINGH B., SINGH B., KRULL E. Biochar carbon stability in four contrasting soils. *Eur. J. Soil Sci.* **65** (1), 60, **2014**.
34. WANG J., XIONG Z., KUZYAKOV Y. Biochar stability in soil: meta-analysis of decomposition and priming effects. *GCB Bioenergy*, **8** (3), 512, **2016**.

35. LEHMANN J., GAUNT J., RONDON M. Bio-char sequestration in terrestrial ecosystems – A review. *Mitig. Adapt. Strat. Gl.* **11** (2), 403, **2006**.
36. SULLIVAN B., SELMANTS P., HART S. Does dissolved organic carbon regulate biological methane oxidation in semiarid soils? *Global Change Biol.* **19** (7), 2149, **2013**.
37. BEARE M., GREGORICH E., ST-GEORGES P. Compaction effects on CO₂ and N₂O production during drying and rewetting of soil. *Soil Biol. Biochem.* **41** (3), 611, **2009**.
38. SINGH B., HATTON B., SINGH B., COWIE A., KATHURIA A. Influence of biochars on nitrous oxide emission and nitrogen leaching from two contrasting soils. *J. Environ. Qual.* **39** (4), 1224, **2010**.
39. CAYUELA M., SÁNCHEZ-MONEDERO M., ROIG A., HANLEY K., ENDERS A., LEHMANN J. Biochar and denitrification in soils: when, how much and why does biochar reduce N₂O emissions? *Sci. Rep.* **3**, 1732, **2013**.
40. CASE S., MCNAMARA N., REAY S., STOTT A., GRANT H., WHITAKER J. Biochar suppresses N₂O emissions while maintaining N availability in a sandy loam soil. *Soil Biol. Biochem.* **81**, 178, **2015**.
41. DAVIDSON E., KELLER M., ERICKSON H., VERCHOT L., VELDKAMP E. Testing a conceptual model of soil emissions of nitrous and nitric oxides. *Bio Science* **50** (8), 667, **2000**.
42. WRAGE N., VELTHOF G., VAN BEUSICHEM M., OENEMA O. Role of nitrifier denitrification in the production of nitrous oxide. *Soil Biol. Biochem.* **33** (12), 1723, **2001**.
43. MØRKVED P., DORSCH P., BAKKEN L. The N₂O product ratio of nitrification and its dependence on long-term changes in soil pH. *Soil Biol. Biochem.* **39** (8), 2048, **2007**.
44. LIU B., MØRKVED P., FROSTEGÅRD Å., BAKKEN L. Denitrification gene pools, transcription and kinetics of NO, N₂O and N₂ production as affected by soil pH. *FEMS Microbiol. Ecol.* **72** (3), 407, **2010**.
45. ZHANG Y., LIN F., JIN Y., WANG X., LIU S., ZOU J. Response of nitric and nitrous oxide fluxes to N fertilizer application in greenhouse vegetable cropping systems in southeast China. *Sci. Rep.* **6**, 20700, **2016**.