

*Review*

# Environmental Quality and Human Health Risk of Urban Groundwater Sources Based on Hydrochemical Analysis: a Case Study of Suzhou, China

Lijuan Wang<sup>1\*</sup>, Yuezan Tao<sup>1</sup>, Bin Su<sup>2</sup>, Lijun Wang<sup>3</sup>, Peigui Liu<sup>1</sup>

<sup>1</sup>School of Civil and Hydraulic Engineering, Hefei University of Technology, Hefei 230009, China

<sup>2</sup>Anhui and Huaihe Water Conservancy Science Research Institute, Hefei 230009, China

<sup>3</sup>Anhui Survey and Design Institute of Water Resources and Hydropower Company Limited, Hefei 230008, China

*Received: 2 June 2022*

*Accepted: 25 July 2022*

## Abstract

Groundwater is the main water supply in Suzhou City in China. In recent years, heavy metal contamination of urban groundwater sources has become a serious threat to water quality and human health. Groundwater samples from 87 monitoring wells were collected in the study area and analyzed using ICP-MS spectrometry, their major heavy metals concentrations have been analyzed by a series of chemical and statistical indexes for the evaluation of pollution status and drinking suitability. We found that heavy metals are evenly distributed in the groundwater source, 10% of the samples were of excellent quality and 75% were deemed good; however, 70% of the samples exceeded the allowed Mn content. In 62% of the samples, the carcinogenic risk of As exceeded the recommended levels. Based on the non-carcinogenic health risk values, the metals could be ordered as follows: Mn>Fe>Pb>Cu>As>Zn. Mn accounted for 71% of health risk, and the total health risk was mostly due to excess As. Therefore, the levels of As, Mn, and other heavy metal elements in this region warrant urgent regulatory strategies and methods to reduce the Mn concentration should be continuously investigated.

**Keywords:** groundwater, environmental geochemistry, heavy metal, environmental quality, human health risk

## Introduction

Groundwater is the main drinking water source for many cities. The increase in population and rapid

economic development are increasing groundwater pollution, which poses risks to the safety of water consumption from this source.

Contamination with heavy metals is among the greatest hazards to human health [1-6]. Mitchell et al. [7] found that the main heavy metal contaminants in groundwater are As and Mn. The risks of groundwater contamination with As are being widely

---

\*e-mail: wanglj201007@163.com

studied in Bangladesh, India, Pakistan, Vietnam, and China [8-10]; many researchers believe that exposure to As causes serious health problems, including skin and bladder cancer [11-13]. Jedrychowski et al. [14-15] found that exposure to Pb can cause neurological disorders, and other researchers found that exposure to Pb can cause kidney diseases, hypertension, liver crisis, and skin irritation [16-17]. Recent studies also revealed the negative environmental effect of contamination with Fe and Mn [18-19]. Groundwater containing high concentrations of Fe, Mn, Cu, and Zn can also pose health risks [20-22].

To assess the environmental health of groundwater, an evaluation method that aims to estimate the impact degree and the probability of damage from contaminants on human health can be used. It can provide scientific support for health risk management, drinking water safety, and water environment protection [23-27]. Focusing on heavy metals in this evaluation can help evaluate the health risks linked to consumption of the contaminated water and provide a basis for the formulation of water safety policies [28-30].

In this study, we focused on Suzhou City (population 5.32 million people), the largest industrial city in the northern Anhui province, and an important base for production of coal and grain in East China. The city's water supply is heavily dependent on groundwater resources, with 80% of the total water supply (780 million m<sup>3</sup>) being extracted from the ground. This caused over extraction of groundwater in Suzhou City, prompting the local government to issue a restriction order on groundwater exploitation [31-32]. Considering the declining groundwater level and the resulting decrease in its flow, there is a risk of high concentrations of contaminants in extracted water. Aiming to provide an improved scientific basis for the protection of groundwater resources, we evaluated the quality of groundwater, specifically concentrating on risks posed by heavy metal contamination to human health.

## Material and Methods

### Research Area

Suzhou City is located in the northeast of Anhui Province, Central Huaibei Plain, China (33°18'-34°38'N; 116°09'-118°10'E). The groundwater aquifer of Suzhou City is of the porous type, formed of loose rock, and the lithology is mainly silty, consisting of silty sand, sub-sand, local fine sand, and silt. The aquifer is vertical and can be divided into three aquifer groups. For this study, samples were collected from the first aquifer.

### Sample Collection and Analysis

Samples were collected from 87 monitoring wells, allowing the sampling to cover the entire study area (Fig. 1). The sampling points covered both centralised and decentralised water sources. The measuring instrument used was the inductively coupled plasma mass spectrometer (ICP-MS)(NexION 300, USA). Laboratory water was prepared for the Millipore ultrapure water preparation device. The groundwater samples included in the current study were collected following the norms and standards prescribed by the Ministry of Environmental Protection of China (MEP) [33] on sampling collection, transportation, storage, preparation, and instrumental analysis. The method detection limit (LD) was 0.12 µg/L for As, 0.11 µg/L for Cr, 0.08 µg/L for Cu, 0.82 µg/L for Fe, 0.12 µg/L for Mn, 0.09 µg/L for Pb, and 0.67 µg/L for Zn. The test results are shown in Table 1.

### Hydrogeochemical Analyses

Hydrogeochemical analysis is the analysis of content and spatial distribution. Analysis of heavy metal hydrogeochemical is based on mathematical statistics




Fig. 1. Sampling locations within of the study area.

Table 1. Heavy metal concentrations in groundwater ( $\mu\text{g/L}$ ).

Sampling point	Mn	Cu	Zn	Pb	Fe	As	Sampling point	Mn	Cu	Zn	Pb	Fe	As
1	45	1.48	14.49	0.52	15.92	<LD	46	281	1.84	2.92	0.42	130	<LD
2	104	0.44	12.57	0.25	169.4	0.45	47	43.03	1.08	7.59	0.10	177	2.45
3	92	0.54	85.95	<LD	19.59	0.60	48	561.7	0.79	4.38	<LD	193	12.01
4	4	1.05	17.26	0.23	30.82	1.10	49	98.71	1.31	6.20	0.28	35	5.16
5	544	0.16	392.20	0.48	471.5	<LD	50	176.7	0.47	2.13	<LD	61	0.62
6	223	0.68	33.98	0.60	79.69	0.72	51	143.1	0.80	5.07	0.19	26	<LD
7	50	0.23	11.03	0.44	183.3	<LD	52	463.1	0.41	8.01	0.24	70	0.63
8	2171	0.53	8.83	0.17	182	0.48	53	68.99	0.26	4.62	<LD	70	0.54
9	5163	1.55	3.69	<LD	1229	<LD	54	386	0.67	2.60	0.25	141	0.80
10	291	0.87	12.92	0.19	36.83	<LD	55	614	1.67	7.77	0.19	20	0.88
11	181	1.82	8.98	<LD	222.1	0.93	56	102	0.26	1.53	0.76	61	0.82
12	128	0.13	85.58	0.09	98.44	0.50	57	260	0.12	<LD	<LD	79	0.69
13	181	0.27	264.40	<LD	81.27	1.01	58	8	0.29	14.77	0.50	111	5.56
14	229	<LD	1.34	0.60	412.8	0.69	59	29	<LD	<LD	<LD	6	<LD
15	251.9	0.37	8.33	<LD	32.04	<LD	60	305	0.68	<LD	<LD	65	0.65
16	81.52	3.52	8.84	<LD	1.543	<LD	61	217	3.69	8.32	0.34	116	<LD
17	347.6	0.60	17.17	0.20	70.27	0.60	62	310	0.40	7.16	0.65	5	1.46
18	103.4	0.36	49.38	0.19	12	0.50	63	249	0.26	<LD	<LD	10	0.71
19	424.7	1.22	18.33	0.21	13.36	0.90	64	46.48	0.13	4.84	<LD	3	1.52
20	441.2	0.68	25.45	1.00	128.7	1.90	65	77.94	1.09	279.90	0.20	76	<LD
21	89.68	0.69	20.27	2.07	73.37	<LD	66	1311	0.86	21.33	0.36	15	1.60
22	136.4	0.35	13.13	0.53	49.7	<LD	67	164.3	0.62	14.52	0.60	146	<LD
23	138.8	1.44	20.46	<LD	28.86	0.42	68	261.4	1.11	5.55	0.80	94	1.17
24	37.84	1.14	14.40	<LD	36.82	<LD	69	240.6	0.69	10.05	<LD	315	0.35
25	1012	1.61	10.85	0.50	89.12	1.82	70	133.6	0.21	8.25	0.50	569	<LD
26	226.3	0.43	4.83	<LD	6.706	0.35	71	206.8	2.38	30.62	<LD	68	1.59
27	45.23	1.06	19.85	<LD	43.29	<LD	72	995.5	0.77	14.57	0.30	81	0.46
28	132.3	0.83	20.08	0.12	156	<LD	73	204.2	0.42	4.97	0.52	93	<LD
29	307.2	0.81	<LD	<LD	71.55	0.50	74	71.3	0.10	<LD	0.20	11	0.48
30	781.3	1.01	12.25	0.16	54.07	0.80	75	2.171	0.25	3.79	0.36	<LD	<LD
31	10.77	1.96	138.90	0.26	35.28	0.60	76	895.7	0.36	14.61	0.88	71	2.70
32	228.1	0.59	18.82	0.86	156.1	1.70	77	139.1	0.32	3.12	0.70	186	<LD
33	476.6	0.25	<LD	<LD	294.9	<LD	78	222.1	0.09	61.33	0.82	175	0.62
34	396.7	1.72	16.53	0.17	233.4	<LD	79	204.4	0.08	1.71	0.33	69.92	<LD
35	71.32	0.55	14.36	0.20	224.7	0.40	80	61.34	0.13	1.95	0.20	2.979	0.80
36	243.2	1.35	12.45	0.22	59.33	<LD	81	195.8	<LD	1.30	0.33	99.13	<LD
37	282.9	0.75	3.74	<LD	39	<LD	82	174.7	1.17	6.17	<LD	39.56	<LD
38	404.4	5.23	10.19	2.14	110	0.48	83	60.48	0.42	44.37	<LD	388.1	1.30
39	745.9	0.86	2.43	<LD	81	10.36	84	71.24	0.10	9.93	<LD	467.3	1.50

Table 1. Continued.

40	30.49	0.72	1.51	0.40	6	<LD	85	561.5	0.35	8.89	<LD	<LD	0.90
41	478.1	0.40	3.36	<LD	205	9.86	86	172.8	0.67	184.10	2.18	32.42	<LD
42	188.3	0.67	1.83	<LD	136	0.87	87	58.79	<LD	2.15	<LD	21.96	<LD
43	19.55	1.35	8.20	1.89	7	<LD							
44	77.78	0.64	2.28	<LD	117	2.27							
45	232.5	0.34	2.41	0.30	83	46.90							

of concentration data, and compared the heavy metal concentration in each sample (see Table 2) with the Class III division of the groundwater quality standards [34] that describe the standard required for groundwater to be suitable for use as a centralized domestic and drinking water source.

### Environmental Water Quality Analyses

The water environment is analyzed by the comprehensive index which was evaluated based on the Nemerow index method [35]. Evaluation criteria of the Nemerow index are shown in Table 3.

### Human Health Risk Assessment

The health risk of carcinogens in drinking water was evaluated using the following equation [36]:

$$R_i^c = \frac{1 - \exp(-(CW_i \cdot IR) / BW \cdot q_i)}{L} \tag{3}$$

Where  $R_i^c$  is the average annual risk of carcinogenesis for individual person produced by the carcinogen  $i$  in drinking water,  $a^{-1}$ ,  $q_i$  is the carcinogenic potency factor of carcinogen  $i$  in drinking water, mg/(kg.d). A value of 15 for As [37] was used in the present study, and  $L$  is the average human lifespan, which was assumed to be 70 years.  $CW$  is the concentration of the heavy metal in the groundwater (mg/L),  $IR$  is the daily intake of drinking water (a value of 2.2 L/d was used),  $EF$  is the exposure frequency (the value used was 365 days/year),  $ED$  is the exposure duration (70 years for the carcinogen and 30 years for the non-carcinogen),  $BW$  is the body weight in kg (70 kg was selected for the adults in Suzhou City), and  $AT$  is the average exposure time (days), which was calculated as  $365 \times ED$ .

The health risk of non-carcinogens in drinking water was assessed using the following equation:

$$R_j^n = \frac{CW_j \cdot IR \cdot 10^{-6}}{RfD_j \cdot L \cdot BW} \tag{4}$$

Table 2. Classification of groundwater by quality.

Order number	Class	Scope of application
1	I class	Various uses
2	II class	Various uses
3	III class	Centralized domestic and drinking water sources, industrial and agricultural water use
4	IV class	Used for agriculture and in the industry, but can be used for drinking after appropriate treatment
5	V class	Choose according to the purpose of use

Table 3. Groundwater quality classification standard and single-component scoring standard.

Water quality classification	Individual component score values $F_i$	The Memero Pollution Index, F	Water quality classification
I	0	<0.80	Excellent
II	1	0.80~2.50	Good
III	3	2.50~4.25	Better
IV	6	4.25~7.20	Poor
V	10	>7.20	Extremely Poor

Table 4. Reference values of the risk level ( $a^{-1}$ ).

Institution name	Maximum acceptable level	Negligible level	Remarks
The Swedish Environmental Protection Agency	$1.0 \times 10^{-6}$	-	Chemical pollutant
The Dutch Ministry of Construction and the Environment	$1.0 \times 10^{-6}$	$1.0 \times 10^{-8}$	Chemical pollutant
Royal Society	$1.0 \times 10^{-6}$	$1.0 \times 10^{-7}$	-
International Commission on Radiological Protection	$5.0 \times 10^{-5}$	-	Radiation
US Environmental Protection Agency	$1.0 \times 10^{-4}$	-	-

Where  $R_i^n$  is the average annual risk to individuals by non-carcinogens  $i$  in drinking water ( $a^{-1}$ ),  $RfD_i$  is the reference dose of the non-carcinogen  $i$  in drinking water, mg/(kg.d). The value of  $RfD_i$  is 0.02 for As,  $5 \times 10^{-3}$  for Cu, 0.3 for Fe,  $1.4 \times 10^{-1}$  for Mn,  $1.4 \times 10^{-3}$  for Pb, and  $3.0 \times 10^{-1}$  for Zn.

The total annual risk of carcinogenesis caused by the carcinogens in groundwater was calculated using Eq. (5). The reference values that we used for each risk level were given by the international standards of various institutions, as given by Table 4.

$$R = \sum_{i=1}^n R_i^c + \sum_{i=1}^n R_i^n \quad (5)$$

## Results and Discussion

### Hydrogeochemical Analyses

#### Content Analysis

Content characteristics analysis of 87 sampling sites in the study area revealed that the content of the six heavy metals in the samples were ranked as Mn>Fe>Zn>As>Cu>Pb (Table 1).

#### Mn

The average Mn content was 330  $\mu\text{g/L}$ , the maximum Mn concentration was 5163  $\mu\text{g/L}$ , and the coefficient of variation was 185%. 61 sampling points (accounting for 70%) contained excessive amounts of Mn, and the maximum excess multiple was 51.36.

Mn is a common natural groundwater contaminant and is present at concentrations much higher than the recommended drinking water guidelines [38]. Superfluous Mn has been found in groundwater in many countries worldwide, such as western Quebec in Canada [39], the lower Kelantan River Basin on the north-eastern coast of Malaysia [40], the Balat-Teneida area in the El-Dakhla Basin in Egypt [41], and Hidalgo in Mexico [42]. Mn also exceeds the standards in groundwater in some cities in China. Mn and Fe are the main contaminants of the groundwater in Suzhou

[43] and excessive amounts of Mn have been found in the southern and northeastern regions of China [44-46].

#### Fe

The average iron (Fe) content was 121  $\mu\text{g/L}$ , the maximum Fe concentration was 1229  $\mu\text{g/L}$ , and the coefficient of variation was 137 %. The concentration value of each sample was compared with the Class III division of the People's Republic of China Quality Standard for Drinking Water, 7 sampling points (accounting for 8%) contained excessive amounts of Fe, maximum excess multiple was 4.10.

Groundwater contamination occurs owing to the ability of water to dissolve aquifers; any change in water acidity and alkalinity affects this process, and Mn and Fe are very sensitive to water redox conditions [47]. Highly industrialised cities are also at risk of exceeding the accepted Mn and Fe concentration limit in groundwater [48]. This previous study also showed that excessive quantities of Mn and Fe in the groundwater of Suzhou are linked to human activities (such as the excessive exploitation of groundwater and mineral resources).

#### As

The average Arsenic (As) content was 2.51  $\mu\text{g/L}$ , the maximum As concentration was 46.9  $\mu\text{g/L}$ , and the coefficient of variation was 261 %. The concentration value of each sample was compared with the Class III division of the People's Republic of China Quality Standard for Drinking Water, 3 sampling points (accounting for 3%) contained excessive amounts of As, maximum excess multiple was 4.69. Monitoring of these three Wells should be strengthened.

#### Zn, Cu, Pb

The average Zinc (Zn) content was 28.44  $\mu\text{g/L}$ , the maximum Zn concentration was 392.2  $\mu\text{g/L}$ . The average Copper (Cu) content was 0.85  $\mu\text{g/L}$ , the maximum Cu concentration was 5.23  $\mu\text{g/L}$ . The average lead (Pb) content was 0.51  $\mu\text{g/L}$ , the maximum Pb concentration was 2.18  $\mu\text{g/L}$ . The contents of these three metals are lower than the limit of the Class III division




Fig. 2. Concentration spatial distribution of heavy metals.

of the People's Republic of China Quality Standard for Drinking Water.

#### *Spatial Characteristic Analysis*

The spatial distribution characteristics can directly reflect the concentration value of heavy metals in a specific area. Mn was generally high in the whole groundwater sources, and there are one point where the Mn was very high, it could be that the well was seriously contaminated (Fig. 2). The sampling point with the highest concentration was found in south part

of Suzhou City where located in the Sunan mining area, there is a high possibility that the water quality at this sampling point has been disturbed by human activities. Fe in the same area shows high concentration as Mn. This result may contribute to their common tendency in redox changes and similar geochemical features [49], and it could be its mixed lithologic and anthropogenic origins [50]. The concentration of six heavy metals varies gently in regional distribution, except for a few points. The uniform concentration distribution indicates that the water source is not affected by regional pollution [51].

## Comprehensive Quality Evaluation

The quality of water samples evaluated using the Nemerow method is shown in Table 5. Among the evaluated sampling sites, 9 sites (10 %) had an excellent water quality classification, 65 sites (75 %) had a good water quality classification, 11 sites (13 %) had a poor water quality classifications, and 2 sites (2 %) had an extremely poor water quality classification. These results are similar to the data given in the Suzhou Water Resources Bulletin and suggests that the study area generally has good water quality. The groundwater

water type in Suzhou city is medium to deep pore water and carbonate karst water, and the system is mainly semi-closed. Because atmospheric precipitation does not seep directly into the sampling wells, and because contaminants from surface water are absorbed by rock and adsorbed by gas, the quality of groundwater is effectively preserved. There are a few water source wells where the water intake layers are closely related to rainfall or surface water. This runoff water may be affected by human activities, causing groundwater pollution.

Table 5. The calculation of F value.

Sample point	$F_{\max}$	$\bar{F}$	$F$	Sample point	$F_{\max}$	$\bar{F}$	$F$
1	0	0.0	0.0	46	3	1.0	2.2
2	3	1.0	2.2	47	3	1.0	2.2
3	3	1.0	2.2	48	6	2.3	4.5
4	3	0.9	2.2	49	3	1.3	2.3
5	6	2.3	4.5	50	3	0.9	2.2
6	3	0.9	2.2	51	3	0.9	2.2
7	1	0.3	0.7	52	3	0.9	2.2
8	10	3.0	7.4	53	3	0.9	2.2
9	10	3.7	7.5	54	3	1.0	2.2
10	3	0.9	2.2	55	3	0.9	2.2
11	3	1.3	2.3	56	3	0.9	2.2
12	3	1.0	2.2	57	3	0.9	2.2
13	3	1.4	2.3	58	3	1.0	2.2
14	6	2.1	4.5	59	0	0.0	0.0
15	3	0.9	2.2	60	3	0.9	2.2
16	3	0.9	2.2	61	3	1.0	2.2
17	3	0.9	2.2	62	3	1.3	2.3
18	3	0.9	2.2	63	3	0.9	2.2
19	3	0.9	2.2	64	3	0.9	2.2
20	3	1.4	2.3	65	3	1.0	2.2
21	3	0.9	2.2	66	6	2.1	4.5
22	3	0.9	2.2	67	3	1.0	2.2
23	3	0.9	2.2	68	3	1.3	2.3
24	0	0.0	0.0	69	6	2.1	4.5
25	6	2.1	4.5	70	6	2.1	4.5
26	3	0.9	2.2	71	3	1.3	2.3
27	0	0.0	0.0	72	3	0.9	2.2
28	3	1.0	2.2	73	3	0.9	2.2
29	3	0.9	2.2	74	3	0.9	2.2

Table 5. Continued.

30	3	0.9	2.2	75	0	0.0	0.0
31	1	0.3	0.7	76	3	1.3	2.3
32	3	1.4	2.3	77	3	1.0	2.2
33	3	1.3	2.3	78	3	1.1	2.3
34	3	1.3	2.3	79	3	0.9	2.2
35	3	1.3	2.3	80	3	0.9	2.2
36	3	0.9	2.2	81	3	0.9	2.2
37	3	0.9	2.2	82	3	0.9	2.2
38	3	1.0	2.2	83	6	2.6	4.6
39	6	2.1	4.5	84	6	2.6	4.6
40	0	0.0	0.0	85	3	0.9	2.2
41	3	1.7	2.4	86	3	1.0	2.2
42	3	1.0	2.2	87	3	0.9	2.2
43	0	0.0	0.0				
44	3	1.4	2.3				
45	6	2.1	4.5				

## Human Health Risk Assessment

### *Carcinogenic Health Risks Assessment*

Of the total sampling points, 33 sampling points measured at a 0 value for carcinogenic risk. However, our results indicated that the cancer risk from As in the study area was high. Among the evaluated sampling sites, 54 sites (62 %) exceeded the level of maximum acceptable carcinogenic risk recommended by the Swedish Environmental Protection Agency, the Ministry of Construction and Environment in the Netherlands and the Royal British Association, 6 sites (7 %) exceeded the level of maximum acceptable carcinogenic risk recommended by the ICRP, and 4 sites (5 %) exceeded the level of maximum acceptable carcinogenic risk recommended by the US.EPA (see Table 6).

A high concentration of arsenic in groundwater is a worldwide problem, high concentration of arsenic in groundwater has been documented as a major health issue around the globe. Exposure to arsenic poses a serious risk of different types of cancer [52-56], and on several occasions the presence of As has been reported to have caused serious threat to health by causing keratosis and melanosis [57].

Arsenic (As) contamination in different areas of the world has been reported, more than 230 million people worldwide, which include 180 million from Asia, are at risk of arsenic poisoning. Southeast Asian countries, Bangladesh, India, Pakistan, China, Nepal, Vietnam, Burma, Thailand and Cambodia, are the most affected [58]. Arsenic has been found in groundwater in many

Chinese cities, Northern China has been identified as high-risk area. Our results are supported by several published studies, where arsenic pollution has been measured in the groundwater around Suzhou [69-62].

High arsenic groundwater is mainly distributed in the areas with reducing environment [63]. Many scholars believe that arsenic aggregation is caused by human activities that use arsenic-containing minerals, and after release in the environment through a redox reaction, arsenic mixes with surface water to contaminate groundwater [64-69].

Considering the carcinogenic health risks caused by As in Suzhou, we recommend that research efforts are directed towards the development of countermeasures that reduce the adverse impact of human activities on the environment.

### *Non-Carcinogenic Health Risks Assessment*

The non-carcinogenic health risks ranged between  $3.20 \times 10^{-13}$  and  $1.66 \times 10^{-8}$  for Mn, Cu, Zn, Pb, Fe, and As (Table 6). Only 1 site measured a health risk value for Mn that exceeded the level set by the Dutch Ministry of Construction and Environment, however this value was still within the acceptable limits of the recommendations set by other institutions. The calculated health risk values can be ranked as Mn>Fe>Pb>Cu>As>Zn, among which Mn presented as the greatest risk, measuring at 71 % of the total risk (Fig. 3). Although Mn is a common element in groundwater, in excessive quantities it poses a hazard to human health. Chronic exposure to excessive Mn levels causes increased infant mortality in mothers with



Table 6. Health risk ( $a^{-1}$ ).

Sample point	Non-carcinogenic risk						Carcinogenic risk	Total health risk
	Mn	Cu	Zn	Pb	Fe	As	As	
2	$3.34 \times 10^{-10}$	$3.95 \times 10^{-11}$	$1.88 \times 10^{-11}$	$8.02 \times 10^{-11}$	$2.54 \times 10^{-10}$	$1.00 \times 10^{-11}$	$7.01 \times 10^{-6}$	$7.01 \times 10^{-6}$
3	$2.95 \times 10^{-10}$	$4.85 \times 10^{-11}$	$1.29 \times 10^{-10}$	0	$2.93 \times 10^{-11}$	$1.35 \times 10^{-11}$	$9.43 \times 10^{-6}$	$9.43 \times 10^{-6}$
4	$1.28 \times 10^{-11}$	$9.43 \times 10^{-11}$	$2.58 \times 10^{-11}$	$7.38 \times 10^{-11}$	$4.61 \times 10^{-11}$	$2.47 \times 10^{-11}$	$1.73 \times 10^{-5}$	$1.73 \times 10^{-5}$
5	$1.74 \times 10^{-9}$	$1.44 \times 10^{-11}$	$5.87 \times 10^{-10}$	$1.54 \times 10^{-10}$	$7.06 \times 10^{-10}$	0	0	$3.21 \times 10^{-9}$
6	$7.15 \times 10^{-10}$	$6.11 \times 10^{-11}$	$5.09 \times 10^{-11}$	$1.92 \times 10^{-10}$	$1.19 \times 10^{-10}$	$1.62 \times 10^{-11}$	$1.13 \times 10^{-5}$	$1.13 \times 10^{-5}$
7	$1.60 \times 10^{-10}$	$2.07 \times 10^{-11}$	$1.65 \times 10^{-11}$	$1.41 \times 10^{-10}$	$2.74 \times 10^{-10}$	0	0	$6.13 \times 10^{-10}$
8	$6.96 \times 10^{-9}$	$4.76 \times 10^{-11}$	$1.32 \times 10^{-11}$	$5.45 \times 10^{-11}$	$2.72 \times 10^{-10}$	$1.08 \times 10^{-11}$	$7.57 \times 10^{-6}$	$7.58 \times 10^{-6}$
9	$1.66 \times 10^{-8}$	$1.39 \times 10^{-10}$	$5.52 \times 10^{-12}$	0	$1.84 \times 10^{-9}$	0	0	$1.85 \times 10^{-8}$
10	$9.33 \times 10^{-10}$	$7.81 \times 10^{-11}$	$1.93 \times 10^{-11}$	$6.09 \times 10^{-11}$	$5.51 \times 10^{-11}$	0	0	$1.15 \times 10^{-9}$
11	$5.80 \times 10^{-10}$	$1.63 \times 10^{-10}$	$1.34 \times 10^{-11}$	0	$3.32 \times 10^{-10}$	$2.08 \times 10^{-11}$	$1.45 \times 10^{-5}$	$1.45 \times 10^{-5}$
12	$4.10 \times 10^{-10}$	$1.17 \times 10^{-11}$	$1.28 \times 10^{-10}$	$2.89 \times 10^{-11}$	$1.47 \times 10^{-10}$	$1.12 \times 10^{-11}$	$7.82 \times 10^{-6}$	$7.82 \times 10^{-6}$
13	$5.80 \times 10^{-10}$	$2.42 \times 10^{-11}$	$3.96 \times 10^{-10}$	0	$1.22 \times 10^{-10}$	$2.27 \times 10^{-11}$	$1.59 \times 10^{-5}$	$1.59 \times 10^{-5}$
14	$7.34 \times 10^{-10}$	0	$2.01 \times 10^{-12}$	$1.92 \times 10^{-10}$	$6.18 \times 10^{-10}$	$1.54 \times 10^{-11}$	$1.08 \times 10^{-5}$	$1.08 \times 10^{-5}$
15	$8.08 \times 10^{-10}$	$3.35 \times 10^{-11}$	$1.25 \times 10^{-11}$	0	$4.80 \times 10^{-11}$	0	0	$9.02 \times 10^{-10}$
16	$2.61 \times 10^{-10}$	$3.16 \times 10^{-10}$	$1.32 \times 10^{-11}$	0	$2.31 \times 10^{-12}$	0	0	$5.93 \times 10^{-10}$
17	$1.11 \times 10^{-9}$	$5.40 \times 10^{-11}$	$2.57 \times 10^{-11}$	$6.38 \times 10^{-11}$	$1.05 \times 10^{-10}$	$1.35 \times 10^{-11}$	$9.43 \times 10^{-6}$	$9.43 \times 10^{-6}$
18	$3.32 \times 10^{-10}$	$3.20 \times 10^{-11}$	$7.39 \times 10^{-11}$	$6.16 \times 10^{-11}$	$1.80 \times 10^{-11}$	$1.12 \times 10^{-11}$	$7.85 \times 10^{-6}$	$7.86 \times 10^{-6}$
19	$1.36 \times 10^{-9}$	$1.09 \times 10^{-10}$	$2.74 \times 10^{-11}$	$6.67 \times 10^{-11}$	$2.00 \times 10^{-11}$	$2.02 \times 10^{-11}$	$1.41 \times 10^{-5}$	$1.41 \times 10^{-5}$
20	$1.41 \times 10^{-9}$	$6.12 \times 10^{-11}$	$3.81 \times 10^{-11}$	$3.21 \times 10^{-10}$	$1.93 \times 10^{-10}$	$4.27 \times 10^{-11}$	$2.98 \times 10^{-5}$	$2.98 \times 10^{-5}$
21	$2.88 \times 10^{-10}$	$6.20 \times 10^{-11}$	$3.03 \times 10^{-11}$	$6.64 \times 10^{-10}$	$1.10 \times 10^{-10}$	0	0	$1.15 \times 10^{-9}$
22	$4.37 \times 10^{-10}$	$3.17 \times 10^{-11}$	$1.97 \times 10^{-11}$	$1.71 \times 10^{-10}$	$7.44 \times 10^{-11}$	0	0	$7.34 \times 10^{-10}$
23	$4.45 \times 10^{-10}$	$1.29 \times 10^{-10}$	$3.06 \times 10^{-11}$	0	$4.32 \times 10^{-11}$	$9.47 \times 10^{-12}$	$6.63 \times 10^{-6}$	$6.63 \times 10^{-6}$
24	$1.21 \times 10^{-10}$	$1.03 \times 10^{-10}$	$2.16 \times 10^{-11}$	0	$5.51 \times 10^{-11}$	0	0	$3.01 \times 10^{-10}$
25	$3.25 \times 10^{-9}$	$1.45 \times 10^{-10}$	$1.62 \times 10^{-11}$	$1.60 \times 10^{-10}$	$1.33 \times 10^{-10}$	$4.09 \times 10^{-11}$	$2.86 \times 10^{-5}$	$2.86 \times 10^{-5}$
26	$7.26 \times 10^{-10}$	$3.85 \times 10^{-11}$	$7.23 \times 10^{-12}$	0	$1.00 \times 10^{-11}$	$7.81 \times 10^{-12}$	$5.47 \times 10^{-6}$	$5.47 \times 10^{-6}$
27	$1.45 \times 10^{-10}$	$9.48 \times 10^{-11}$	$2.97 \times 10^{-11}$	0	$6.48 \times 10^{-11}$	0	0	$3.34 \times 10^{-10}$
28	$4.24 \times 10^{-10}$	$7.41 \times 10^{-11}$	$3.01 \times 10^{-11}$	$3.72 \times 10^{-11}$	$2.33 \times 10^{-10}$	0	0	$7.99 \times 10^{-10}$
29	$9.85 \times 10^{-10}$	$7.30 \times 10^{-11}$	$3.20 \times 10^{-13}$	0	$1.07 \times 10^{-10}$	$1.12 \times 10^{-11}$	$7.85 \times 10^{-6}$	$7.86 \times 10^{-6}$
30	$2.51 \times 10^{-9}$	$9.10 \times 10^{-11}$	$1.83 \times 10^{-11}$	$5.10 \times 10^{-11}$	$8.09 \times 10^{-11}$	$1.80 \times 10^{-11}$	$1.26 \times 10^{-5}$	$1.26 \times 10^{-5}$
31	$3.45 \times 10^{-11}$	$1.76 \times 10^{-10}$	$2.08 \times 10^{-10}$	$8.21 \times 10^{-11}$	$5.28 \times 10^{-11}$	$1.35 \times 10^{-11}$	$9.43 \times 10^{-6}$	$9.43 \times 10^{-6}$
32	$7.32 \times 10^{-10}$	$5.28 \times 10^{-11}$	$2.82 \times 10^{-11}$	$2.74 \times 10^{-10}$	$2.34 \times 10^{-10}$	$3.82 \times 10^{-11}$	$2.67 \times 10^{-5}$	$2.67 \times 10^{-5}$
33	$1.53 \times 10^{-9}$	$2.22 \times 10^{-11}$	$6.23 \times 10^{-13}$	0	$4.41 \times 10^{-10}$	0	0	$1.99 \times 10^{-9}$
34	$1.27 \times 10^{-9}$	$1.55 \times 10^{-10}$	$2.47 \times 10^{-11}$	$5.45 \times 10^{-11}$	$3.49 \times 10^{-10}$	0	0	$1.86 \times 10^{-9}$
35	$2.29 \times 10^{-10}$	$4.97 \times 10^{-11}$	$2.15 \times 10^{-11}$	$6.54 \times 10^{-11}$	$3.36 \times 10^{-10}$	$9.07 \times 10^{-12}$	$6.35 \times 10^{-6}$	$6.35 \times 10^{-6}$
36	$7.80 \times 10^{-10}$	$1.21 \times 10^{-10}$	$1.86 \times 10^{-11}$	$6.93 \times 10^{-11}$	$8.88 \times 10^{-11}$	0	0	$1.08 \times 10^{-9}$
37	$9.07 \times 10^{-10}$	$6.73 \times 10^{-11}$	$5.60 \times 10^{-12}$	0	$5.84 \times 10^{-11}$	0	0	$1.04 \times 10^{-9}$
38	$1.30 \times 10^{-9}$	$4.70 \times 10^{-10}$	$1.53 \times 10^{-11}$	$6.86 \times 10^{-10}$	$1.65 \times 10^{-10}$	$1.09 \times 10^{-11}$	$7.60 \times 10^{-6}$	$7.61 \times 10^{-6}$
39	$2.39 \times 10^{-9}$	$7.70 \times 10^{-11}$	$3.64 \times 10^{-12}$	$2.79 \times 10^{-11}$	$1.21 \times 10^{-10}$	$2.33 \times 10^{-10}$	$1.62 \times 10^{-4}$	$1.62 \times 10^{-4}$
40	$9.78 \times 10^{-11}$	$6.44 \times 10^{-11}$	$2.25 \times 10^{-12}$	$1.28 \times 10^{-10}$	$8.98 \times 10^{-12}$	0	0	$3.02 \times 10^{-10}$

Table 6. Continued.

41	$1.53 \times 10^{-9}$	$3.56 \times 10^{-11}$	$5.03 \times 10^{-12}$	0	$3.07 \times 10^{-10}$	$2.21 \times 10^{-10}$	$1.54 \times 10^{-4}$	$1.54 \times 10^{-4}$
42	$6.04 \times 10^{-10}$	$6.01 \times 10^{-11}$	$2.74 \times 10^{-12}$	0	$2.04 \times 10^{-10}$	$1.96 \times 10^{-11}$	$1.37 \times 10^{-5}$	$1.37 \times 10^{-5}$
43	$6.27 \times 10^{-11}$	$1.21 \times 10^{-10}$	$1.23 \times 10^{-11}$	$6.05 \times 10^{-10}$	$1.05 \times 10^{-11}$	0	0	$8.11 \times 10^{-10}$
44	$2.49 \times 10^{-10}$	$5.72 \times 10^{-11}$	$3.41 \times 10^{-12}$	0	$1.75 \times 10^{-10}$	$5.10 \times 10^{-11}$	$3.57 \times 10^{-5}$	$3.57 \times 10^{-5}$
45	$7.46 \times 10^{-10}$	$3.09 \times 10^{-11}$	$3.61 \times 10^{-12}$	$9.62 \times 10^{-11}$	$1.24 \times 10^{-10}$	$1.05 \times 10^{-9}$	$7.18 \times 10^{-4}$	$7.18 \times 10^{-4}$
46	$9.01 \times 10^{-10}$	$1.65 \times 10^{-10}$	$4.37 \times 10^{-12}$	$1.35 \times 10^{-10}$	$1.95 \times 10^{-10}$	0	0	$1.40 \times 10^{-9}$
47	$1.38 \times 10^{-10}$	$9.66 \times 10^{-11}$	$1.14 \times 10^{-11}$	$3.21 \times 10^{-11}$	$2.65 \times 10^{-10}$	$5.50 \times 10^{-11}$	$3.85 \times 10^{-5}$	$3.85 \times 10^{-5}$
48	$1.80 \times 10^{-9}$	$7.07 \times 10^{-11}$	$6.55 \times 10^{-12}$	0	$2.89 \times 10^{-10}$	$2.70 \times 10^{-10}$	$1.87 \times 10^{-4}$	$1.87 \times 10^{-4}$
49	$3.17 \times 10^{-10}$	$1.18 \times 10^{-10}$	$9.28 \times 10^{-12}$	$8.82 \times 10^{-11}$	$5.24 \times 10^{-11}$	$1.16 \times 10^{-10}$	$8.08 \times 10^{-5}$	$8.08 \times 10^{-5}$
50	$5.67 \times 10^{-10}$	$4.18 \times 10^{-11}$	$3.18 \times 10^{-12}$	0	$9.13 \times 10^{-11}$	$1.39 \times 10^{-11}$	$9.74 \times 10^{-6}$	$9.74 \times 10^{-6}$
51	$4.59 \times 10^{-10}$	$7.19 \times 10^{-11}$	$7.59 \times 10^{-12}$	$6.16 \times 10^{-11}$	$3.89 \times 10^{-11}$	0	0	$6.39 \times 10^{-10}$
52	$1.49 \times 10^{-9}$	$3.65 \times 10^{-11}$	$1.20 \times 10^{-11}$	$7.63 \times 10^{-11}$	$1.05 \times 10^{-10}$	$1.42 \times 10^{-11}$	$9.96 \times 10^{-6}$	$9.96 \times 10^{-6}$
53	$2.21 \times 10^{-10}$	$2.33 \times 10^{-11}$	$6.91 \times 10^{-12}$	0	$1.05 \times 10^{-10}$	$1.22 \times 10^{-11}$	$8.55 \times 10^{-6}$	$8.55 \times 10^{-6}$
54	$1.24 \times 10^{-9}$	$6.03 \times 10^{-11}$	$3.89 \times 10^{-12}$	$7.86 \times 10^{-11}$	$2.11 \times 10^{-10}$	$1.79 \times 10^{-11}$	$1.25 \times 10^{-5}$	$1.25 \times 10^{-5}$
55	$1.97 \times 10^{-9}$	$1.50 \times 10^{-10}$	$1.16 \times 10^{-11}$	$5.97 \times 10^{-11}$	$2.99 \times 10^{-11}$	$1.98 \times 10^{-11}$	$1.39 \times 10^{-5}$	$1.39 \times 10^{-5}$
56	$3.27 \times 10^{-10}$	$2.31 \times 10^{-11}$	$2.29 \times 10^{-12}$	$2.44 \times 10^{-10}$	$9.13 \times 10^{-11}$	$1.85 \times 10^{-11}$	$1.29 \times 10^{-5}$	$1.29 \times 10^{-5}$
57	$8.34 \times 10^{-10}$	$1.03 \times 10^{-11}$	0	0	$1.18 \times 10^{-10}$	$1.55 \times 10^{-11}$	$1.09 \times 10^{-5}$	$1.09 \times 10^{-5}$
58	$2.57 \times 10^{-11}$	$2.60 \times 10^{-11}$	$2.21 \times 10^{-11}$	$1.61 \times 10^{-10}$	$1.66 \times 10^{-10}$	$1.25 \times 10^{-10}$	$8.72 \times 10^{-5}$	$8.72 \times 10^{-5}$
59	$9.30 \times 10^{-11}$	0	0	0	$8.98 \times 10^{-12}$	0	0	$1.02 \times 10^{-10}$
60	$9.78 \times 10^{-10}$	$6.06 \times 10^{-11}$	0	0	$9.73 \times 10^{-11}$	$1.45 \times 10^{-11}$	$1.02 \times 10^{-5}$	$1.02 \times 10^{-5}$
61	$6.96 \times 10^{-10}$	$3.31 \times 10^{-10}$	$1.24 \times 10^{-11}$	$1.10 \times 10^{-10}$	$1.74 \times 10^{-10}$	0	0	$1.32 \times 10^{-9}$
62	$9.94 \times 10^{-10}$	$3.58 \times 10^{-11}$	$1.07 \times 10^{-11}$	$2.08 \times 10^{-10}$	$7.48 \times 10^{-12}$	$3.27 \times 10^{-11}$	$2.29 \times 10^{-5}$	$2.29 \times 10^{-5}$
63	$7.99 \times 10^{-10}$	$2.31 \times 10^{-11}$	0	0	$1.50 \times 10^{-11}$	$1.59 \times 10^{-11}$	$1.11 \times 10^{-5}$	$1.11 \times 10^{-5}$
64	$1.49 \times 10^{-10}$	$1.16 \times 10^{-11}$	$7.25 \times 10^{-12}$	0	$4.49 \times 10^{-12}$	$3.42 \times 10^{-11}$	$2.39 \times 10^{-5}$	$2.39 \times 10^{-5}$
65	$2.50 \times 10^{-10}$	$9.77 \times 10^{-11}$	$4.19 \times 10^{-10}$	$6.54 \times 10^{-11}$	$1.14 \times 10^{-10}$	0	0	$9.46 \times 10^{-10}$
66	$4.20 \times 10^{-9}$	$7.70 \times 10^{-11}$	$3.19 \times 10^{-11}$	$1.14 \times 10^{-10}$	$2.24 \times 10^{-11}$	$3.60 \times 10^{-11}$	$2.52 \times 10^{-5}$	$2.52 \times 10^{-5}$
67	$5.27 \times 10^{-10}$	$5.55 \times 10^{-11}$	$2.17 \times 10^{-11}$	$1.93 \times 10^{-10}$	$2.19 \times 10^{-10}$	0	0	$1.02 \times 10^{-9}$
68	$8.38 \times 10^{-10}$	$9.92 \times 10^{-11}$	$8.30 \times 10^{-12}$	$2.55 \times 10^{-10}$	$1.41 \times 10^{-10}$	$2.62 \times 10^{-11}$	$1.83 \times 10^{-5}$	$1.83 \times 10^{-5}$
69	$7.72 \times 10^{-10}$	$6.21 \times 10^{-11}$	$1.50 \times 10^{-11}$	0	$4.71 \times 10^{-10}$	$7.81 \times 10^{-12}$	$5.47 \times 10^{-6}$	$5.47 \times 10^{-6}$
70	$4.28 \times 10^{-10}$	$1.91 \times 10^{-11}$	$1.23 \times 10^{-11}$	$1.61 \times 10^{-10}$	$8.52 \times 10^{-10}$	0	0	$1.47 \times 10^{-9}$
71	$6.63 \times 10^{-10}$	$2.13 \times 10^{-10}$	$4.58 \times 10^{-11}$	0	$1.02 \times 10^{-10}$	$3.56 \times 10^{-11}$	$2.49 \times 10^{-5}$	$2.49 \times 10^{-5}$
72	$3.19 \times 10^{-9}$	$6.95 \times 10^{-11}$	$2.18 \times 10^{-11}$	$9.62 \times 10^{-11}$	$1.21 \times 10^{-10}$	$1.03 \times 10^{-11}$	$7.23 \times 10^{-6}$	$7.23 \times 10^{-6}$
73	$6.55 \times 10^{-10}$	$3.74 \times 10^{-11}$	$7.44 \times 10^{-12}$	$1.67 \times 10^{-10}$	$1.39 \times 10^{-10}$	0	0	$1.01 \times 10^{-9}$
74	$2.29 \times 10^{-10}$	$8.53 \times 10^{-12}$	0	$6.41 \times 10^{-11}$	$1.65 \times 10^{-11}$	$1.07 \times 10^{-11}$	$7.48 \times 10^{-6}$	$7.48 \times 10^{-6}$
75	$6.96 \times 10^{-12}$	$2.26 \times 10^{-11}$	$5.67 \times 10^{-12}$	$1.15 \times 10^{-10}$	0	0	0	$1.51 \times 10^{-10}$
76	$2.87 \times 10^{-9}$	$3.27 \times 10^{-11}$	$2.19 \times 10^{-11}$	$2.82 \times 10^{-10}$	$1.06 \times 10^{-10}$	$6.05 \times 10^{-11}$	$4.23 \times 10^{-5}$	$4.23 \times 10^{-5}$
77	$4.46 \times 10^{-10}$	$2.88 \times 10^{-11}$	$4.67 \times 10^{-12}$	$2.24 \times 10^{-10}$	$2.78 \times 10^{-10}$	0	0	$9.82 \times 10^{-10}$
78	$7.12 \times 10^{-10}$	$8.08 \times 10^{-12}$	$9.18 \times 10^{-11}$	$2.63 \times 10^{-10}$	$2.62 \times 10^{-10}$	$1.40 \times 10^{-11}$	$9.77 \times 10^{-6}$	$9.77 \times 10^{-6}$
79	$6.56 \times 10^{-10}$	$7.45 \times 10^{-12}$	$2.57 \times 10^{-12}$	$1.06 \times 10^{-10}$	$1.05 \times 10^{-10}$	0	0	$8.76 \times 10^{-10}$
80	$1.97 \times 10^{-10}$	$1.19 \times 10^{-11}$	$2.92 \times 10^{-12}$	$6.41 \times 10^{-11}$	$4.46 \times 10^{-12}$	$1.80 \times 10^{-11}$	$1.26 \times 10^{-5}$	$1.26 \times 10^{-5}$

Table 6. Continued.

81	$6.28 \times 10^{-10}$	0	$1.95 \times 10^{-12}$	$1.06 \times 10^{-10}$	$1.48 \times 10^{-10}$	0	0	$8.84 \times 10^{-10}$
82	$5.60 \times 10^{-10}$	$1.05 \times 10^{-10}$	$9.24 \times 10^{-12}$	0	$5.92 \times 10^{-11}$	0	0	$7.34 \times 10^{-10}$
83	$1.94 \times 10^{-10}$	$3.79 \times 10^{-11}$	$6.64 \times 10^{-11}$	0	$5.81 \times 10^{-10}$	$2.92 \times 10^{-11}$	$2.04 \times 10^{-5}$	$2.04 \times 10^{-5}$
84	$2.28 \times 10^{-10}$	$8.53 \times 10^{-12}$	$1.49 \times 10^{-11}$	0	$6.99 \times 10^{-10}$	$3.37 \times 10^{-11}$	$2.36 \times 10^{-5}$	$2.36 \times 10^{-5}$
85	$1.80 \times 10^{-9}$	$3.10 \times 10^{-11}$	$1.33 \times 10^{-11}$	0	0	$2.02 \times 10^{-11}$	$1.41 \times 10^{-5}$	$1.41 \times 10^{-5}$
86	$5.54 \times 10^{-10}$	$6.02 \times 10^{-11}$	$2.76 \times 10^{-10}$	$7.00 \times 10^{-10}$	$4.85 \times 10^{-11}$	0	0	$1.64 \times 10^{-9}$
87	$1.89 \times 10^{-10}$	0	$3.22 \times 10^{-12}$	0	$3.29 \times 10^{-11}$	0	0	$2.25 \times 10^{-10}$
Mean	$1.06 \times 10^{-9}$	$7.28 \times 10^{-11}$	$3.91 \times 10^{-11}$	$1.04 \times 10^{-10}$	$1.77 \times 10^{-10}$	$3.50 \times 10^{-11}$	$2.42 \times 10^{-5}$	$2.42 \times 10^{-5}$




Fig. 3. Non-carcinogenic health risk ratios.

pregnancy-induced iron deficiency anemia [70], and deterioration of child health [71].

#### Total Health Risk Assessment

The total health risk values at 54 of the sites between  $5.47 \times 10^{-6}$  and  $7.18 \times 10^{-4}$ , whereas the remaining 33 sites had negligible values in relation to total health risk (Table 6). We found that As was the primary pollutant of concern, measuring as 99.99 % of the total health risk.

#### Water Protection Countermeasures

In total, 70% of the sample points in the study area exceeded the Mn standards for centralized domestic and drinking water sources and industrial and agricultural water use. Mn removal equipment or chemicals must be used in groundwater. Additionally, 8% and 3% of the sample points exceed the standards for Fe and As, respectively. Water treatment facilities could be added to wells or the use of wells could be suspended. Where As content meets the national standards of groundwater to be suitable for use as a centralized domestic and drinking water source; thus, the number of As treatment facilities does not need to be increased. However, at 62% of the sample points, where the health risk was high, monitoring needs to be strengthened and the passage of As into groundwater needs to be obstructed.

## Conclusions

The results of groundwater quality study showed that 70% of the sampling points exceeded the minimum safe levels of Mn and, in 8% and 7% of the sampling points, respectively, Fe and As contents exceeded the limits. Heavy metals are evenly distributed in the groundwater source, there was no regional contamination. There is a significant correlation between Mn and Fe, and a significant correlation between Cu and Pb, however, the correlation of the other two heavy metals is not obvious. Groundwater quality was good when using the comprehensive evaluation method. The human health risk study showed that 62% of the sampling sites had carcinogenic risk values exceeding the maximum acceptable level recommended by various institutions, with As being the major carcinogenic factor in the study area. Non-carcinogenic health risk values were below the negligible standard level, and elements were ranked according to their health risk value as  $Mn > Fe > Pb > Cu > As > Zn$ , of which Mn accounted for 71%. The carcinogenic health risk is almost entirely caused by the presence of As (99.99%). The focus of future research efforts should be directed towards reducing the content of Mn and monitoring the concentration of As in the groundwater source of Suzhou. The results of this study will assist environmental researchers to increase awareness, business owners and homeowners to receive safe water at a low cost; however, appropriate water safety strategies at the household level need to be created to enhancing water Supply safety. Further research is needed to acquire a better understanding of the individual health risks associated with heavy metals in drinking water sources.

## Acknowledgments

This work was supported by the Belt and Road Special Foundation of the State Key Laboratory of Hydrology-Water Resources and Hydraulic Engineering (No. 2018nkms06), and the National Natural Science Foundation of China Grants (Young Scientists Fund) (No. 51509064 and 51309071).

### Conflict of Interest

The authors declare no conflicts of interest.

### References

- ZHANIBEKOV A., ISSAYEVA R., GOLOVATYI S., TASPOLTAYEVA A., AITIMBETOVA A., NURTAYEVA A., KURGANBEKOV Z., TULBASIYEVA A. Assessment of Soil Contamination by Heavy Metals: A Case of Turkistan Region. *Polish Journal of Environmental Studies*. **31** (2), 1985, **2022**.
- SUN Q., SUN Z., XING W., HAO G., LI X., DU J., LI X., DU J., LI C., TIAN H., LI X. Ecological Health Risk Assessment of Heavy Metals in Farmland Soil of Changchun New Area. *Polish Journal of Environmental Studies*. **30** (6), 5775, **2021**.
- SEGHIER T.B., BOUHADJERA K. Pollution Assessment of Heavy Metals in Roadside Agricultural Soils. *Polish Journal of Environmental Studies*. **29** (4), 2855, **2020**.
- SHU W., WANG P., ZHAO J., YU X., XU Q. Characteristics, Sources and Risk Assessment of Heavy Metals in the Ganjiang River Basin, China. *Polish Journal of Environmental Studies*. **29** (2), 1849, **2020**.
- HAN Z., WAN D., TIAN H., HE W., WANG Z., LIU Q. Pollution Assessment of Heavy Metals in Soils and Plants around a Molybdenum Mine in Central China. *Polish Journal of Environmental Studies*. **28** (1), 123, **2019**.
- NKPAA K.W., AMADI B.A., WEGWU M.O. Hazardous metals levels in groundwater from Gokana, Rivers State, Nigeria: Non-cancer and cancer health risk assessment. *Human and Ecological Risk Assessment: An International Journal*. **24** (1), 214, **2018**.
- MITCHELL E., FRISBIT S., SARKAR B. Exposure to multiple metals from groundwater – a global crisis: Geology, climate change, health effects, testing, and mitigation. *Metallomics*. **3** (9), 874, **2011**.
- KRICKOVIC E., NOVKOVIC I., LUKIC T., KRICKOVIC Z. Using DPSEEA Framework to Investigate the Relation between Arsenic Concentration Values in Waters and Cancer Rates in AP Vojvodina (North Serbia). *Polish Journal of Environmental Studies*. **31** (3), 2117, **2022**.
- COOMAR P., MUKHERJEE A., BHATTACHARYA P., BUNDSCHUH J., VERMA S., FRYAR A.E., RAMOS O.E.R., MUNOZ M.O., GUPTA S., MAHANTA C., QUINO I., THUNVIK R. Contrasting controls on hydrogeochemistry of arsenic-enriched groundwater in the homologous tectonic settings of Andean and Himalayan basin aquifers, Latin America and South Asia. *Sci Total Environ*. **689**, 1370, **2019**.
- JOHANNESSON K.H., YANG N., TRAHAN A.S., TELFEYAN K., MOHAJERIN T.J., ADEBAYO S.B., AKINTOMIDE O.A., CHEVIS D.A., DATTA S., WHITE C.D. Biogeochemical and reactive transport modeling of arsenic in groundwaters from the Mississippi River delta plain: An analog for the As-affected aquifers of South and Southeast Asia. *Geochim Cosmochim Acta*. **264**, 245, **2019**.
- MUSHTAQ N., MASOOD N., KHATTAK J.A., HUSSAIN I., KHAN Q., FAROOQI A. Health risk assessment and source identification of groundwater arsenic contamination using agglomerative hierarchical cluster analysis in selected sites from upper Eastern parts of Punjab province, Pakistan. *Human and Ecological Risk Assessment: An International Journal*. **27** (4), 999, **2021**.
- WAQAS H., SHAN A., KHAN Y.G., NAWAZ R., RIZWAN M., SAIF-UR-REHMAN M., SHAKOOR M.B., AHMED W., JABEEN M. Human health risk assessment of arsenic in groundwater aquifers of Lahore, Pakistan. *Human and Ecological Risk Assessment: An International Journal*. **23** (4), 836, **2017**.
- SUSHANT K.S., ASHOK K.G. Health Risk Assessment Due to Groundwater Arsenic Contamination: Children Are at High Risk. *Human and Ecological Risk Assessment: An International Journal*. **18** (4), 751, **2012**.
- KIM T.H., SEO J.W., HONG Y.S., SONG K.H. Case-control study of chronic low-level exposure of inorganic arsenic species and non-melanoma skin cancer. *Journal of Dermatology*. **44** (12), 1374, **2017**.
- MURKO M., ELEK B., STYBL M., THOMAS D.J., FRANCECONI K.A. Dose and Diet -Sources of Arsenic Intake in Mouse in Utero Exposure Scenarios. *Chemical Research in Toxicology*. **31** (2), 156, **2018**.
- USEPA (US Environmental Protection Agency) Baseline Human Health Risk assessment, Vasquez Boulevard and I-70 Superfund Site, U.S. Public Health Service, Denver, CO, **2001**.
- WHO. Guidelines for Drinking-Water Quality, first addendum to third edition. Vol. 1: Recommendations Available at: [http://www.who.int/water\\_sanitation\\_health/dwq/gdwq0506.pdf](http://www.who.int/water_sanitation_health/dwq/gdwq0506.pdf). **2006**.
- AGUSTINA U., HERMAWATI E. Environmental health risk assessment of manganese and cadmium in the ground water around Cipayang Landfill in Depok, Indonesia. 6th International Conference on Water Resource and Environment; **612**, Japan, **2020**.
- HUILI Q., HERONG G. Heavy metals contamination in shallow groundwater of a coal-mining district and a probabilistic assessment of its human health risk. *Human and Ecological Risk Assessment: An International Journal*. **25** (3), 548, **2019**.
- LI R., ZHANG Y., CHU W., DENG Z., CHEN Z., TIAN D. Natural Mineral for Remediation of Iron and Manganese-Contaminated Groundwaters. *Polish Journal of Environmental Studies*. **30** (3), 2161, **2021**.
- JAREDA G., MAHAPATRA S. P., DHEKNE P. Y. Water quality index, heavy metal pollution index and seasonal variation correlation of groundwater of Bailadila iron ore mine area and its peripherals: Dantewada district, Chhattisgarh, India. *Desalination and Water Treatment*. **101**, 7, **2018**.
- SRILERT C., POKKATE W., WATTASIT S., MARIJJA B., MARK R. Non-Carcinogenic Hazard Maps of Heavy Metal Contamination in Shallow Groundwater for Adult and Aging Populations at an Agricultural Area in Northeastern Thailand. *Human and Ecological Risk Assessment: An International Journal*. **20** (3), 689, **2014**.
- SAMUEL O.A., THEOPHILUS T.I., PRAISE GOD E.C., WINIFRED A.U. Computation of human health risk in surface water in Ado-Odo Ota, Ogun State, Nigeria. *Data in brief*. **19**, 1574, **2018**.
- PINTO M.M.S.C., ORDENS C.M., DE MELO M.T.C. INACIO M., ALMEIDA A., PINTO E., DA SILVA E.A.F. An Inter-disciplinary Approach to Evaluate Human Health Risks Due to Long-Term Exposure to Contaminated Groundwater Near a Chemical Complex. *Exposure and Health* **12** (2), 199, **2020**.
- WONGSANIT J., TEARTISUP P., KERDSUEB P., THARPOOPHASIAM P., WORKHUNPISET S.

- Contamination of nitrate in groundwater and its potential human health: a case study of lower Mae Klong river basin. Thailand. *Environ Sci Pollut Res.* **22** (15), 11504, **2015**.
26. WEI F., JINLONG Z., YINZHU Z., YANYAN Z., YUNFEI C., YING S. Water quality and health risk assessment of shallow groundwater in the southern margin of the Tarim Basin in Xinjiang, P. R. China. *Human and Ecological Risk Assessment: An International Journal.* **27** (20), 483, **2021**.
  27. RUI Z., LI M., BIN W., JINSHENG W., FEI D., XUERU G., SHUHE J., YANGUO T. Pollution risk assessment based on source apportionment in a groundwater resource area, NE China. *Human and Ecological Risk Assessment: An International Journal.* **24** (5), 1197, **2018**.
  28. NKPA K.W., AMADI B.A., WEGWU M.O. Hazardous metals levels in groundwater from Gokana, Rivers State, Nigeria: Non-cancer and cancer health risk assessment. *Human and Ecological Risk Assessment: An International Journal.* **24** (1), 214, **2018**.
  29. MIRZABEYGI M., ABBASIA A., YUNESIAN M., NODEHI R.N., YOUSEFI N., HADI M., MAHVI A.H. Heavy metal contamination and health risk assessment in drinking water of Sistan and Baluchistan, Southeastern Iran. *Human and Ecological Risk Assessment: An International Journal.* **23** (8), 189, **2017**.
  30. NAZ A., CHOWDHURY A., MISHRA B.K., GUPTA S.K. Metal pollution in water environment and the associated human health risk from drinking water: A case study of Sukinda chromite mine, India. *Human and Ecological Risk Assessment: An International Journal.* **22** (7), 1433, **2016**.
  31. YUEZAN T., DAOYING X.. Overdraft evaluation and pumping-limit planning of groundwater in Huaibei Plain of Anhui Pvince. *Geotech. Investig. Surv.* **2006** (1), 30, **2006** [In Chinese].
  32. GUOQIANG Y., JINGYING M., XIAOSI S. A briefly review of evaluation index and method of groundwater overdraft. *Water Saving Irrig.* **2012** (8), 34, **2012** [In Chinese].
  33. CHINA MINISTRY OF ENVIRONMENTAL PROTECTION. Water Quality-Determination of 65 Elements-Inductively Coupled Plasma-Mass Spectrometry. **2014**. Available online: <https://www.mee.gov.cn/>.
  34. SAC (Standardization Administration of the People's Republic Of China). Standard for groundwater quality, China. **2017**. Available at:<http://www.mwr.gov.cn/>
  35. LIJUAN W., YUEZAN T., BIN S., LIJUN W., PEIGUI L. Environmental and Health Risks Posed by Heavy Metal Contamination of Groundwater in the Sunan Coal Mine, China. *Toxics.* **10** (7), 390, **2022**.
  36. ZIVKOVI N., TAKIC L., DJORDJEVIC L., DJORDJEVIC A., MLADENOVIC-RANISAVLJEVIC I., GOLUBOVIC T., BOZILOV A. Concentrations of Heavy Metal Cations and a Health Risk Assessment of Sediments and River Surface Water: A Case Study from a Serbian Mine. *Polish Journal of Environmental Studies.* **28** (3), 2009, **2019**.
  37. IRIS (Integrated Risk Information System). US Environmental Protection Agency. Cincinnati, OH, USA. Available at: <http://www.epa.gov/iris>. **2005**.
  38. HABIB M.E., ALI M.A., FATTAH K.P. Modified sand for theremoval of manganese and arsenic from groundwater. *Journal of Environmental Engineering and Science.* **15** (4), 189, **2021**.
  39. BONDU R., CLOUTIER V., ROSA E. Occurrence of geogenic contaminants in private wells from a crystalline bedrock aquifer in western Quebec, Canada. *Geochemical sources and health risks. Journal of Hydrology.* **559**, 627, **2018**.
  40. USMAN U.A., YUSOFF I., RAOOV M., ALIAS Y., HODGKINSON J., ABDULLAH N., HUSSIN N.H. Natural sources of iron and manganese in groundwater of the lower Kelantan River Basin, North-eastern coast of Peninsula Malaysia: water quality assessment and an adsorption-based method for remediation. *Environmental Earth Sciences.* **80**, 12, **2022**.
  41. KHOZYEM H., HAMDAN A., TANTAWY A.A., EMAM A., ELBADRY E. Distribution and origin of iron and manganese in groundwater: case study, Balat-Teneida area, El-Dakhla Basin, Egypt. *Arabian Journal of Geosciences.* **12**, 6, **2019**.
  42. RIVERA-RODRIGUEZ D.A., BELTRAN-HERNANDEZ R.I., LUCHOCONSTANTINO C.A., CORONEL-OLIVARES C., HERNANDEZ-GONZALEZ S., VILLANUEVA-IBANEZ M., NOLASCO-ARIZMENDI V., VAZQUEZ-RODRIGUEZ G.A. Water quality indices for groundwater impacted by geogenic background and anthropogenic pollution: case study in Hidalgo, Mexico. *International Journal of International Science and technology.* **16** (5), 2201, **2019**.
  43. ZHAO W.D., ZHAO L., GONG J.S., ZHOU W., QIAN J. Pollution assessment and source apportionment of shallow groundwater in Suzhou mining area, China. *Earth Science Frontiers.* **28** (5), 1, **2021** [In Chinese].
  44. HUANG Z.N., TANG Y.K., ZHANG K.X., CHEN Y.H., WANG Y., KONG L.M., YOU T.Y., GU Z.Q. Environmental risk assessment of manganese and its associated heavy metals in a stream impacted by manganese mining in South China. *Human and Ecological Risk Assessment: An International Journal.* **22** (6), 1341, **2016**.
  45. YI F.H., CHEN L., YAN F. The health risk weighting model in groundwater quality evaluation. *Human and Ecological Risk Assessment: An International Journal* **25** (8), 2089, **2019**.
  46. ZUO R., MENG L., WANG B., WANG J.S., DING F., GUO X.R., JIN S.H., TENG Y.G. Pollution risk assessment based on source apportionment in a groundwater resource area, NE China. *Human and Ecological Risk Assessment: An International Journal* **24** (5), 1197, **2018**.
  47. BROWN C.J., BARLOW J.R.B., CRAVOTTA C.A., LINDSEY B.D. Factors affecting the occurrence of lead and manganese in untreated drinking water from Atlantic and Gulf Coastal Plain aquifers, eastern United States-Dissolved oxygen and pH framework for evaluating risk of elevated concentrations. *applied Geochemistry.* **101**, 88, **2019**.
  48. ZHENGANG X., YI D., HUIMIN H., LIANG W., YUNLIN Z., GUIYAN Y. Biosorption Characteristics of Mn (II) by *Bacillus cereus* Strain HM-5 Isolated from Soil Contaminated by Manganese Ore. *Polish Journal of Environmental Studies.* **28** (1), 463, **2019**.
  49. ESMAEILI S., MOGHADDAM A.A., TZIRITIS E. Multivariate statistics and hydrogeochemical modeling for source identification of major elements and heavy metals in the groundwater of Qareh-Ziaeddin plain, NW Iran. *Arabian Journal of Geosciences.* **11** (1), **2018**.
  50. EGBI C.D., ANORNU G.K., APPIAH-ADJEI E.K., GANYAGLO S.Y., DAMPARE S.B. Trace Metals Migration and Contamination Assessment of Groundwater in the Lower Volta River Basin, Ghana. *Exposure and Health.* **13** (3), 487, **2021**.
  51. LIU X.H., YE Q.H., CHEN K., PENG W.H. Characterization Haracterization and control mechanism

- of main ions heavy metals and multi-isotopes in groundwater of coal-bearing aquifer a case study in Qianyingzi Coal Mine. Characteristics and control mechanism of main ions, heavy metals and multi-isotopes in groundwater of coal-bearing aquifers: A case study of Qianyingzi Coal Mine. *Fresenius Environment Bulletin*. **31** (1A), 1407, **2022**.
52. MAJOUNI S., KIM J.S., SWEENEY E., KELTIE E., ABIDI S.S.R. Applying Machine Learning to Arsenic Species and Metallomics Profiles of Toenails to Evaluate Associations of Environmental Arsenic with Incident Cancer Cases. *Studies in health technology and informatics*. **294**, 3, **2022**.
  53. SAWADA N. Association between Arsenic Intake and Cancer-From the Viewpoint of Epidemiological Study. *Nihon eiseigaku zasshi. Japanese journal of hygiene*. **72** (3), 265, **2018**.
  54. KRICKOVIC E., NOVKOVIC I., LUKIC T., KRICKOVIC Z. Using DPSEEA Framework to Investigate the Relation between Arsenic Concentration Values in Waters and Cancer Rates in AP Vojvodina (North Serbia). *Polish Journal of Environmental Studies*. **31** (3), 2117, **2022**.
  55. GUO Y.J., WANG Y., LIU Y.G., HOU L., YANG G.Y., LI M.Y., WEN G.J. Effects of Environmental Factors on Arsenic Fractions in Plateau Lakeside Wetland Sediments. *Polish Journal of Environmental Studies*. **27** (5), 202, **2018**.
  56. DIZAJI A.R., HOSSEINI S.A., REZAVERDINEJAD V., SHARAFATI A. Assessing Pollution Risk in Ardabil Aquifer Groundwater of Iran with Arsenic and Nitrate Using the SINTACS Model. *Polish Journal of Environmental Studies*. **29** (4), 2609, **2020**.
  57. SINGH R., GAYEN A., KUMAR S., DEWANGAN R. Geo-spatial distribution of arsenic contamination of groundwater resources in intricate crystalline aquifer system of Central India: Arsenic toxicity manifestation and health risk assessment. *Human and Ecological Risk Assessment: An International Journal*. **27** (6), 1588, **2021**.
  58. SHAJI E., SANTOSH M., SARATH K.V., PRAKASH P., DEEPCHAND V., DIVYA B.V. Arsenic contamination of groundwater: A global synopsis with focus on the Indian Peninsula. *Geoscience Frontiers*. **12** (3), **2021**
  59. ZHANG Y.R., GUI H.R., HUANG Y.H., YU H., LI J., WANG M.C., FANG H.X., JIANG Y.Q., WANG C.L., CHEN C. Characteristics of Soil Heavy Metal Contents and its Source Analysis in Affected Areas of Luning Coal Mine in Huaibei Coalfield. *Polish Journal of Environmental Studies*. **30** (2), 1465, **2021**.
  60. HU Y.H., WANG X.M., DONG Z.B., LIU G.J., WANG M.H., LIU M.C. Groundwater Quality at the Huaibei Coalfield, China. *Analytical Letters*. **48** (10), 1654, **2015**.
  61. LIANG B., YANG Y.J., WANG X.L., QIAN Y., LU Z.I., GUO C., WU J.B., ZHAO S.L. Ecological risk evaluation of heavy metals in soils at the high environmental risk area of the sub-branches of Huaihe River. *Chinese Journal of Applied and Environmental Biology*. **22** (2), 257, **2017**.
  62. LI X.H., HOU G.Q., YUAN C.S., LIU Y., ZHANG L., YU B., CUI L.Q. Research on relationship between the concentration of water fluoride, water As and depth of well in the water-drinking endemic fluorosis areas of Qixian. *Henan. J. Prev. Med.* **21**, 133, **2010**.
  63. HANG L.M., YANG Q.C., WANG H., GU Q.B., ZHANG Y.L. Genetic interpretation and health risk assessment of arsenic in Hetao Plain of inner Mongolia, China. *ENVIRONMENTAL RESEARCH*. **2022**, 208
  64. HUANG K., LIU Y.Y., YANG C., DUAN Y.H., YANG X.F., LIU C.X. Identification of Hydrobiogeochemical Processes Controlling Seasonal Variations in Arsenic Concentrations Within a Riverbank Aquifer at Jiangnan Plain, China. *Water Resources Research*. **54** (7), 4294, **2018**.
  65. NGHIEM A.A., STAHL M.O., MAILLOUX B.J., MAI T.T., TRANG P.T., VIET P.H., HARVEY C.F., VAN GEEN A., BOSTICK B.C. Quantifying Riverine Recharge Impacts on Redox Conditions and Arsenic Release in Groundwater Aquifers Along the Red River, Vietnam. *Water Resources Research*. **55** (8), 6712, **2019**.
  66. XIE X., JOHNSON T. M., WANG Y., LUNDSTROM C.C., ELLIS A., WANG X., DUAN M. Mobilization of arsenic in aquifers from the Datong Basin, China: evidence from geochemical and iron isotopic data. *Chemosphere*. **90** (6), 1878, **2012**.
  67. HAGIWARA H., AKAI J., TERASAKI K., YOSHIMURA T., LUO H. Black colored sandy sediments caused by bacterial action, and the mechanism for arsenic enrichment of groundwater in Inner Mongolia. *Applied Geochemistry*. **26**, 380, **2011**.
  68. MAITY J.P., NATH B., KAR S., CHEN C.Y., BANERJEE S., JEAN J.S., LIU M.Y., CENTENO J.A., BHATTACHARYA P., CHANG C.L., SANTRA S.C. Arsenic induced health crisis in peri-urban Moyna and Ardebok villages, West Bengal, India: an exposure assessment study. *Environ. Geochem. Health*. **34**, 563, **2012**.
  69. CHEN T.T., SU Y.H., YUAN X.H. Influences of the petroleum-recovering activity on the arsenic level in groundwater of Kuitun, Xinjiang, China. *Human and Ecological Risk Assessment*. **24** (8), 2195, **2018**.
  70. HAFEMAN D., FACTOR-LITVAK P., CHENG Z.Q., VAN GEEN A., AHSAN H. Association between manganese exposure through drinking water and infant mortality in Bangladesh. *Environ. Health Perspect*. **115**, 1107, **2007**.
  71. AHMDE J., WONG LP., CHUA Y.P., CHANNA N., MEMON U.U.R., GARN J.V., YASMIN A., VANDERLICE J.A. Heavy metals drinking water contamination and health risk assessment among primary school children of Pakistan. *Journal of Environmental Science and Health Part A-Toxic/Hazardous Substances & Environmental Engineering*. **56** (6), 667, **2021**.