















Table 8. Nitrogen, phosphorus, potassium, and protein concentrations in milled rice grains as affected by the interaction between rice variety and nitrogen level in 2020 and 2021.

| Treatments |          | N %    |        | Pv%   |       | K%    |       | Protein |        |
|------------|----------|--------|--------|-------|-------|-------|-------|---------|--------|
| Variety    | Kg N /ha | 2019   | 2020   | 2019  | 2020  | 2019  | 2020  | 2019    | 2020   |
| Giza 179   | 0        | 1.24g  | 1.13e  | 0.21k | 0.20k | 0.13l | 0.13l | 7.13g   | 6.47j  |
|            | 50       | 1.46de | 1.33d  | 0.27i | 0.26i | 0.18j | 0.17i | 8.4de   | 7.64g  |
|            | 100      | 1.54cd | 1.41c  | 0.29h | 0.28g | 0.2h  | 0.2h  | 8.86c   | 8.13e  |
|            | 150      | 1.65b  | 1.51b  | 0.3g  | 0.31e | 0.22f | 0.22f | 9.46b   | 8.71c  |
|            | 200      | 1.79a  | 1.65a  | 0.33f | 0.33d | 0.23e | 0.23d | 10.3a   | 9.51a  |
| Sakha 109  | 0        | 1.34f  | 1.20e  | 0.23j | 0.21j | 0.14l | 0.13k | 7.73f   | 6.90h  |
|            | 50       | 1.52cd | 1.37cd | 0.30g | 0.30f | 0.20h | 0.17i | 8.75cd  | 7.89f  |
|            | 100      | 1.6bc  | 1.45bc | 0.32f | 0.35c | 0.24d | 0.2g  | 9.19bc  | 8.33d  |
|            | 150      | 1.65b  | 1.50b  | 0.34e | 0.38b | 0.26b | 0.24d | 9.46b   | 8.63c  |
|            | 200      | 1.76a  | 1.63a  | 0.38c | 0.42a | 0.29a | 0.25c | 10.12a  | 9.34b  |
| Giza 182   | 0        | 0.98h  | 0.87f  | 0.21k | 0.21j | 0.15k | 0.14j | 5.65h   | 4.99k  |
|            | 50       | 1.29fg | 1.17e  | 0.31g | 0.27h | 0.19i | 0.19h | 7.43fg  | 6.74i  |
|            | 100      | 1.44e  | 1.31d  | 0.36d | 0.30f | 0.21g | 0.23e | 8.28e   | 7.54g  |
|            | 150      | 1.57bc | 1.43bc | 0.39b | 0.35c | 0.24e | 0.27b | 9.03bc  | 8.24de |
|            | 200      | 1.78a  | 1.64a  | 0.42a | 0.37b | 0.26c | 0.29a | 10.22a  | 9.45ab |

### Conclusions

The studied varieties showed different behavior under the various nitrogen levels. Giza182 (Indica Type) recorded a high grain yield with better nutritional value (protein, carbohydrate, lipids, ash, phosphorus, and potassium contents). Higher grain quality characteristics (milling recovery, gelatinization temperature, kernel elongation, and amylose) were obtained from Sakha 109. Generally, increasing the nitrogen level up to 200 kg N/ha increased rice grain yield, hulling %, milling %, and head rice % as well as carbohydrate %, oil %, ash %, amylose %, nitrogen %, phosphorus %, potassium %, and protein % in milled rice grains.

### Acknowledgments

The authors would like to thank King Saud University, Riyadh, Saudi Arabia for supporting this research through the Researchers Supporting Project number (RSP2024R505).

### Funding

This research was financially supported by the Researchers Supporting Project (number RSP2024R505), King Saud University, Riyadh, Saudi Arabia.

### Conflict of Interest

The authors declare no conflict of interest

### References and Notes

1. ABDEL-LATTEF A.S., ELKHOBAY W.M., ABDELAAL KH., MOHAMED A. Genetic behavior of morphology yield yield components and grain quality in Rice (*Oryza sativa* L). Journal of Plant Production, 5 (11), 1945, 2014.
2. EL-HITY M.A., METWALLY T.F., MOHAMMED A.A., ELSAKA M.A.M., HASSAN F., HAFEZ Y.M., ABDELAAL KH. Integrated fertilizer management on growth and nutrient uptake of Egyptian hybrid rice. Fresenius Environmental Bulletin, 29 (12A), 11588, 2020.
3. HAFEZ Y., ELKHOBAY W., MAZROU Y.S.A., GHAZY M., ELGAMAL A., ABDELAAL KH. Alleviating the detrimental impacts of salt stress on morpho-physiological and yield characters of rice plants (*Oryza sativa* L) using actosol Nano-Zn and Nano-Si. Fresenius Environmental Bulletin, 29 (8), 6882, 2020.
4. OMAR A., ZAYED B., ABDEL SALAM A., HAFEZ Y., ABDELAAL K. Folic acid as foliar application can improve growth and yield characters of rice plants under irrigation with drainage water. Fresenius Environmental Bulletin, 29 (10), 9420, 2020.
5. MOHAMED A.A., ZAYED B., HAFEZ E., ELTOBGY R., HAFEZ Y., ABDELAAL KH. Effect of Potassium Fertilizer Treatments and Harvest Date on Rice Grain Yield and Quality of Rice Plants (Giza 179 Variety). Fresenius Environmental Bulletin, 30 (2), 1010, 2021.
6. MOHAMED A.A., MAZROU Y., ELHAG D., SAAD A., HAMDY R., ABDELAAL K.H. Response of some faba bean varieties to amino acids and potassium foliar application under delta soil conditions. Fresenius Environmental Bulletin, 30 (11), 11816, 2021.
7. ABDELAAL K.H., MAZROU Y., MOHAMED A.A., GHAZY M., BARAKAT M., HAFEZ Y., GABALLAH M. The different responses of rice genotypes to heat stress associated with morphological chlorophyll and yield characteristics Notulae Botanicae Horti Agrobotanici Cluj-Napoca, 49 (4), 12550, 2021.
8. ELHITY M.A., OMER A.M., ZAYED B.A., ASSRA A.A., HASSAN M.M., HAFEZ Y.M., ABDELAAL KH.

