

Original Research

Spatial Distribution Characteristics and Source Analysis of Ions in Lancang River and Yangtze River Source Region During Wet Period

Wei Deng^{1,2}, Min Liu^{1,2}, Cheng Han^{1,2}, Liangyuan Zhao^{1,2,3*}, Yuting Zhang^{1,2}, Yuan Hu^{1,2}, Mingli Wu^{1,2}, Luyi Nan¹

¹ Basin Water Environmental Research Department, Yangtze River Scientific Research Institute, Wuhan 430010, China

² Key Lab of Basin Water Resource and Eco-Environmental Science in Hubei Province, Changjiang River Scientific Research Institute, Wuhan 430010, China

³ Innovation Team for Basin Water Environmental Protection and Governance of Changjiang Water Resources Commission, Wuhan, 430010, China

Received: 06 November 2023

Accepted: 30 January 2024

Abstract

The security of the environment in the headwaters is vital to the whole basin. This study focused on the main ion composition, spatial distribution characteristics, and sources of river water and surface sediments in the source regions of the Lancang River and the Yangtze River and analyzed hydrochemical type and control factors. The results showed that the main cations in the water of the source region were Ca^{2+} and Na^+ , with a combined proportion of over 89%. The main anions were Cl^- , SO_4^{2-} , and HCO_3^- , with a combined proportion of over 97%. There were some differences in the content of metallic elements in the water. The average contents of As, Cu, Pb, Cr, Cd, Ni, and Ti in the river water of the source region of the Yangtze River (SRYR) were higher than those in the source region of the Lancang River (SRLR). The main hydrochemical types in the source region were Na-Cl and Ca- SO_4 - HCO_3 , which were controlled by rock weathering and evaporating crystallization. The ions in the river water of the source region were less affected by atmospheric transport and human activities, but more affected by rock weathering. In the SRLR, the water ions were mainly affected by evaporites and carbonate rocks. And in the SRYR, those were mainly affected by evaporites, silicates, and carbonate rocks. The average contents of Ti, Mn, Fe, Sr, and Ba in the surface sediments of rivers were higher than those of other metallic elements. The metallic elements in sediments were controlled by natural and traffic pollution. The study could provide basic data and a scientific basis for the study of the geochemical processes of the water environment and the protection of river ecological environments in plateau rivers.

Keywords: The Source Region of the Yangtze River and the Lancang River, Surface Water, Surface Sediment, Distribution Characteristics of Ions, Hydrochemical Characteristics

* e-mail: zhaoliangyuannew@163.com

Introduction

Due to the influence of geology, climate, and human activities [1, 2], the ion composition and distribution of river water could directly reflect the basic characteristics of the basin, even the history of water formation and migration [3]. After different physical and geochemical processes, the content and distribution of different ions not only reflected the different hydrochemical characteristics, but also reflected the relationship between the source, migration, and transformation of solutes and regional natural conditions [4-6]. Generally, the solutes of surface water are mainly derived from rock weathering and the erosion of soils in the drainage basin; these processes are the major components of the exogenic cycle of elements in the continent-river-ocean system [7, 8]. The Ca^{2+} , Mg^{2+} , SO_4^{2-} , and HCO_3^- in river water can be attributed mainly to rock weathering. The types and influence degree of rock weathering can be qualitatively analyzed by the Piper plot and Gibbs distribution [9]. Quantitative analysis of rock weathering rates requires the analysis of strontium and its isotopic characteristics [10]. Soil erosion in river basins leads to some metal elements in soil entering water, such as Al, Fe, Mn, and Mo, which are abundant in the earth's crust. Some ions still might come from human activities, such as NO_3^- , Pb, and Zn, even though there were relatively few human activities in the area. Sediments are produced by natural erosion and biological processes. Ions that were in river water could enter sediments through physical and chemical actions, including adsorption, sedimentation, and complexation [11]. When the external conditions changed, the ions in the sediment would be released into the water again, which changed the ion concentration and affected the basin characteristics and eco-environment [12-16]. Therefore, it is of great significance to study the river water and sediments together for the evaluation of water environment quality and the protection of the river ecological environment.

The Qinghai-Tibet Plateau is known as the "Asian Water Tower" [17]. It is the birthplace of many major rivers in East Asia, Southeast Asia, and South Asia. Its natural environment and ecosystem hold a special position globally. Due to its unique climate conditions, complex geological structures, and active crustal activity, the region has always been a hot topic of widespread concern and research by scientists. In recent years, with global warming, the accelerated melting of glaciers located on the Qinghai-Tibet Plateau has had an impact on the hydrology and hydrochemistry of the basin [18, 19]. Many studies found that the hydrochemistry of rivers in the Qinghai-Tibet Plateau was mainly controlled by rock weathering [20-22]. Global climate change accelerates the weathering and erosion of rocks and influences the ionic content and distribution characteristics in rivers to some extent [23, 24]. At the same time, the intensification of human activities has also increased the environmental pressure on plateau rivers. It has become one of the main sources of ions [25].

The source region, located on the Qinghai-Tibet Plateau, is the starting point of the hydrologic cycle in the Lancang River and Yangtze River basins. Its water resource status and changes affect the sustainable utilization of basin water resources [26-28]. The geographical location of the SRYR and the SRLR is relatively close. They are both in the Qinghai-Tibet, and the Dangqu in SRYR and the SRLR both originated from glaciers in the Tanggula Mountains. Therefore, there might be some similarities in the physical and chemical properties and the composition of ions in the river between them. However, most studies about the SRYR or the SRLR were analyzed separately or as a whole, and there were few studies in other fields on comparison between them, except the distribution characteristics of phytoplankton and isotopes [29-36]. Especially, there was little comparative analysis between the ion composition, spatial distribution characteristics, and source of river water and sediment. This study focused on the main ion composition, spatial distribution characteristics, and sources of river water and surface sediments in the SRYR or SRLR, hydrochemical type, and control factors, and comparative analysis was conducted. It can provide basic data and a scientific basis for geochemical processes and the protection of the ecological environment in plateau rivers.

Materials and Methods

Study Area

Located in the central region of the Qinghai-Tibet Plateau [16], the source region of the Lancang River and the Yangtze River are not only important ecological security barriers in China, but also sensitive and fragile areas for climate and ecological environment changes [16, 37, 38]. The Lancang River, also known as the Mekong River, is a famous international river. It originated in the northeast of the Tanggula Mountains in Qinghai Province, China. Its main stream in Qinghai Province is 454 km long, and its catchment area is $3.7 \times 10^4 \text{ km}^2$. The source region of the Lancang River has a dense river network. The first-class tributaries include Zanaqu, Ayong, Budangqu, Shaqu, Banyong, Geruoyong, Ningqu, Ziqu, and Angqu [39]. The Yangtze River is considered to have three sources, namely, Dangqu in the South, Chumaer River in the North, with the Tuotuo River being the main source. The Dangqu originated from the eastern part of Tanggula Mountain in the south of the Qinghai-Tibet Plateau. The Chumaer River originated from Hoh Xil Mountain in the south branch of Kunlun Mountain, which is in the north of the Qinghai-Tibet Plateau. And the Tuotuo River originated from Gladandong Ice Peak, which is the main peak of Tanggula Mountain. The basin area of the source region is about $13.82 \times 10^4 \text{ km}^2$. Its river system is fan-shaped. The main tributaries include the Beilu River, Sewuqu, Moqu, Buqu, Keqiaqu, and Nieqiaqu. There are over 1.1×10^4 lakes with a total area of approximately 1027 km^2 [40].

The source region is located in the transition zone of a cold, semi-arid, and semi-humid climate, with typical climate characteristics of an inland plateau [39]. It is dry and cold, with strong solar radiation and a short frost-free period. Precipitation is concentrated in the summer, accounting for 60% to 70% of the annual precipitation [41]. The source region is high in altitude, low in air pressure, and thin in air. Influenced by the westerly circulation in the upper air, the westerly wind prevails. The water quality of the rivers in the source region is good. Due to the influence of sediment and regional geological conditions, there are certain differences in water quality among different rivers [42, 43]. The river sediment in the source region is controlled by the geological background. Some metallic elements are enriched in river sediments from certain regions, such as Ni, Cd, and Pb. In recent years, with the rapid construction of local road transportation and the development of the social economy, human activities have gradually become one of the most important sources of ions in the Qinghai-Tibet Plateau [44].

Data Collection

According to the actual situation, 17 monitoring sections were selected for research. There were 13 points at the SRYSR (JY-01~JY-13), and 4 points at the SRLR (JY-14~JY-17). The location is shown in Figure 1, and the specific information is shown in Table 1. The sampling time was the wet period (July 2022). After the sample collection was completed, it was sealed and packaged before being refrigerated and transported back to the laboratory.

Sample Pretreatment and Detection Method

Surface Water Sample

Surface water samples were collected at a depth of about 10cm and filtered through pre-washed 0.45 μm Millipore nitrocellulose filters. The initial portion of the filtration was discarded to clean the membrane, and the following ones were acidified to $\text{pH} < 2$ with ultra-purified 6 M HNO_3 and then stored in plastic bottles for trace metal analyses. Cleaning of plastic bottles was carried out by soaking in 20% (v/v) HNO_3 for 24 h and then rinsing with milli-Q deionized water ($\sim 18 \text{ M}\Omega/\text{cm}$ resistivity; $\text{TOC} < 5 \text{ mg/L}$) from Milli-Q (Millipore, Direct 8).

Field indicators were determined by a Portable Multi-parameter Water Quality Monitoring Analyzer (EX01, USA). Metallic elements were detected by an Inductively Coupled Plasma Mass Spectrometer (NexION 300X, USA) with a detection limit of 0.06~0.41 $\mu\text{g/L}$ and a recovery rate of 86.8~99.8%. Hg was detected by the Atomic Fluorescence Spectrophotometer (AFS-3100, China) with a detection limit of 0.01 $\mu\text{g/L}$ and a recovery rate of 93.0~104.0%. The contents of cations (K^+ , Na^+ , Ca^{2+} , and Mg^{2+}) were analyzed by a Microwave Plasma Atomic Emission Spectrometer (Agilent 4200, USA) with a detection limit of 1.94~6.61 $\mu\text{g/L}$ and a recovery rate of 95.6~108.7%.

The contents of anions (F^- , Cl^- , SO_4^{2-} , and NO_3^-) were analyzed by Ion Chromatography (ICS900, USA) with a detection limit of 0.006~0.018 mg/L and a recovery rate of 99.1~116.3%. The CO_3^{2-} and HCO_3^- were analyzed by acid-




Fig. 1. Sampling sites in the study area

Table 1. Basic information table of sampling points

Simple Sites			Longitude and Latitude		Altitude (m)		
River system	Name	Number	Longitude	Latitude			
The SRYR	Dangqu	Dangqu Upstream	JY-01	94°32'42"	32°41'08"	4881	
		Dangqu Bridge 1	JY-02	94°12'55"	32°52'14"	4676	
		Dangqu Bridge 2	JY-03	93°50'51"	32°53'03"	4653	
		Yanshiping	JY-04	92°04'26"	33°36'02"	4525	
		Buqu	JY-05	92°21'57"	33°51'43"	4525	
		Gaerqu	JY-06	92°21'58"	33°51'57"	4662	
	Chumaer River		JY-07	93°18'15"	35°18'23"	4479	
	Tuotuo River		JY-08	92°26'37"	34°19'51"	4483	
	Tongtian River	Nangjibalong		JY-09	93°01'00"	34°07'40"	4401
		Keqianqu		JY-10	94°50'40"	34°11'14"	4486
		Nieqiaqu		JY-11	95°49'29"	34°01'11"	4016
		Qumalai		JY-12	95°49'16"	34°01'30"	4024
		Zhimenda		JY-13	97°14'17"	33°01'00"	3494
The SRLR	Zhaqu	Moyun Town	JY-14	94°14'53"	33°09'52"	4497	
		Zhaqu Bridge	JY-15	94°36'44"	33°12'01"	4335	
		Zaduo	JY-16	95°15'26"	32°54'07"	4031	
		Nangqian	JY-17	96°27'08"	32°18'53"	3609	

base titration (methyl orange as an indicator and diluted hydrochloric acid titration) with an error of less than 5%.

Sediment Sample

After being naturally dried, sand and gravel were removed from the sediment. It was ground before passing through a mesh sieve with a pore size of 0.075 mm. According to the literature method [45–47], sediment samples were digested by a Microwave Digestion Instrument (MARS6, USA). Metallic elements in the sediment were detected by an Inductively Coupled Plasma Mass Spectrometer with a detection limit of 0.04–2.0 mg/kg and a recovery rate of 75–120%. Hg in the sediment was detected by an Atomic Fluorescence Spectrophotometer with a detection limit of 0.002 mg/kg and a recovery rate of 80%.

Analytical Method

The hydrochemical characteristics, distribution characteristics, and source of the main ions were analyzed by the Piper plot, Gibbs diagram, and Ion ratio. The possible sources of metallic elements in sediment were analyzed by combining correlation analysis with principal component analysis. The analysis software was SPSS 26. The mapping software was Origin 2021.

Results and Discussion

Distribution Characteristics and Source of Ions in Water

Physical and Chemical Properties Analysis

According to the study results (Table 2), the main rivers in the SRYR were alkaline, and the

temperature difference was large. The pH range was 8.2–8.8. The water temperature range was 7.5–18.6°C. The dissolved oxygen (DO) range was 7.0–9.3 mg/L. The redox potential (ORP) range was 16.0–109 mV. The conductivity range was 333–5326 µS/cm. The total dissolved solids (TDS) range was 145–2875 mg/L.

Similar to the SRYR, the main rivers in the SRLR were alkaline, and the temperature difference was large. The pH range was between 8.3 and 8.6. The water temperature range was 9.7–17.3°C. The DO range was 7.6–8.8 mg/L. The ORP range was 22.3–56.9 mV. The conductivity range was 709–2160 µS/cm. The TDS range was 423–1281 mg/L.

The average value of TDS in the source region was 647.3 mg/L and 702.2 mg/L, both of which were much larger than the world average of 150 mg/L [48]. It might be caused by geotectonic movement [34]. The uplift of the Qinghai-Tibet Plateau is the result of tectonic movement, which caused landform ups and downs, and led to the exposure of fresh rocks to accelerate the weathering process [49, 50]. It had caused an increase in the content of ions, such as K⁺, Na⁺, Ca²⁺, and Mg²⁺, entering the river basin through weathering, and TDS was closely related to these ions. Therefore, the tectonic movement had affected the TDS content, making it higher than the world average. The total hardness (TH) range in the SRYR was 279–693 mg/L, and the average value was 647.3 mg/L. The TH range in the SRLR was 338–812 mg/L, and the average value was 620.2 mg/L. It was much larger than the average value of the Yangtze River and the Lancang River basins (291.6 mg/L and 164.4 mg/L), which indicated that the water hardness in the source area was relatively high [51–53].

Most of the river water in the source region was fresh water, with medium or high hardness, and only a small

Table 2. Physical and chemical properties in river water

River system	Name	Number	Temperature (°C)	DO (mg/L)	pH	ORP (mv)	Conductivity (μS/cm)	TDS (mg/L)	
The SRYR	Dangqu Upstream	JY-01	/	/	/	/	/	333	
	Dangqu	Dangqu Bridge 1	JY-02	17.82	7.24	8.63	19.4	539.4	315
		Dangqu Bridge 2	JY-03	18.55	7.72	8.52	59.4	489.5	356
		Yanshiping	JY-04	10.38	8.45	8.35	21.1	340.9	298
		Buqu	JY-05	8.14	8.87	8.31	53.1	407.4	335
		Gaerqu	JY-06	9.02	8.60	8.23	72.5	333.7	145
		Chumaer River	JY-07	16.31	7.52	8.47	37.7	5326	2875
	Tuotuo River	JY-08	17.90	7.06	8.35	16.0	1557	836	
	Tongtian River	Nangjibalong	JY-09	15.80	7.57	8.53	27.8	1461	782
		Keqianqu	JY-10	7.54	9.32	8.27	79.8	442.3	244
		Nieqiaqu	JY-11	12.04	8.68	8.42	109.2	445.8	251
		Qumalai	JY-12	14.13	8.18	8.24	58.5	1386	775
		Zhimenda	JY-13	14.41	9.03	8.75	87.7	1531	871
The SRLR	Zhaqu	Moyun Town	JY-14	17.31	7.61	8.38	22.3	2160	1281
		Zhaqu Bridge	JY-15	9.77	8.83	8.39	56.9	709.2	423
		Zaduo	JY-16	15.03	7.98	8.56	46.2	818.6	548
		Nangqian	JY-17	13.50	8.73	8.61	22.6	815.1	557




Fig. 2. TDS and TH distribution of river water in the source region

fraction was brackish water with high hardness (Figure 2). This might be caused by complex topography and a changeable local climate [54]. It was reported that the Qinghai Provincial Government implemented an

emergency drainage project in 2019. The high salinity lake water located in the upper reaches was guided into the Chumaer River, which led to an increase in salinity in the Chumaer River [55, 56].

Table 3. Anions and cations content of the river water in source region (unit: mg/L)

Water system		Cations				Anions			
		K ⁺	Na ⁺	Ca ²⁺	Mg ²⁺	Cl ⁻	SO ₄ ²⁻	CO ₃ ²⁻	HCO ₃ ⁻
The SRYR	Dangqu	2.34~ 9.68	12.1~ 26.9	124~ 251.5	11.6~ 15.6	5.39~ 20.2	40.8~ 123.9	2.68~ 7.78	85.2~ 182.5
	Chumaer River	12.2	893.9	160.2	66.9	1116.7	358.6	23.6	97.8
	Tuotuo River	16.0	233.8	169.7	23.0	339.8	118.2	10.9	116.4
	Tongtian River	1.11~ 10.6	12.8~ 211.5	61.0~ 179.4	16.1~ 31.0	3.71~ 342.9	56.2~ 136.7	4.86~ 17.3	92.2~ 139.8
The SRLR		2.83~ 6.62	43.6~ 262.0	92.7~ 252.2	26.1~ 44.2	62.2~ 358.7	123~355	2.43~ 7.05	84.8~ 172.9




Fig. 3. Proportion of anions and cations content (a. Cation Percentage; b. Anion Percentage)

Characteristics of Ionic Content and Spatial Distribution in Water

(1) Content and Distribution Characteristics of Main Ions

K^+ , Na^+ , Ca^{2+} , Mg^{2+} , Cl^- , SO_4^{2-} , CO_3^{2-} , and HCO_3^- were the main anions and cations in the river water of the source region (Table 3). The ionic content varied among different regions. The average content of the main cations in the river water of the source region was the same, which was $Ca^{2+} > Na^+ > Mg^{2+} > K^+$. However, the average content of anions was different. The SRYR was $Cl^- > SO_4^{2-} > HCO_3^- > CO_3^{2-}$, and that of the Lancang River was $SO_4^{2-} > Cl^- > HCO_3^- > CO_3^{2-}$.

The content difference and spatial distribution characteristics were clarified by calculating the proportion of ions (Figure 3). In the source region, the main cations in river water were Ca^{2+} and Na^+ , with a combined proportion of over 89%. The main anions were Cl^- , SO_4^{2-} , and HCO_3^- , with a combined proportion of over 97%.

In the SRYR, the main ions in Dangqu were Ca^{2+} (81.9%), SO_4^{2-} (40.5%), and HCO_3^- (51.3%). In the Chumaer River, they were Na^+ (78.9%) and Cl^- (69.9%). In the Tuotuo River, they were Ca^{2+} (38.3%), Na^+ (52.8%), and Cl^- (58.1%). In the Tongtian River, they were Na^+ (42.2%), Ca^{2+} (47.7%), and Cl^- (47.7%). The main ions, which were in the river water of the SRLR, were similar to those of the Tongtian River, but the proportions of Ca^{2+} (58.9%) and SO_4^{2-} (44.3%) were higher. It might be due to the different sources of ions leading to differences in ion content, especially the processes of rock weathering and evaporation-crystallization, which had a great influence on the content of anion and cation in water [3, 57].

(2) Content and Distribution Characteristics of Metallic Elements

From Figure 4, it was found that in the SRYR, the order of the average contents of metallic elements in river water was $Sr > Ba > Ti > Zn > Pb > As > Cu > Cr > Ni > Se > Sb > Cd$, and in the SRLR, it was $Sr > Ba > Zn > As > Cu > Se > Ni > Cr > Sb > Ti > Pb > Cd$. Combined with the average contents of Sr, Ba, and Zn in the source region of the Yellow River, it was found that the average contents of these three metallic elements, which were in the Sanjiangyuan Region, were obviously higher than in other regions. This was due to the Sanjiangyuan Region being located in the Qinghai-Tibet Plateau and influenced by glacial meltwater, rock weathering, and the geological background of the Qinghai-Tibet Plateau [58-60].

In the SRYR, the average content of Zn, Se, Ba, Sr, or Sb in river water was flat with the SRLR, while As, Cu, Pb, Cr, Cd, Ni, or Ti were higher (Figure 4). In particular, the average contents of As, Cu, Pb, Cr, and Ti are several times or even dozens of times higher. This might be due to the fact that there were three sources in the SRYR, but only Dangqu originated from glaciers in the Tanggula Mountain, the same as the SRLR [30]. The Tuotuo River and Chumaer River were different from the SRLR in

origins and geographical locations [61, 62], resulting in the average contents of these metallic elements in the SRYR being different from those in the SRLR. According to Figure 5, it could be found that the ratio of average contents of As, Cu, Pb, Cr, and Ti in the Tuotuo River and Chumaer River were relatively high in the SRYR, which was the reason why these metallic elements in the SRYR were higher than those in the SRLR.

Compared with the source region of the Yellow River [63, 64], the average contents of Cd, Se, Ni, or Sr in the SRYR were flat, while those of As, Zn, Ba, or Sb were lower and Cu, Pb, Cr, or Ti were higher. The average contents of Cu, Cr, Cd, Se, Ba, or Sr in the SRLR were flat with those in the source region of the Yellow River, while As, Zn, Pb, Ni, Ti, or Sb were lower. Except for Sb, the average contents of other metallic elements in river water of the SRYR were higher than the average values of the Yangtze River basin and the World Rivers [48, 65].

From Table 4. [66, 67], it was found that the historical background values of elements in the source region, such as Ni, Cd, Cr, Pb, Zn, Cu, and As, were higher than those in the Yangtze River basin and the World Rivers, which showed that the content difference of these elements was influenced by the background value. The source region was rich in mineral resources. Copper mine reserves account for about 40% of the total reserves in China, and lead ore and zinc ore reserves account for about 30% [68]. Under the natural effects of rainfall, leaching, and rock weathering, these metal elements entered the river, which made the contents in the source region higher than those in other areas. In addition, human activities such as railway and road construction, tourism development, and the increase of animal husbandry in the source region also lead to an increase in Pb, Ti, and Se contents. To sum up, the different background values of rivers, as well as the influence of natural factors and human activities, might lead to the differential distribution of metallic elements in the source region [69-71].

There were differences in the spatial distribution of metallic elements in different rivers in the SRYR. The average contents of Hg, Cr, Cd, Se, Ni, Sb, Co, V, Mo, or Be in different rivers were flat, while the average contents of other ions varied. The average contents of Cu, Zn, Pb, Ba, Ti, or Mn in the Dangqu River were higher than those in the Tuotuo River, Chumaer River, or Tongtian River. The Sr, Li, Tl, or Fe in the Chumaer River were higher than those in other rivers. Except for Se, Sr, Sb, Mo, Tl, and Fe, the average contents of other metallic elements in the Tuotuo River were higher than those in other rivers. In the Tongtian River, only Hg was higher than that in other rivers.

Analysis of Hydrochemical Characteristics and Main Control Factors

The Piper plot (Figure 6) provides a way to visualize the compositions and water types of rivers and therefore, can show the relative importance of different weathering regimes [29]. On the cation diagram, samples from Zhaqu and Dangqu lie in the lower-left corner, where




Fig. 4. Metallic elements' average content of river water in different regions

Table 4. The background value of metallic elements in the river (unit: µg/L)

Rivers	As	Cu	Zn	Pb	Cr	Cd	Ni
The SRYR	2~38	0.73~23.1	0.99~66.4	1.12~57.1	0.26~48.8	0.04~0.5	0.33~6.1
The Yangtze River Basin	1.89	1.17	0.98	0.15	0.21	0.07	1.55
The World Rivers	0.6~13	1~6	4~10	0.2~6	0.08~2.1	0.01~0.2	/




Fig. 5. Distribution of metallic elements in different rivers in the SRYR

the Ca values are high (more than 50%), revealing that the dominant cation in these samples is calcium. On the contrary, samples from the Chumaer River and Tuotuo River fall on the lower-right corner, where the dominant ion is sodium, showing the characteristics of evaporites. The values of samples from the Tongtian River are relatively scattered, but still show that the dominant ions are Ca and Na. On the anion diagram, the samples from Dangqu and a part of the Tongtian River fall on the left, showing carbonate rock weathering features. However, samples from the Chumaer River, the Tuotuo River, and a part of the Tongtian River fall on the lower-right corner, revealing chloride-dominated features. The values of samples from Zhaqu are relatively scattered. According to the Piper plot (Figure 6), the main hydrochemical types in the source region are Na-Cl and Ca-SO₄•HCO₃, but they are diverse in different rivers in the SRYSR. The hydrochemical type of the river water in Dangqu is Ca-SO₄•HCO₃, which indicates that it may be affected by the evaporites, carbonate, and silicate rocks [72]. That of river water in the Tuotuo River and Chumaer River is Na-Cl, affected by evaporite dissolution [34, 73, 74]. In the Tongtian River, the hydrochemical types are Na-Cl and Ca-HCO₃. That of the trunk stream is Na-Cl, which has the same hydrochemical type as the Chumaer River and Tuotuo River, indicating that their ion has a great impact on that of the Tongtian River.

Gibbs defined the mechanisms controlling world surface water chemistry as three end-members: atmospheric precipitation, rock dominance, and the evaporation-crystallization process. To distinguish the contribution of the three end-members, TDS versus Na⁺/(Na⁺+Ca²⁺) or Cl⁻/(Cl⁻+HCO₃⁻) are used [75]. The Gibbs schematic diagram is shown in Figure 7. The river water of the source region is characterized by moderate or high concentrations of TDS, indicating that the dominant factor is rock weathering and evaporation-crystallization processes, and none of the rivers are affected by precipitation [76]. The diagram reveals that the river waters of Dangqu are characterized by low ratios of Na⁺/(Na⁺+Ca²⁺) and Cl⁻/(Cl⁻+HCO₃⁻) (less than 0.5) and moderate concentrations of TDS (about 100 to 400 mg/L), typical of rock-dominant rivers. Chumaer River is characterized by high ratios of Na⁺/(Na⁺+Ca²⁺) and Cl⁻/(Cl⁻+HCO₃⁻) (approaching 1) and high concentrations of TDS, showing that the controlling factor is evaporation-crystallization processes. However, Tuotuo River and Tongtian River are in an intermediate state, having moderate values of TDS and ratios of Na⁺/(Na⁺+Ca²⁺) and Cl⁻/(Cl⁻+HCO₃⁻), as is the river in the SRLR. They are controlled by rock weathering and evaporation-crystallization processes [32]. These results and conclusions are consistent with the Piper plot (Figure 6).




Fig. 6. Piper plot of major ion composition




Fig. 7. Variation of the ratio of $\text{Na}^+(\text{Na}^++\text{Ca}^{2+})$ and $\text{Cl}/(\text{Cl}+\text{HCO}_3^-)$ as a function of TDS in the Gibbs diagram (a. TDS- $\text{Na}^+(\text{Na}^++\text{Ca}^{2+})$; b. TDS- $\text{Cl}/(\text{Cl}+\text{HCO}_3^-)$)

Analysis of Ion Sources in Water

The way of ions' sources in river water can be divided into natural or anthropogenic. Natural sources are atmospheric transmission and rock weathering. Anthropogenic sources are influenced by human activities such as mineral development, agriculture, and animal husbandry, and so on [70, 77, 78]. The source of river ions can be determined by calculating the ratio of different ions.

(1) Atmospheric Transmission

In the process of atmospheric transport, a large amount of water vapor with many ions comes from the evaporation of seawater. By comparing the values of Na^+/Cl^- and K^+/Cl^- in the source region with the average value of seawater in the world, the contribution of atmospheric transport to the ions in the source region could be judged [79]. The average values of Na^+/Cl^- (1.46) and K^+/Cl^- (0.20) in the river water of the SRYR were much larger than the World Seawater Average ($\text{Na}^+/\text{Cl}^- = 0.86$, $\text{K}^+/\text{Cl}^- = 0.02$). It shows that the Cl was less affected by seawater. Because the SRYR was far away from the ocean and controlled by westerly circulation and monsoon circulation, the contribution and influence of ocean sources to the Cl were both less, and it had no obvious contribution to other ions [80, 81]. Therefore, it was considered that atmospheric transport was not the main way of ions' source in the SRYR.

In the source region of Lancang River, the average value of Na^+/Cl^- (0.74) in river water was slightly less than the World Seawater Average, while K^+/Cl^- (0.04) was slightly larger, which indicated that the natural source route of ions was influenced by other factors besides atmospheric transport [31, 82].

(2) Rock Weathering

Rock weathering with different lithologies can produce different ions, which can reflect how that affects ions [83]. Na^+ and K^+ in natural water mainly come from evaporites or silicate rock weathering. Ca^{2+} and Mg^{2+} may come from evaporites, carbonate, or silicate rock weathering. Cl^- and SO_4^{2-} mainly come from evaporite weathering. And HCO_3^- mainly comes from carbonate rock weathering [84, 85].

In the SRYR, the molar equivalent ratios of Cl^-/Na^+ in the river water were less than or equal to one (Figure 8), which indicated that Na^+ and Cl^- were greatly influenced by the evaporites. It should be paid special attention to that the Cl^-/Na^+ ratio of Dangqu and some areas of the Tongtian River was located at the lower right side of the isogram, which indicated that those areas might be affected not only by evaporites, but also by silicate rock weathering. The ratio of all samples was located above the $(\text{Ca}^{2+}+\text{Mg}^{2+})/\text{HCO}_3^-$ and $(\text{Ca}^{2+}+\text{Mg}^{2+})/(\text{SO}_4^{2-}+\text{HCO}_3^-)$ isograms, and closer to the $(\text{Ca}^{2+}+\text{Mg}^{2+})/(\text{SO}_4^{2-}+\text{HCO}_3^-)$ isograms, indicating that Ca^{2+} and Mg^{2+} were affected not only by carbonate rock weathering, but also by evaporites [86].




Fig. 8. The molar equivalent ratios of different ions (a. Cl^-/Na^+ ; b. $(\text{Ca}^{2+}+\text{Mg}^{2+})/\text{HCO}_3^-$; c. $(\text{Ca}^{2+}+\text{Mg}^{2+})/(\text{SO}_4^{2-}+\text{HCO}_3^-)$; d. $\text{HCO}_3^-/(\text{Cl}^-+\text{SO}_4^{2-})$)

The ratios of the Dangqu and part of the Tongtian River were located around the $\text{HCO}_3^-/(\text{Cl}^-+\text{SO}_4^{2-})$ isogram, which indicated that those areas were affected by carbonate rocks and evaporites [87]. While the ratios of Chumaer River, the Tuotuo River, and part of the Tongtian River were located at the lower right side of the $\text{HCO}_3^-/(\text{Cl}^-+\text{SO}_4^{2-})$ isogram, showing that those areas were effected by evaporites more than carbonate rocks.

In the SRLR, the molar equivalent ratio of Cl^-/Na^+ was about 1 and the ratios of $(\text{Ca}^{2+}+\text{Mg}^{2+})/\text{HCO}_3^-$ and $(\text{Ca}^{2+}+\text{Mg}^{2+})/(\text{SO}_4^{2-}+\text{HCO}_3^-)$ were located at the upper left of the isogram, which indicated that Na^+ and Cl^- were effected by evaporites and Ca^{2+} and Mg^{2+} were influenced by carbonate rocks and evaporites. In addition, according to the ratios of $\text{HCO}_3^-/(\text{Cl}^-+\text{SO}_4^{2-})$, the main ions were more affected by the evaporites than carbonate rocks.

(3) Human Activities Affect

In recent years, with the development of tourism and the increase of grazing activities in Tibetan areas, the hydrochemical characteristics in the source region may change accordingly [88]. The NO_3^- can serve as an indicator of the degree of impact of human activities on the water environment [48, 89]. The NO_3^- content in the river water of the source region was low as a whole. In the SRLR, the NO_3^- content ranged from 1.77 to 2.88 mg/L, with an average of 2.19 mg/L. In the SRYP, that was

0.40~2.71 mg/L, with an average of 1.46 mg/L. The NO_3^- concentration in the natural background value is generally 0.08~0.16 mg/L [90]. In the SRLR and the SRYP, the NO_3^- concentration is higher than the natural background value. Therefore, it may be affected by human input.

To further investigate the impact of human activities on the river water in the source region, the ratios of NO_3^- , SO_4^{2-} , and Cl^- normalized by Na^+ were used to distinguish between human sources and nature sources. However, there was no obvious correlation between $\text{NO}_3^-/\text{Na}^+$, $\text{SO}_4^{2-}/\text{Na}^+$, and Cl^-/Na^+ in river water (Figure 9), which showed that the sources of NO_3^- were different from SO_4^{2-} and Cl^- . The Cl^- and SO_4^{2-} mainly come from evaporite weathering, which belongs to a kind of natural source. Therefore, the NO_3^- may come from human inputs. Anthropogenic sources are usually divided into industrial sources and agricultural sources. Generally, industrial sources are associated with higher $\text{NO}_3^-/\text{Cl}^-$ and $\text{SO}_4^{2-}/\text{Ca}^{2+}$, while agricultural sources are characterized by low NO_3^- and high Cl^- . However, the geographical location of the study area is special, with an average elevation of more than 4000 m, and no industrial sources were found in the study area during the field investigation, so the input of agricultural sources was considered. In Figure 9, the $\text{NO}_3^-/\text{Na}^+$ was obviously much lower than the Cl^-/Na^+ which meant high $\text{Cl}^-/\text{NO}_3^-$. In other words, it meant low NO_3^- and high Cl^- . Therefore, the river water of the source region was affected by agricultural sources to a




Fig. 9. The ratios of NO_3^- , SO_4^{2-} and Cl^- normalized by Na^+ (a. $\text{SO}_4^{2-}/\text{Na}^+ - \text{NO}_3^-/\text{Na}^+$; b. $\text{Cl}^-/\text{Na}^+ - \text{NO}_3^-/\text{Na}^+$)

certain extent [54]. Due to the low content of NO_3^- , it was speculated that human activities had a relatively small impact on the ions.

Distribution Characteristics and Source of Metallic Elements in Sediments

Content and Distribution Characteristics of Metallic Elements in Sediments

Metallic elements are a special group of ions because heavy metals have persistence and potential toxicity to the environment, which can change the activity processes of microorganisms and can be accumulated through sediments and aquatic food chains [91]. The content of metals in sediments is an important factor affecting

river water, because metals can be transformed into each other in sediment water. Analyzing the content, species, and distribution characteristics of metal elements in sediments is helpful to determine the pollution degree of metals in rivers and their potential pollution risks. Therefore, the analysis of metallic elements in sediment is very important for assessing the state of the river.

Compared with other metallic elements, the average contents of Ti, Mn, Fe, Sr, and Ba in the sediment of the source region were relatively higher (Figure 10). In the SRYR, the average content of Fe in the sediment was 19758 mg/kg and Ti was 2581 mg/kg. In the SRLR, that of Fe was 18736 mg/kg and Ti was 2533mg/kg. It was mainly due to the high background value of some metallic elements in river sediments in the source region [46, 92].




Fig. 10. The box-plot of the metallic elements' average contents in sediment

In the SRYR, the average contents of Cd, Mo, Sr, Se, Cu, Co, Fe, Mn, Cr, V, Ti, and Li in the sediments of different rivers were basically flat, while those of the others were different in spatial distribution (Figure 11). In the sediments of the Dangqu River, the average contents of Hg, As, Ba, Sb, Ni, and Be were higher than those in the Tuotuo River, Chumaer River, and Tongtian River. In the sediments of the Chumaer River, the average contents of Hg (below the detection limit), As, Pb, Sb, Zn, Ni, and Be were lower than those in other rivers. In the sediments of the Tuotuo River, Hg, and Ba were below the detection limit, and the average contents of other metallic elements were flat with those of other rivers. In the sediments of the Tongtian River, only the average content of Hg was obviously higher than that of other rivers, and others were flat or lower.

Compared with the average background value of Chinese river sediment [93, 94], in the sediments of the Dangqu River, the average contents of Pb, Ba, Zn, Cu, Co, Fe, Mn, V, and Ti were obviously lower, Cd, Mo, and Ni were basically flat, and Hg, As, Sb, Sr, Se, Cr, Be, and Li were higher. Except for Se and Cr, the average contents of other metallic elements in the Chumaer River were lower than the average background value of Chinese river sediment. In the Tuotuo River, the average contents of As, Sr, Se, and Cr in sediments were obviously higher, while Sb, Cd, Mo, and Cr were flat and other metallic elements were lower. In the Tongtian River, the average contents of Hg, As, Sr, Se, and Cr in sediments were

relatively higher, while Sb, Cd, Mo, and Li were flat. The other metallic elements were lower than the average background value of Chinese river sediment.

In the SRLR, the average contents of Cr and Cd in sediments were slightly higher than the background values of sediments in the China River system (Figure 12), while V, Ni, Cu, Zn, Pb, Mn, and Fe were lower. Some studies indicate that it was the rocks containing Cd and Cr that affected the content of metallic elements in sediments [95, 96].

As shown in Figure 12, the average contents of metallic elements in the sediments in the SRYR were basically flat compared to those in the SRLR. Compared with the source regions of the Yellow River and other rivers in alpine and cold regions, the average contents of V, Ni, and Zn in the sediments of the source regions were slightly lower, while Cr was slightly higher. Cd was lower than that in the source region of the Yellow River, but higher than that in rivers in alpine and cold regions. Pb, Mn, and Fe were higher than those in the source region of the Yellow River, but lower than those in rivers in alpine and cold regions.

Analysis of the Source of Metallic Elements in Sediments

Correlation analysis can be used to infer the homology between various metallic elements and reveal the source relationship of ions based on the significant level of




Fig. 11. Distribution of metallic elements in river sediments in the SRYR




Fig. 12. Metallic elements' average contents in sediments of different areas

correlation [60, 92]. Considering that the amount of samples affects the analysis accuracy, the metallic elements in the sediments of rivers in the source region measured by our study team in the past five years (2019~2022) were selected to complete the relevant analysis.

The Pearson Correlation Coefficient is shown in Table 5. In the SRLR, the results show that there was a significant positive correlation ($p < 0.05$) between Fe and Ti, V; Pb, V, Co, Sr, Sb, Co, Cr, and Ni were pairwise correlated; Cu, Sr, Cb, and Sb were pairwise correlated; Be, Mn, and Ni were pairwise correlated. It indicated that these metallic elements had a certain degree of correlation with each other, and their sources might have similarities and homology. In the SRYR, Li and Be were significantly positively correlated with most ions; T, V, Mn, Fe, Co, and Cu were pairwise correlated; Mn and Co were significantly positively correlated with Ni, Se, and Pb; there was a significant positive correlation in Se-Sr and Sr-Cd-Pb. It indicated that these metallic elements were highly correlated ($p < 0.01$) and might have similar sources. However, not all metallic elements had a significant correlation, which indicated that their sources were not completely consistent.

In order to further analyze the main sources of metallic elements, the principal component analysis method was used to analyze the sediments in the source region. Under the premise of a 95% confidence level, three principal components (eigenvalues > 1) of sediments were extracted, respectively. In the source region of the Lancang River,

the contribution rates of the first and second principal components were 67.15% and 20.65%, and the cumulative contribution rate was 87.8%. In the SRYR, the contribution rates of the first and second principal components of sediments were 45.97% and 35.05%, and the cumulative contribution rate was 81.02%.

The results of the principal component analysis are shown in Table 6. In the SRLR, the first principal component contribution rate of the metallic elements in river sediments (67.15%) was much higher than that of other principal components. The metallic elements with a higher load were Be, Ti, V, Cr, Fe, Co, Ni, Cu, Zn, Sr, Cd, Sb, and Pb, and the metallic elements of the second principal component were Li and Mn. In the SRYR, the first principal components of metallic elements with a higher load in river sediments were Li, Be, V, Mn, Fe, Co, and Ni, and the metallic elements of the second principal component were Zn, Sr, Cd, Sb, and Pb. According to the correlation analysis, these metallic elements had a significant correlation, which indicated that their sources were similar or the same. According to the literature research, Fe and Mn were abundant macroelements in the crust [97, 98]; Ni and Co were controlled by the geological background of the Qinghai-Tibet Plateau [99]; Pb, Zn, and Ba were often used as identification elements of traffic pollution sources [100, 101]. Although the values of Pb, Zn, and Ba in the study area were lower than the background values of sediments in China, there was a significant correlation between Sr and Sb with Pb and Zn, while the values of Sr and Sb were higher than

Table 5. Pearson correlation coefficient of metallic elements in sediments

Areas	Li	Be	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Se	Sr	Cd	Sb	Pb
The SRLR	Li	1														
	Be	0.089	1													
	Ti	-0.42	0.543	1												
	V	-0.443	0.774	0.92	1											
	Cr	-0.111	0.924	0.823	0.937	1										
	Mn	0.297	0.977*	0.454	0.659	0.872	1									
	Fe	-0.299	0.696	0.980*	0.954*	0.918	0.623	1								
	Co	-0.188	0.928	0.809	0.952*	0.994**	0.857	0.902	1							
	Ni	0.063	0.976*	0.693	0.848	0.978*	0.955*	0.823	0.967*	1						
	Cu	-0.704	0.643	0.681	0.876	0.729	0.466	0.70	0.795	0.637	1					
	Zn	-0.872	0.185	0.807	0.754	0.481	0.006	0.708	0.52	0.292	0.784	1				
	Se	0.706	-0.297	-0.084	-0.382	-0.215	-0.115	-0.095	-0.32	-0.175	-0.778	-0.459	1			
	Sr	-0.573	0.764	0.74	0.934	0.841	0.612	0.784	0.892	0.768	0.983*	0.728	-0.682	1		
	Cd	-0.789	0.533	0.754	0.881	0.688	0.347	0.737	0.749	0.562	0.981*	0.889	-0.715	0.952*	1	
	Sb	-0.655	0.664	0.845	0.961*	0.822	0.504	0.853	0.865	0.711	0.967*	0.85	-0.596	0.976*	0.978*	1
	Pb	-0.438	0.819	0.865	0.993**	0.941	0.7	0.915	0.965*	0.867	0.909	0.715	-0.464	0.964*	0.893	0.966*
The SRYR	Li	1														
	Be	0.836**	1													
	Ti	0.641*	0.521	1												
	V	0.720**	0.529	0.960**	1											
	Cr	0.431	0.760**	0.214	0.102	1										
	Mn	0.842**	0.957**	0.584*	0.581*	0.765**	1									
	Fe	0.836**	0.680*	0.919**	0.967**	0.268	0.731**	1								
	Co	0.894**	0.901**	0.790**	0.782**	0.615*	0.894**	0.883**	1							
	Ni	0.700**	0.900**	0.468	0.391	0.867**	0.847**	0.538	0.866**	1						
	Cu	0.748**	0.827**	0.850**	0.781**	0.582*	0.815**	0.818**	0.926**	0.817**	1					
	Zn	0.316	0.186	0.001	0.182	-0.25	0.146	0.232	0.085	-0.137	-0.035	1				
	Se	0.438	0.699**	0.416	0.268	0.869**	0.688**	0.394	0.707**	0.896**	0.696**	-0.449	1			
	Sr	0.19	0.589*	-0.082	-0.156	0.758**	0.50	0.01	0.391	0.713**	0.364	-0.007	0.645*	1		
	Cd	-0.178	-0.362	-0.411	-0.23	-0.580	-0.366	-0.237	-0.471	-0.628	-0.565	0.806**	-0.830	-0.311	1	
	Sb	0.346	0.07	-0.073	0.147	-0.351	0.11	0.234	0.032	-0.212	-0.177	0.716**	-0.418	-0.223	0.672*	1
	Pb	0.629*	0.776**	0.176	0.268	0.422	0.724**	0.433	0.569*	0.528	0.44	0.653*	0.262	0.553*	0.188	0.455

Note: **, extremely significant correlation ($p < 0.01$); *, significant correlation ($p < 0.05$).

Table 6. Principal component analysis of metallic elements in sediments

Areas	Principal Constituent	Load Factor															
		Li	Be	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Se	Sr	Cd	Sb	Pb
The SRLR	I	-0.554	0.755	0.845	0.98	0.885	0.612	0.878	0.922	0.796	0.954	0.78	-0.558	0.985	0.947	0.991	0.99
	II	0.828	0.579	0.115	0.136	0.463	0.738	0.269	0.386	0.598	-0.233	-0.489	0.547	-0.052	-0.322	-0.124	0.136
	III	-0.09	-0.308	0.522	0.145	0.041	-0.285	0.396	-0.027	-0.093	-0.189	0.391	0.624	-0.166	-0.017	0.041	0.023
The SRYR	I	0.924	0.962	0.548	0.714	-0.864	0.879	0.835	0.89	0.948	0.581	0.546	-0.729	-0.378	0.315	0.589	0.358
	II	0.207	0.063	-0.755	-0.688	0.118	0.086	-0.544	-0.454	-0.219	-0.663	0.73	-0.564	0.701	0.852	0.703	0.744
	III	-0.115	0.188	0.137	0.012	-0.444	0.365	-0.016	0.022	-0.203	0.228	-0.078	0.23	0.423	-0.289	0.015	0.564

Note: I, II and III represent the first, the second and the third principal component; Bold data indicates high load

the background values of sediments in China. Therefore, Sr and Sb might be affected by traffic pollution sources. In conclusion, it was considered that the metallic elements in the sediments of the source region were controlled by both natural and traffic pollution.

Conclusion

In general, the river water in the source region was medium or high-hardness fresh water. The water of the Chumaer River was high-hardness brackish water due to geological conditions and engineering impacts. The main cations in the water of the source region were Ca^{2+} and Na^+ ; and the main anions were Cl^- , SO_4^{2-} and HCO_3^- . There were some differences in the content of metallic elements in the water. The average contents of As, Cu, Pb, Cr, Cd, Ni, and Ti in the river water of the SRYR were higher than those in the SRLR.

The main hydrochemical types in the source region are Na-Cl and $\text{Ca-SO}_4\cdot\text{HCO}_3$, which are mainly controlled by rock weathering and evaporating crystallization. However, it was diverse in different regions. The hydrochemical type of the river water in Dangqu was $\text{Ca-SO}_4\cdot\text{HCO}_3$, controlled by rock weathering; in the Tuotuo River and the Chumaer River, this was Na-Cl, controlled by evaporating crystallization; and in Tongtian River, this was Na-Cl and Ca-HCO_3 , controlled by rock weathering and evaporating crystallization. The ions in the river water of the source region were less affected by atmospheric transport and human activities, but more affected by rock weathering. In the SRLR, the water ions were mainly affected by evaporites and carbonate rocks. And in the SRYR, those were mainly affected by evaporites, silicates, and carbonate rocks.

In the source region, the average contents of Ti, Mn, Fe, Sr, and Ba in the surface sediments of rivers were higher than those of other metallic elements. Compared with the background value of sediments in China, the average contents of Cr and Cd in the source area were higher. Compared with the source region of the Yellow River and other rivers in alpine and cold regions, the average contents

of V, Ni, and Zn were slightly lower. In the SRYR, the average contents of metallic elements in the sediments of different rivers were different. In Dangqu, Hg, As, Ba, Sb, Ni, and Be were higher. In the Chumaer River, Hg was below the detection limit; in the Tuotuo River, Hg and Ba were both below the detection limit; and in the Tongtian River, the average content of Hg was higher than in other rivers. A preliminary analysis of the sources of metallic elements in the sediments showed that the metallic elements were controlled by natural and traffic pollution.

Acknowledgments

This study was supported by the State-level Public Welfare Scientific Research Institutes Basic Scientific Research Business Project of China (Grant No. CKSF2023311+SH, CKSF2021743/HL, CKSF2023337/SH.).

Conflict of Interest

The authors declare no conflict of interest.

References

- LIU H., LIU E., YU Z., LIN Q., ZHANG E., SHEN J. Spatio-temporal accumulation patterns of trace metals in sediments of a large plateau lake (Erhai) in Southwest China and their relationship with human activities over the past century. *Journal of Geochemical Exploration*, **234**, 106943, 2022.
- VIANELLO P., ANSORGE I., ROUAULT M., OSTROWSKI M. Transport and transformation of surface water masses across the Mascarene Plateau during the Northeast Monsoon season. *African Journal of Marine Science*, **39** (4), 453, 2017.
- LIU J., GUO H. Hydrochemical Characteristics and Ion Source Analysis of the Yarlung Tsangpo River Basin. *Water*, **15** (3), 537, 2023.
- HUANG J., LIU R., WANG Q., GAO X., HAN Z., GAO J., GAO H., ZHANG S., WANG J., ZHANG L., XIA X. Climate factors affect N_2O emissions by influencing the migration and transformation of nonpoint source nitrogen in an agricultural watershed. *Water Research*, **223**, 119028, 2022.

5. JAHAN A., KHAN M.U., RAI N., KUMAR S., ALI D.T. Geochemical characterization, its controlling factors, and comparison between the upstream and downstream segments of the Himalayan Satluj River basin, India. *Geochemistry*, **83** (2), 125974, **2023**.
6. LI Z., FENG Q., WANG Q., YONG S., LI J., LI Y., WANG Y. Quantitative evaluation on the influence from cryosphere meltwater on runoff in an inland river basin of China. *Global and Planetary Change*, **143**, 189, **2016**.
7. JIANG L., YAO Z., LIU Z., WANG R., WU S. Hydrochemistry and its controlling factors of rivers in the source region of the Yangtze River on the Tibetan Plateau[J]. *Journal of Geochemical Exploration*, **155**, 76, **2015**.
8. KUMARASAMY P., ARTHUR JAMES R., DAHMS H., BYEON C., RAMESH R. Multivariate water quality assessment from the Tamiraparani river basin, Southern India[J]. *Environment Earth Science*, **71**, 2441, **2014**.
9. LIU W., SUN G., LI Y., XU Z. Hydrochemistry and carbon isotope characteristics of Nujiang River water: Implications for CO₂ budgets of rock weathering in the Tibetan Plateau[J]. *Scientia Sinica Terrae*, **53** (12), 2992, **2023** [In Chinese].
10. WOOJIN S., JONGSIK R., YOUNGYUN P., KWANG-SIK L. Chemical weathering and associated CO₂ consumption in six major river basins, South Korea[J]. *Geomorphology*, **129** (3-4), 334, **2011**.
11. SABINE H., JAN S. Pollution history revealed by sedimentary records: a review. *Environmental chemistry letters*, **11** (3), 255, **2013**.
12. GUO B., LIU Y., ZHANG F., HOU J., ZHANG H. Characteristics and Risk Assessment of Heavy Metals in Core Sediments from Lakes of Tibet. *Environmental Science*, **37** (02), 490, **2016** [In Chinese].
13. SAHA AJOY., VIJAYKUMAR M.E., DAS B.K., SAMANTA S., KHAN M.F., KAYAL T., JANA C., CHOWDHURY A.R. Geochemical distribution and forms of phosphorus in the surface sediment of Netravathi-Gurupur estuary, southwestern coast of India. *Marine Pollution Bulletin*, **187**, 114543, **2023**.
14. SHAO Y., HE Q., FU Y., LIU Y. Construction of the comprehensive evaluation system of waterbody pollution degree and the response of sedimentary microbial community. *Environmental Pollution*, **317**, 120837, **2023**.
15. YANG Z., LI C., CHEN H., SHAN X., CHEN J., ZHANG J., LIU S., LIU Q., WANG X. Source-oriented ecological and resistome risks associated with geochemical enrichment of heavy metals in river sediments. *Chemosphere*, **336**, 139119, **2023**.
16. ZHAO L., LI W., LIN L., GUO W., ZHAO W., TANG X., GONG D., LI Q., XU P. Field Investigation on River Hydrochemical Characteristics and Larval and Juvenile Fish in the Source Region of the Yangtze River. *Water*, **11** (7), 1342, **2019**.
17. WU Y., WANG S., NI Z., LI H., MAY L., PU J. Emerging water pollution in the world's least disturbed lakes on Qinghai-Tibetan Plateau. *Environmental Pollution*, **272**, 116032, **2020**.
18. LIU Z., PAN B., HAN X., LI G., WANG, T. Water Environmental Characteristics and Water Quality Assessment of Lakes in Tibetan Plateau. *Environmental Science*, **43** (11), 5073, **2022** [in Chinese].
19. WANG C., ZHAO W., CUI Y. Changes in the Seasonally Frozen Ground Over the Eastern Qinghai-Tibet Plateau in the Past 60 Years. *Frontiers in Earth Science*, **8**, 270, **2020**.
20. CUI B., LI X. Characteristics of stable isotopes and hydrochemistry of river water in the Qinghai Lake Basin, northeast Qinghai-Tibet Plateau, China. *Environmental Earth Sciences*, **73** (8), 4251, **2015**.
21. HAN R., LIU W., ZHANG J., ZHAO T., SUN H., XU Z. Hydrogeochemical characteristics and recharge sources identification based on isotopic tracing of alpine rivers in the Tibetan Plateau. *Environmental Research*, **229**, 115981, **2023**.
22. SHI W., LIU J., LI B., CHEN X. Hydrochemical characteristics and solute sources of the Yanhu Areas on the Qinghai-Tibet Plateau. *Environmental Chemistry*, **42** (01), 101, **2023** [In Chinese].
23. KAUSHIK H., RAMANATHAN A.L., SOHEB M., SHAMURAILATPAM M.S., BISWAL K., MANDAL A., SINGH C. Climate change-induced high-altitude lake: Hydrochemistry and area changes of a moraine-dammed lake in Leh-Ladakh. *Acta Geophysica*, **69** (6), 1, **2021**.
24. SHI X., CHEN X., GAO M., LI G. Study on the Spatiotemporal Pattern and Controlling Factors of River Water Hydro-Chemical Characteristics in the Lhasa River Basin, Tibet Plateau. *Resources and Environment in the Yangtze Basin*, **32** (01), 183, **2023** [In Chinese].
25. WANG P., ZHAO J., XIAO H., YANG W., YU X. Bacterial community composition shaped by water chemistry and geographic distance in an anthropogenically disturbed river. *Science of the Total Environment*, **655**, 61, **2019**.
26. LI Q., SHI X., ZHAO Z., WU Q. Ecological restoration in the source region of Lancang River: Based on the relationship of plant diversity, stability and environmental factors. *Ecological Engineering*, **180**, 106649, **2022**.
27. LI Y., LI B., YUAN Y., LEI Q., JIANG Y., LIU Y., LI R., LIU W., ZHAI D., XU J. Trends in total nitrogen concentrations in the Three Rivers Headwater Region. *The Science of the Total Environment*, **852**, 158462, **2022**.
28. RONNY B., DU Y., DONG A., LINUS Z., YUAN F., HAO Z. Integrated large-scale circulation impact on rainy season precipitation in the source region of the Yangtze River[J]. *International Journal of Climatology*, **40** (4), 2285, **2020**.
29. BIN Q., MIKE S., ZHANG Y., GUO J., MAHMOUD S.M., ABDEL W., KANG S. Water chemistry of the headwaters of the Yangtze River. *Environmental Earth Sciences*, **74**, 6443, **2015**.
30. CHENG H., ZHANG Z. Discussion on the Source of the Yangtze River, Yellow River, and Lancang River. *Management & Strategy of Qinghai Land & Resources*, **2**, 31, **2009** [In Chinese].
31. HUANG C., LI Y., LIU G., ZHANG H., LIU Q. Recent climate variability and its impact on precipitation, temperature, and vegetation dynamics in the Lancang River headwater area of China. *International Journal of Remote Sensing*, **35** (8), 2822, **2014**.
32. JIANG L., YAO Z., LIU Z., WANG R., WU S. Hydrochemistry and its controlling factors of rivers in the source region of the Yangtze River on the Tibetan Plateau[J]. *Journal of Geochemical Exploration*, **155**, 76, **2015**.
33. LI H., ZHAO W., ZHAO L., LI W., CAO X. Study on the Distribution Characteristics of Phytoplankton in Summer in the Source Region of Yangtze River and Lancang River. *Journal of Hydroecology*, **43** (03), 77, **2022** [In Chinese].
34. LIU M., ZHAO L., LI Q., ZOU J., HU Y., ZHANG Y., XU P., WU Z., DENG W., TAO J.X. Hydrochemical characteristics, main ion sources of main rivers in the source region of Yangtze River. *China Environmental Science*, **41** (03), 1243, **2021** [In Chinese].
35. YAO Z., SUN G., LIN L., ZHANG W., ZHANG J., GAO L., SHANG L. Distribution, Source Identification, and Output flux of Barium in Surface Waters in the Sanjiangyuan Region and Qilian Mountain Region of Tibetan Plateau. *Bulletin of Environmental Contamination and Toxicology*, **111** (1), 7, **2023**.
36. ZHAO L., DENG W., LIU M., HU Y., XIE L. Preliminary Research on Hydrogen and Oxygen Isotope Characteristics

- of River Waters in the Source Region of the Yangtze River and the Lancang River. Springer Proceedings in Earth and Environmental Sciences, pp. 67, **2021**.
37. WALTER I., LUDOVICUS B., MARC B. Climate Change Will Affect the Asian Water Towers. *Science*, **328** (5984), 1382, **2010**.
 38. ZHANG X., JIN X. Vegetation dynamics and responses to climate change and anthropogenic activities in the Three-River Headwaters Region, China. *Ecological Indicators*, **131**, 108223, **2021**.
 39. XU P., LI Q., HUANG Z., ZHAO W., ZHOU Y., REN F., ZHAO D., LU Y. Comprehensive scientific investigation of ecological environment in the source areas of the Yangtze River and Lancang River (2012~2016). Science Press: Beijing, **2019** [In Chinese].
 40. WU Z., XU P., ZHAO L. Integrated Scientific Expedition Report on the Headwaters of the Yangtze River 2019. Changjiang Press: Wuhan, **2020** [In Chinese].
 41. LIU X., WU Z., LIU Y., ZHAO X., RUI Y., ZHANG J. Spatial-temporal characteristics of precipitation from 1960 to 2015 in the Three Rivers' Headstream Region, Qinghai, China. *Acta Geographica Sinica*, **74** (09), 1803, **2019** [In Chinese].
 42. HUANG Z., LIU Y., ZHAO W., ZHAO L. Discussion on Recent Spatial-temporal Distribution of Water Quality in Changjiang River Source Area. *Journal of Changjiang River Scientific Research Institute*, **33** (07), 46, **2016** [In Chinese].
 43. LU D., ZHANG J., GENG Z., MIN M. Community structures of zooplankton and assessment of water quality in important reaches and lakes of Three-River head-waters region. *Yangtze River*, **50** (05), 41, **2019** [In Chinese].
 44. YANG A., WANG Y., HU J., LIU X., LI J. Evaluation and Source of Heavy Metal Pollution in Surface Soil of Qinghai-Tibet Plateau. *Environmental Science*, **41** (02), 886, **2020** [In Chinese].
 45. LIU M., LEI J., LIN L., HUANG Z., CAO X., TAO J., YANG W. Metal element analysis in complicated media based on MP-AES. *Yangtze River*, **49** (S1), 43, **2018** [In Chinese].
 46. LIU M., DENG W., ZHAO L., HU Y., H H., GAO F. Distribution Characteristics and Sources of Heavy Metals in Surface Sediments and Bank Soils of Major Rivers in Source Region of Yangtze River. *Journal of Changjiang River Scientific Research Institute*, **38** (07), 143, **2021** [In Chinese].
 47. WEN Z., SU R., XIAN P. Determination of Multiple Elements in Stream Sediment by Inductively Coupled Plasma-Mass Spectrometry with Microwave Digestion. *The Administration and Technique of Environmental Monitoring*, **28** (04), 39, **2016** [In Chinese].
 48. GAILLARDETA J., VIERS J., DUPRÉ B. *Treatise on Geochemistry*, 2nd ed., Elsevier Ltd., pp. 195, **2014**.
 49. RAYMO M.E., RUDDIMAN W.F. Tectonic forcing of late Cenozoic climate[J]. *Nature*, **359** (6391), 117, **1992**.
 50. JOEL D.B. The Effect of Late Cenozoic Glaciation and Tectonic Uplift on Silicate Weathering Rates and the Marine ⁸⁷Sr/⁸⁶Sr Record[A] // WILLIAM F.R. Tectonic Uplift and Climate Change[C]. Plenum Press: New York, pp. 260, **1997**.
 51. LU T., LI R., FERRER A.S., XIONG S., ZOU P., PENG H. Hydrochemical characteristics and quality assessment of shallow groundwater in Yangtze River Delta of eastern China. *Environmental Science and Pollution research International*, **29** (38), 57215, **2022**.
 52. MA B., ZHANG Q., LI S. Hydrochemical characteristics and controlling factors of trans-boundary rivers in China. *Quaternary Sciences*, **43** (02), 425, **2023** [In Chinese].
 53. ZHANG X., MA P., GAO Q., YAN H., QIAN B. Spatial-temporal variations of water quality in upstream and downstream of Three Gorges Dam. *Journal of Lake Sciences*, **31** (03), 633, **2019** [In Chinese].
 54. LI Z., LI Z., SONG L., GUI J., XUE J., ZHANG B., GAO W. Precipitation chemistry in the Source Region of the Yangtze River. *Atmospheric Research*, **245**, 105073, **2020**.
 55. FU C., LIU C. Hydrochemical characteristics and formation mechanism of saline lakes in Hoh Xil region. *Yangtze River*, **53** (7), 36, **2022** [In Chinese].
 56. LIU W., XIE C., WANG W., ZHANG Y., YANG G., LIU G. Analysis on expansion trend and outburst risk of the Yanhu Lake in Hoh Xil region, Qinghai-Tibet Plateau. *Journal of Glaciology and Geocryology*, **41** (06), 1467, **2019** [In Chinese].
 57. MA B., ZHANG Q., LI S. Hydrochemical characteristics and controlling factors of trans-boundary rivers in China. *Quaternary Sciences*, **43** (02), 425, **2023** [In Chinese].
 58. LI M., SUN X., LI S., ZHANG Q. Research progress on inorganic hydrochemical characteristics of glacial meltwater runoff in Qinghai-Tibet Plateau and its surrounding areas. *Journal of Glaciology and Geocryology*, **42** (02), 562, **2020** [In Chinese].
 59. MAL., BAI Y., NIU Z., YAO H., DUAN Q. Strontium and neodymium isotope geochemistry of Late Triassic volcanic rocks in Geladandong area, the source of the Yangtze River, and its significance. *Geology in China*, **2**, 174, **2004** [In Chinese].
 60. QIAO S., WANG T., ZHANG Q., LIU X., ZHAO M. Distribution Characteristics and Risk Assessment of Heavy Metals in the Source Region of Yangtze River (In Chinese) [J]. *Acta Scientiarum Naturalium Universitatis Pekinensis*, **58** (02), 297, **2022**.
 61. XIAO D., SHAO X., QIN N., HUANG X. Tree-ring-based reconstruction of streamflow for the Zaqu River in the Lancang River source region, China, over the past 419 years. *International Journal of Biometeorology*, **61** (7), 1173, **2017**.
 62. ZHU J. Exploring the source of the Yangtze River (Yangtze River Tour). *Journal of Yangtze River Scientific Research Institute*, **40** (04), 2, **2023** [In Chinese].
 63. CHENG H. Characteristics of heavy metals in surface water and evaluation of water environment quality in the upper reaches of reservoirs in the source region of the Yellow River in recent 10 years. *Chinese Society for Environmental Sciences*, 660, **2019** [In Chinese].
 64. ZUO H., MA X., YANG K., CHEN Y., CHEN J., GUO Y., ZHAO J., WANG R., FANG F., LIU Y. Distribution and Risk Assessment of Metals in Surface Water and Sediment in the Upper Reaches of the Yellow River, China. *Soil and Sediment Contamination: An International Journal*, **25** (8), 917, **2016**.
 65. WU W., RAN X., LI J., WANG H., LI M., LIU J., ZANG J. Sources, Distribution, and Fluxes of Major and Trace Elements in the Yangtze River. *Environmental Science*, **40** (11), 4900, **2019** [In Chinese].
 66. FU D., WANG X., LIU J., ZHANG A., SUN Z., FU Q., SHU J. The Investigate and Analysis about Background Value of Water Environmental in the Source Area of Yangtze River[J]. *Environmental Monitoring in China*, **1**, 11, **1998** [In Chinese].
 67. ZHANG L., ZHUN K. Background values of trace elements in the source area of the Yangtze River[J]. *Science of the Total Environment*, **125**, 391, **1992**.
 68. MAO J., LI X., LI H., QU X., ZHANG C., XUE C., WANG Z., YU J., ZHANG Z., FENG C., WANG R. Types and characteristics of endogenetic metallic deposits in orogenic belts in China and their metallogenic processes[J]. *Acta Geologica Sinica*, **3**, 342, **2005** [In Chinese].
 69. CHU H., WEI J., QIU J., LI Q., WANG G. Identification of the impact of climate change and human activities on rainfall-runoff relationship variation in the Three-River Headwaters region. *Ecological Indicators*, **106**, 105516, **2019**.
 70. LI Z., MA J., YU H., YANG H., SONG L., LI Z., GUI J. Geochemical evidence of ions' sources and influences of meteorological factors on hydrochemistry of glacier snow meltwa-

- ter in the source region of the Yangtze River. *Environmental Earth Sciences*, **79** (10), 235, **2020**.
71. LIU M., ZHAO L., LI Q., HU Y., HUANG H., ZOU J., GAO F., TAO J., ZHANG Y., XU P., WU Z., YU C. Distribution characteristics, enrichment patterns and health risk assessment of dissolved trace elements in river water in the source region of the Yangtze River. *Journal of Water and Climate Change*, **12** (6), 2288, **2021**.
 72. WANG Q., YU S., JIANG P., SUN P. Water Chemical Characteristics and Influence of Exogenous Acids in the Yangtze River Basin. *Environmental Science*, **42** (10), 4687, **2021** [In Chinese].
 73. CAO Y., ZHANG M., QU D., WANG S., QIU X., MA Z., DU Q., CHE C. Temporal-spatial variation of surface and underground water chemistry in the eastern part of Qilian Mountains. *China Environmental Science*, **40** (4), 1667, **2020** [In Chinese].
 74. ZHANG L., ZHAO Z., ZHANG W., TAO Z., HUANG L., YANG J., WU Q., LIU C. Characteristics of water chemistry and its indication of chemical weathering in Jinshajiang, Lancangjiang and Nujiang drainage basins. *Environmental Earth Sciences*, **75** (6), 506, **2015**.
 75. JAHAN A., KHAN M.U., RAI N., KUMAR S., ALI D.T. Geochemical characterization, its controlling factors, and comparison between the upstream and downstream segments of the Himalayan Satluj River basin, India. *Geochemistry*, **83** (2), 125974, **2023**.
 76. LIU J., LI Y., GAO Z., WANG M., MA Y., SHI M., ZHANG H. Ionic composition characteristics and their controlling factors in the main stream of the Yarlung Zangbo River. *Journal of Glaciology and Geocryology*, **41** (03), 697, **2019** [In Chinese].
 77. WANG S., HE X., WU J., DING Y., HU Z., WANG L., YANG G. Chemical Characteristics and Ionic Sources of Precipitation in the Source Region of the Yangtze River. *Environmental Science*, **40** (10), 4431, **2019**.
 78. YAN Y., NIU F., LIU J., L X., LI Y., PENG H., YAN D., XIAO S. Hydrochemical characteristics and sources of the upper Yarlung Zangbo River in summer. *China Environmental Science*, **42** (02), 815, **2022** [In Chinese].
 79. BERNER E.K., BERNER R. *Global Environment: Water, Air, and Geochemical Cycles*, 2nd ed., Princeton University Press, pp. 221, **2012**.
 80. LI Z.J., LI Z.X., SONG L.L., GUI J., XUE J., ZHANG B., GAO W. Precipitation chemistry in the Source Region of the Yangtze River. *Atmospheric Research*, **245**, 105073, **2020**.
 81. LI Z., DUAN R., KE H., LIU X., XIE G., GAO W., SONG L., ZHANG B., GUI J., XUE J., NAN F., LIANG P., LI Y. Research progress of ecological hydrology based on hydrochemical characteristics in the source region of the Yangtze River. *Journal of Glaciology and Geocryology*, **44** (01), 288, **2022** [In Chinese].
 82. MA B., ZHANG Q., LI S. Hydrochemical characteristics and controlling factors of trans-boundary rivers in China. *Quaternary Sciences*, **43** (02), 425, **2023** [In Chinese].
 83. RAMESH R.P., ZHANG F., REHMAN F.U., WANG G., YE M., CHEN Z., TANG H. Spatiotemporal variations of hydrogeochemistry and its controlling factors in the Gandaki River Basin, Central Himalaya Nepal. *Science of the Total Environment*, **622**, 770, **2018**.
 84. LI B., BAO Y., HU M., WANG Y., SUN M. Hydrochemical characteristics and carbon sink calculation in Yarlung Zangbo River Basin. *Yangtze River*, **54** (04), 49, **2023** [In Chinese].
 85. NOH H., HUH Y., QIN J., ANDRE E. Chemical weathering in the Three Rivers region of Eastern Tibet. *Geochimica et Cosmochimica Acta*, **73** (7), 1857, **2009**.
 86. JOBIN T., SABU J., THRIVIKRAMJI K.P., MANJUSREE T.M., ARUNKUMAR K.S. Seasonal variation in major ion chemistry of a tropical mountain river, the southern Western Ghats, Kerala, India. *Environmental earth sciences*, **71** (5), 2333, **2014**.
 87. MOHAMED E.A., AREF L., FATHY A., ABDULAZIZ A.B. Assessing the hydrogeochemical processes affecting groundwater pollution in arid areas using an integration of geochemical equilibrium and multivariate statistical techniques. *Environmental Pollution*, **229**, 760, **2017**.
 88. LI Z.J., LI Z.X., SONG L., YONG J. Environment significance and hydrochemical characteristics of supra-permafrost water in the source region of the Yangtze River. *Science of the Total Environment*, **644**, 1141, **2018**.
 89. ZHOU L., JUN X., JAIVIME E., ZHI L. Spatiotemporal variations in the hydrochemical characteristics and controlling factors of stream flow and groundwater in the Wei River of China. *Environmental Pollution*, **254**, 113006, **2019**.
 90. FEI C., LIJUN H., MIN L., YANLING Z., GUOYU Y., XIANBIAO L., XIAOFEI L., HAIBO Z., FENGYU D., JUAN G., XIAOFEN J. Net anthropogenic nitrogen inputs (NANI) into the Yangtze River basin and the relationship with riverine nitrogen export[J]. *Journal of Geophysical Research: Biogeosciences*, **121** (2), 451, **2016**.
 91. LI S., SUN W., CHEN R., ZHANG Z., NING D., NI Z. A historical record of trace metal deposition in northeastern Qinghai-Tibetan Plateau for the last two centuries. *Environmental Science and Pollution Research*, **29** (17), 24716, **2021**.
 92. YANG A., XING W., WANG X., HU J., LIU X., LI J. Source and risk assessment of heavy metals in surface sediments of rivers, lakes and their surrounding soils in central Tibet. *China Environmental Science*, **40** (10), 4557, **2020** [In Chinese].
 93. SHI C., LIANG M., FENG B. Average Background Values of 39 Chemical Elements in Stream Sediments of China. *Earth Science*, **41** (02), 234, **2016** [In Chinese].
 94. YAN M., CHI Q., GU T., WANG C. Average Element Content of various sediments in China. *Geophysical and Geochemical Exploration*, **6**, 468, **1995** [In Chinese].
 95. LI H., YANG D., TIAN Y., LI Y., WANG L., LI B. Genesis and Its Geodynamic Significance of Late Cretaceous Granites in North Lancang River Suture. *Earth Science*, **48** (04), 1330, **2023** [In Chinese].
 96. ZHOU H., ZHOU H., XIAO H., MA Y., LI L., ZHANG D., CHEN J. The Variation Characteristics of Heavy Metal Content, Nutrient and Enzyme Activity in Soil of Alpine Steppe with Different Degradation Gradient in the Three River-Headwaters Region. *Acta Agrestia Sinica*, **28** (03), 784, **2020** [In Chinese].
 97. LI T. Statistical characteristics of crustal element abundance. *Geology and Exploration*, **10**, 7, **1992** [In Chinese].
 98. LI T. Element abundances of China's continental crust and its sedimentary layer and upper continental crust. *Chinese Journal of Geochemistry*, **14** (1), 26, **1995**.
 99. ROBERT S., JOSIP H., ZORAN P., LIDIJA G., JASMINKA A. Assessment of the natural and anthropogenic sources of chemical elements in alluvial soils from the Drava River using multivariate statistical methods. *Journal of Geochemical Exploration*, **110** (3), 278, **2011**.
 100. LI S., SUN W., CHEN R., ZHANG Z., NING D., NI Z. A historical record of trace metal deposition in northeastern Qinghai-Tibetan Plateau for the last two centuries. *Environmental Science and Pollution Research*, **29** (17), 24716, **2021**.
 101. WU J., DUAN D., LU JI., LUO Y., BRIAN J.B. Inorganic Pollution around the Qinghai-Tibet Plateau: An Overview of the Current Observations. *Science of the Total Environment*, **550**, 628, **2016**.

- 1 LIU H., LIU E., YU Z., LIN Q., ZHANG E., SHEN J. Spatio-temporal accumulation patterns of trace metals in sediments of a large plateau lake (Erhai) in Southwest China and their relationship with human activities over the past century. *Journal of Geochemical Exploration*, **234**, 106943, **2022**.
- 2 VIANELLO P., ANSORGE I., ROUAULT M., OSTROWSKI M. Transport and transformation of surface water masses across the Mascarene Plateau during the Northeast Monsoon season. *African Journal of Marine Science*, **39** (4), 453, **2017**.
- 3 LIU J., GUO H. Hydrochemical Characteristics and Ion Source Analysis of the Yarlung Tsangpo River Basin. *Water*, **15** (3), 537, **2023**.
- 4 HUANG J., LIU R., WANG Q., GAO X., HAN Z., GAO J., GAO H., ZHANG S., WANG J., ZHANG L., XIA X. Climate factors affect N₂O emissions by influencing the migration and transformation of nonpoint source nitrogen in an agricultural watershed. *Water Research*, **223**, 119028, **2022**.
- 5 JOEL D.B. The Effect of Late Cenozoic Glaciation and Tectonic Uplift on Silicate Weathering Rates and the Marine ⁸⁷Sr/⁸⁶Sr Record[A] // WILLIAM F.R. Tectonic Uplift and Climate Change[C]. Plenum Press: New York, pp. 260, **1997**.
- 6 XIAO D., SHAO X., QIN N., HUANG X. Tree-ring-based reconstruction of streamflow for the Zaqu River in the Lancang River source region, China, over the past 419 years. *International Journal of Biometeorology*, **61** (7), 1173, **2017**.
- 7 LI T. Statistical characteristics of crustal element abundance. *Geology and Exploration*, **10**, 7, **1992** [In Chinese].
- 8 LI T. Element abundances of China's continental crust and its sedimentary layer and upper continental crust. *Chinese Journal of Geochemistry*, **14** (1), 26, **1995**.
- 9 ROBERT S., JOSIP H., ZORAN P., LIDIJA G., JASMINKA A. Assessment of the natural and anthropogenic sources of chemical elements in alluvial soils from the Drava River using multivariate statistical methods. *Journal of Geochemical Exploration*, **110** (3), 278, **2011**.