

Original Research

Characterising Soil Ecological Stoichiometry in Different Vegetation in Southwest China Karst

Xianghuan Gong, Yingge Shu*, Changmin Wang, Hua Cai,
Xiulong Luo, Yuanhang Liao

College of Agriculture, Guizhou University, Guiyang 550025, China

Received: 03 March 2024

Accepted: 27 May 2024

Abstract

Investigate the nutrient content and ecological stoichiometry of Karst soils in southwest China to provide scientific evidence for nutrient cycling among Karst ecosystem components. The results showed that the mean values of soil SOC, TN, and TP were 26.68 (g Kg⁻¹), 2.34 (g Kg⁻¹), and 0.13 (g Kg⁻¹), respectively. The means for C:N, C:P, and N:P were 12.64, 211.56, and 18.78, respectively. Soil SMC, sand, and AN showed a highly significant positive correlation ($P < 0.01$) with SOC and TN, while BD and clay showed a highly significant negative correlation ($P < 0.001$) with SOC. TN content and storage were less influenced by vegetation type. Soil C:N values were relatively stable, while soil C:P and N:P values correlated with changes in SOC and TN. Soil carbon and nitrogen contents are higher under PF vegetation type than other vegetation types, so shrubs are more suitable for ecological restoration and environmental reconstruction in this area, and shrub vegetation type should be considered first for vegetation restoration in Karst areas in southwest China.

Keywords: Karst area, Vegetation, Soil stoichiometry, Soil nutrients

Introduction

Vegetation is an important part of terrestrial ecosystems and plays an important role in water conservation and environmental protection [1]. In recent years, the global climate is changeable, natural disasters occur frequently, unreasonable human activities occur, and vegetation degradation occurs locally or globally, resulting in a reduction of vegetation coverage, soil organic carbon (SOC), and total nitrogen (TN) content [2, 3]. Therefore, carrying out regional vegetation related research and investigation is the premise and foundation of natural ecosystem restoration, which is very effective in improving regional soil

quality and curbing the deterioration of the ecological environment. Ecological stoichiometry focuses on the balance and coupling of the main elements, such as carbon, nitrogen, and phosphorus, and organically unifies the research theories of different levels of molecules, cells, organisms, populations, and ecosystems based on the ratio between elements and elements [4–6]. Soil ecological stoichiometry provides a basis for understanding ecosystem processes, such as primary production and nutrient cycling, through the relative proportion and balance of C elements in soil, N, and P [7]. Soil C:N:P stoichiometry can represent the characteristics of nutrient cycling in different vegetation soil microenvironments [8]. C:N and C:P affect the growth rate of vegetation [9], which also reflects the utilization efficiency of vegetation on soils N and P. Previous studies have conducted a lot of research on soil stoichiometry in the central Himalayas [10], the Loess Plateau of China [11], southwestern China [12],

*e-mail: ygshu@gzu.edu.cn

and South China [13], with few studies on Karst areas [14, 15]. Most of the existing studies have focused on the effects of different land use patterns [16–18], vegetation recovery patterns [19], and fertilization measures [20–22] on soil stoichiometric changes. Climate conditions, terrain, parent material, anthropogenic, and many other factors will affect the soil stoichiometry [2, 23–25]. For example, warming increases in soil N:P and vectorial angle (reflecting P limitation by soil microbes) of 4.2% and 2.0%, respectively [26]. A large number of studies have shown that soil carbon, nitrogen, and phosphorus content and reserve changes respond to vegetation [19, 22], and different vegetation have found inconsistent C, N, and P content and their stoichiometry.

The ecosystem of the Karst region in southwest China is extremely fragile, with low disaster threshold elasticity and sensitivity that is susceptible to adverse factors. Vegetation restoration is an important ecological management measure in this region, and vegetation restoration is considered to be an effective measure to increase the retention of SOC [27, 28]. Vegetation species can affect the number, composition, and law of litter and ultimately affect soil carbon, nitrogen, and phosphorus [17, 19, 28].

At present, the research on soil stoichiometry in Karst areas mainly focuses on different vegetation types [19, 29] and restoration measures [15]. However, the C, N, and P content, reserves, and stoichiometric evolution of limestone soil in Karst areas are rarely reported. We hypothesized that the C, N, and P stoichiometric characteristics in Karst lime soil varied according to the vegetation. This paper verifies the hypothesis of different vegetation recovery types, recovery years (15a), terrain, and soil types and discusses the stoichiometry of soil C:N:P under different vegetation of adjacent agricultural land control (ZM), grassland (IC), shrub land (PF), and orchard (AP).

Experimental

Study Area

Pingba District, Anshun City, Guizhou Province, in the central part of Guizhou Province, Southwest China,

is the study area of this project (26°11'07"~26°36'56"N, 105°56'20"~106°39'10"E), and has a humid subtropical climate with an average annual rainfall of 1165 mm and a temperature range of 12.80 to 16.20°C. The mountain system is part of the Miaoling mountain range, which is mostly northeast-southwest oriented, non-parallel, and poorly continuous. The mountain system belongs to the Miaoling mountain range and is mostly north-east-south-west oriented, non-parallel, and poorly continuous. Most of the area is hilly, and the Karst landscape is very pronounced, with a thin layer of soil and a high proportion of gravel in the soil. The parent rocks of the area are mainly limestone and muddy gray rocks, and the soil type is predominantly calcareous. There are diverse vegetation types, dominated by scrub, arable, woodland, and grassland, with vegetation dominated by hawthorn, artemisia, raspberry, rowan, parkland, and white fescue.

Sample Sampling and Determination

Soils of three vegetation types (15a) were selected in the study area through a survey of the main vegetation types in the area in August 2022, identifying grassland (*Imperata cylindrical* (L.) Beauv., IC), shrubs (*Pyracantha fortuneana*, PF), and orchards (*Amygdalus persica* L., AP), with an adjacent agricultural field as a control (*Zea mays*, ZM) (Table 1). The selected sampling sites had the same time of vegetation restoration and geographic characteristics, including elevation, slope, and parent material, except for the different vegetation types. The vegetation types and years of restoration at the sample sites were determined by research and visits.

Soil sampling is to select a typical sample point according to the 0–5, 5–10, 10–20, 20–30, and 30–40 cm soil layers. Using the “S” shaped distribution points, each sample point repeated sampling 3 times. Roots, pebbles, and plant and animal remains visible to the naked eye were removed from the soil samples to ensure that the mass of each soil sample was ≥1 kg. After collection, soil samples were naturally air-dried to remove roots, stems, and pebbles. They were then ground and sieved through 2 mm mesh for indoor chemical analysis.

Table 1. Basic information on the sample plots in the southwest karst area

Vegetation types	Altitude (m)	Repair period (years)	Latitude and longitude	Primary Vegetation
ZM	1211	0	26°20'52"N 106°32'19"E	<i>Zea mays</i>
IC	1285	15	26°20'05"N 106°27'56"E	<i>Imperata cylindrical</i> (L.) Beauv.
PF	1243	15	26°19'15"N 106°28'41"E	<i>Pyracantha fortuneana</i> , <i>Artemisia annua</i> , <i>Rubus idaeus</i> L. and <i>Zanthoxylum simulans</i>
AP	1215	15	26°21'04"N 106°32'21"E	<i>Amygdalus persica</i> L.

Abbreviations: *Zea mays* (ZM), *Imperata cylindrical* (L.) Beauv. (IC), *Pyracantha fortuneana* (PF), *Amygdalus persica* L. (AP). The same is true below.

Soil organic carbon (SOC) was determined using external heating, total nitrogen (TN) using semi-micro Kjeldahl, total phosphorus (TP) using NaOH melting, alkaline nitrogen (AN) was determined by the alkaline diffusion method, and quick-acting phosphorus (AP) by the 0.5 mol L⁻¹ NaHCO₃ method [30], pH using potentiometry (water:soil = 2.5:1), and soil moisture content (SMC) using the desiccation method, Soil bulk density (BD) was determined by the ring knife method [31]. Soil texture was estimated using the hydrometer method [31].

Statistical Analysis

Soil SOC_{stock}(Cs), TN_{stock}(Ns), and TP_{stock}(Ps) (t hm⁻¹) were calculated [32, 33] as follows:

$$\text{Soil (Cs, Ns, Ps)}_i = \text{Soil (C,N,P)}_i \times \text{BD}_i \times \text{Di}/10$$

where C_i, N_i, and P_i are the SOC, TN, and TP contents in soil depth *i* (g kg⁻¹), BD_i is the BD of soil depth *i*, and Di is the soil depth of soil depth *i*.

Excel 2019, SPSS 22.0, and OriginPro 2021 software were used for data processing and plotting. One-way analysis of variance (ANOVA) with the LSD method was used to test the significance ($\alpha = 0.05$) of SOC, TN, and TP stock characteristics and soil stoichiometric ratios in the 0–40 cm range of the soil layer under different vegetation. The effects of vegetation type and soil depth on soil physico-chemical properties and on SOC, TN, and TP stocks and stoichiometric ratios were investigated using a two-way ANOVA.

Results and Discussion

Results

Different Vegetation Soil Profile Stoichiometry

The study found that the carbon, nitrogen, and phosphorus content of ZM, IC, PF, and AP soils all decreased with the increase in soil depth. According to Fig. 1, the content of carbon, nitrogen, and phosphorus in 0–10 cm in each vegetation is significantly higher than that at the bottom (10–20 cm, 20–30 cm, 30–40 cm), showing the surface aggregation phenomenon. PF soil SOC content peaked at 0–5 cm, and IC and PF soil SOC content were significantly greater than AP at 0–20 cm ($P < 0.05$). IC, PF, and AP soil TN content varied not significantly between different soil layers. The TP content of AP soil was significantly greater than IC and PF ($P < 0.05$). IC and PF soil C:P values decrease with the deepening of the soil layer and are significantly greater than the C:P values of ZM and AP. From 0–40 cm, C:N, N, P, C:N:ZM soil were not significantly different ($P > 0.05$). In the soil profile, the orders of SOC and TP content in different vegetation were PF > IC > AP > ZM, AP > ZM = PF > IC, and the difference in TN content was not significant ($P > 0.05$) (Table 2). The mean SOC content of IC, PF, and AP compared to ZM increased by 80.93%, 37.59%, and 101.34% at 0–40 cm soil depth, respectively.

Table 2. C, N, P content, ratio and stocks of 0–40cm soil layer in different vegetation(V).

V	SOC (g Kg ⁻¹)	TN (g Kg ⁻¹)	TP (g Kg ⁻¹)	C:N	C:P	N:P	Cs (Mg ha ⁻¹)	Ns (Mg ha ⁻¹)	Ps (Mg ha ⁻¹)
ZM	17.16±2.03b	2.08±0.44a	0.12±0.01ab	12.61±3.44a	138.44±18.28b	16.15±8.39a	17.32±6.37b	1.93±0.01a	0.11±0.04ab
IC	31.39±5.95a	2.38±0.31a	0.11±0.01b	14.02±6.07a	278.71±91.62a	21.79±7.86a	27.86±10.72a	2.12±0.79a	0.14±0.03a
PF	34.55±8.99a	2.49±0.39a	0.12±0.01ab	14.28±7.05a	267.29±105.37a	20.36±8.00a	27.47±39.14a	2.14±0.83a	0.10±0.04b
AP	23.61±1.54ab	2.45±0.14a	0.15±0.02a	9.64±0.50a	161.80±39.26b	16.83±4.14a	22.58±6.22ab	2.34±0.63a	0.13±0.04ab
Average value	26.68±4.14	2.35±1.05	0.13±0.04	12.64±8.18	211.56±94.85	18.78±7.50	23.81±9.20	2.13±0.82	0.12±0.04

Note: different lowercase letters in the same column indicate significant differences between different vegetation types ($P < 0.05$).




Fig. 1. Soil stoichiometry changes in different vegetation species

Note: Different capital letters indicate significant differences among different vegetation types at the same depth ($P < 0.05$). Different lowercase letters indicate significant differences among different soil depths under the same vegetation ($P < 0.05$). The same is true below. Abbreviations: soil organic carbon (SOC), total nitrogen (TN), total phosphorus (TP), SOC:TN ratio (C:N), SOC:TP ratio (C:P), and TN:TP ratio (N:P).

Stoichiometric Stocks of Different Vegetation Soil Profiles

The $\text{SOC}_{\text{Stock}}$, TN_{Stock} , and TP_{Stock} ranges of the soil profile depths in the study area were 12.65–45.40, 1.36–3.11, and 0.08–0.18 Mg ha^{-1} , respectively, and from 0–40 cm soil depth, their reserves were 131.88, 11.70, and 0.64 Mg ha^{-1} . IC, PF, and AP soil $\text{SOC}_{\text{Stock}}$ and TN_{Stock} were higher than those of ZM compared to the control. The C, N, and P reserves of IC, PF, and AP all peaked at 10–20 cm in the soil depth, and in terms of carbon accumulation capacity, the trend of soil $\text{SOC}_{\text{Stock}}$ was IC, PF > AP, with no significant difference in $\text{SOC}_{\text{Stock}}$ between IC and PF, making them important soil “carbon sinks” in the region”. From the effect of N fixation, the trend of TN_{Stock} of IC, PF, and AP was not significant. From the effect of P fixation, the trend of TP_{Stock} change was AP > PF > IC, and AP was a soil “phosphorus sink” in the region (Table 2, Fig. 2).

Soil Carbon, Nitrogen, and Phosphorus Relationships Among Different Vegetation

As shown in Table 3–3, by two-way ANOVA, vegetation type had a highly significant ($P < 0.001$) influence on soil SOC, TP, C:P, and $\text{SOC}_{\text{Stock}}$, and soil depth had a significant influence on SOC, TN, TP, C:P, $\text{SOC}_{\text{Stock}}$, TN_{Stock} , and TP_{Stock} indicators. Soil SOC, C:P, and $\text{SOC}_{\text{Stock}}$ were significantly ($P < 0.01$) influenced by the combination of vegetation type and soil depth. C:N and N:P are not affected by the interaction of vegetation, soil layer, or SD ($P > 0.05$). From 0 to 10 cm, the C:N ratio size is sorted as IC > PF > AP > ZM (Fig. 3a). The C of PF and IC:P values decreased with the deepening of the soil layer (Figures 3–3b). From 0–10 cm, the N:P value of IC, ZM, and PF is greater than the bottom layer (Fig. 3c). C:N for IC, ZM, and PF:P values decreased with the deepening of the soil layer, but the opposite was true for AP (Fig. 3d).




Fig. 2. Stoichiometric stocks of different vegetation soil profiles

Note: Different capital letters indicate significant differences among different vegetation types at the same depth ($P < 0.05$). Different lowercase letters indicate significant differences among different soil depths under the same vegetation ($P < 0.05$). The same is true below. Abbreviations: $\text{SOC}_{\text{Stock}}$ (C_s), TN_{Stock} (N_s), TP_{Stock} (P_s).

Table 3. ANOVA results of soil C:N-P stoichiometries influenced by restored vegetative (S), soil depth (D) and their interaction.

		SOC	TN	TP	C:N	C:P	N:P	SOCStock	TNStock	TPStock
F										
	S	22.15	0.59	7.66	1.10	53.99	2.20	13.43	0.69	6.92
	D	29.12	6.22	13.48	1.21	14.72	1.52	16.62	3.191	20.40
	S×D	5.06	0.53	0.88	1.29	9.20	1.07	2.98	0.893	0.73
P										
	S	< 0.001***	0.62	< 0.001***	0.36	< 0.001***	0.10	< 0.001***	0.565	< 0.01**
	D	< 0.001***	< 0.01**	< 0.001***	0.32	< 0.001***	0.22	< 0.001***	< 0.05*	< 0.001***
	S×D	< 0.001***	0.88	0.58	0.26	< 0.001***	0.41	< 0.01**	0.561	0.715

Abbreviations: Soil organic carbon (SOC), Total nitrogen (TN), Total potassium (TP), SOC:TN ratio (C:N), SOC:TP ratio (C:P) and TN:TP ratio (N:P). * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

As can be seen in Fig. 4, SOC showed a highly significant positive correlation ($P < 0.01$) with TN ($R^2 = 0.63$), TP ($R^2 = 0.45$), N:P ($R^2 = 0.38$), C:P ($R^2 = 0.84$), Cs ($R^2 = 0.57$), SMC ($R^2 = 0.37$), Sand ($R^2 = 0.64$), and AN ($R^2 = 0.93$), and highly significant negative correlation ($P < 0.01$) with BD ($R^2 = 0.61$), Clay ($R^2 = 0.56$) were highly significant negatively correlated ($P < 0.001$); and significantly negatively correlated with Ps ($R^2 = 0.26$); TN showed highly significant positive correlation ($P < 0.001$) with SOC ($R^2 = 0.63$), TP ($R^2 = 0.50$), N:P ($R^2 = 0.77$), C:P ($R^2 = 0.45$), Ns ($R^2 = 0.56$), AN ($R^2 = 0.62$); significant positive correlation ($P < 0.001$) with SMC ($R^2 = 0.30$), Sand ($R^2 = 0.31$), AP ($R^2 = 0.26$) were significantly positively correlated ($P < 0.05$); TP showed highly significant positive correlation ($P < 0.001$) with SOC ($R^2 = 0.45$), TN ($R^2 = 0.50$), AP ($R^2 = 0.80$), AN ($R^2 = 0.47$); and significant positive correlation ($P < 0.05$) with Ps ($R^2 = 0.28$).

C:N showed highly significant negative correlation ($P < 0.01$) with TN ($R^2 = 0.36$), N:P ($R^2 = 0.43$), Ns ($R^2 = 0.48$) and significant negative correlation ($P < 0.05$) with Clay ($R^2 = 0.29$); C:P showed highly significant positive correlation ($P < 0.001$) with SOC ($R^2 = 0.84$), TN ($R^2 = 0.45$), N:P ($R^2 = 0.57$), Cs ($R^2 = 0.56$), pH ($R^2 = 0.39$), SMC ($R^2 = 0.48$), AN ($R^2 = 0.79$), Sand ($R^2 = 0.67$), and highly significant positive correlation ($P < 0.001$) with BD ($R^2 = 0.53$), Ps ($R^2 = 0.46$), and Clay ($R^2 = 0.63$) were highly significant negatively correlated ($P < 0.001$); and with AP ($R^2 = 0.28$) were significantly negatively correlated ($P < 0.05$); N:P was significantly positively correlated ($P < 0.01$) with SOC ($R^2 = 0.38$), TN ($R^2 = 0.77$), C:P ($R^2 = 0.57$), Ns ($R^2 = 0.56$), SMC ($R^2 = 0.43$), AN ($R^2 = 0.38$); significantly positively correlated ($P < 0.05$) with Sand ($R^2 = 0.27$); significantly positively correlated ($P < 0.05$) with BD ($R^2 = 0.42$), Ps ($R^2 = 0.45$), C:N ($R^2 = 0.43$) were highly significant negative correlation ($P < 0.001$); and with Clay ($R^2 = 0.25$) were significant negative correlation ($P < 0.05$).

Discussion

Characteristics of C, N, and P Content and Stocks of Different Vegetation Soil Profiles

The average contents of SOC, TN, and TP in the 0–40 cm soil layer in the study area were 26.68 ± 4.14 , 2.35 ± 1.05 , and 0.13 ± 0.04 g kg⁻¹, respectively, which were comparable to the mean values of SOC, TN, and TP in terrestrial soils of China (29.51 , 2.30 , and 0.53 g kg⁻¹), and to the soils of the Karst region of southwestern China (6.44 – 45.17 g kg⁻¹, 0.56 – 2.57 g kg⁻¹, and 0.30 – 1.83 g kg⁻¹). TN, TP range values (6.44 – 45.17 g kg⁻¹, 0.56 – 2.57 g kg⁻¹, and 0.30 – 1.83 g kg⁻¹) were compared with the mean values of SOC, TN, and TP in Chinese terrestrial soils [34, 35]. Chinese terrestrial soil mean, SOC content was lower than the Chinese terrestrial soil mean in the Southwest China Karst area range value, and TP content was lower than the Chinese terrestrial soil mean and the Southwest China Karst area range value in the Southwest China Karst area range value.





Fig. 3. Change of soil stoichiometric ratio under different vegetation

Abbreviations: SOC:TN ratio (C:N), SOC:TP ratio (C:P) and TN:TP ratio (N:P).

In this study, the contents of SOC, TN, and TP decreased with the deepening of the soil profile, and the positive correlation between SOC, TN, TP, and AN was highly significant ($P < 0.001$), and the positive correlation between TN, TP, and AP was significant ($P < 0.05$), which indicated that the total amount and effectiveness of carbon, nitrogen, and phosphorus in the top 0–5 cm layer had the phenomenon of “surface aggregation”, which retarded the cycling of these elements to a certain extent. This indicates that the total amount and effectiveness of carbon, nitrogen, and phosphorus in the top 0–5 cm layer had the phenomenon of “surface aggregation”, which retarded the cycling of these elements to a certain extent. This may be due to the fact that plant and animal residues mainly accumulate in the soil surface layer, and thus higher biological activity in the surface soil may be a key factor contributing to this phenomenon [7]. In addition, the presence of surface

apoplastic litter can significantly improve soil structure and nutrient cycling [36].

In the study area, soil SOC content and storage between 0–20 cm soil layers were significantly higher in natural recovery (PF, IC) than in anthropogenic intervention (ZM, AP), on the one hand, because the vegetation was richer under natural recovery and the nutrients would be concentrated in the soil surface layer before being transported to the lower layer [37]. On the other hand, the upper layer of the soil had more microorganisms in the soil relative to the bottom layer, better aeration of soil structure, and faster nutrient cycling, which are conducive to the accumulation of organic carbon, suggesting that biological action has a more significant role in the regulation of SOC and TN [19]. In soils where ZM and AP are highly influenced by humans, the decrease in organic carbon may be due to the increase in soil permeability caused by agricultural



Note: * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$
Fig. 4. Correlation analysis between soil physicochemical properties and stoichiometric ratio

operations, which promotes soil SOC mineralization and release, leading to a decrease in soil carbon stock. The TN content in the study area showed $PF > AP > IC > ZM$, but there was no significant difference, indicating that vegetation restoration in the area had little effect on elemental N cycling. The TP content showed that AP was higher than PF, IC, and ZM, and there was no significant difference in the soil P content of each vegetation type among the soil layers except for the 0–5 and 5–10 cm soil layers, indicating that the P cycle was greatly affected by anthropogenic influences; however, in natural restoration, the soil P content of PF was higher than the soil P content of IC soil P content was higher, which may be related to the low bioavailability of soil P [38], suggesting that perennial shrub vegetation is more conducive to the biosmall cycle of P than annual herbaceous vegetation.

Ratios of Soil Profiles C, N, and P in Different Vegetation

The total stoichiometric mass ratio of soil carbon, nitrogen, and phosphorus can not only be used as a judgment indicator of soil nutrient supply, but also reflect soil quality and composition. Soil carbon and nitrogen are key elements required for vegetation growth, most of which exist in the form of organic matter, and their ratios are relatively stable due to strong coupling [39, 40]. Although organisms may not be able to directly regulate soil TP, TP can indirectly link SOC and TN by directly affecting the reactive phosphorus available for plant productivity and leading to strong coupling between SOC, TP, and TN [41]. However, due to the different turnover rates of SOC, TN, and TP along the soil profile, C:N, C:P, and N:P ratios may also increase with vertical depth, decreasing significantly, suggesting that

changes in SOC, TN, and TP did not vary proportionally across different soil horizons [19].

The average value of soil C:N in China is in the range of 10.00–12.00 [34], and the average value of soil C:N in the study area is 12.64, which is higher than the national average. The higher C:N indicates that the decomposition and mineralization rate of soil organic matter in the study area is slower, which is conducive to the accumulation of soil nutrients. Soil C:P is a sign of phosphorus mineralization capacity [42]. The mean value of soil C:P in this study area was 211.56, which was significantly higher than the average level in China (61) [34], and the C:P ratio differed significantly among vegetation types and depths, and the C:P ratio was greater than 200 and significantly higher than that of anthropogenic interventions (ZM, AP) in the profile (0–40 cm) and among soil layers (Fig. 1, Table 2–3), indicating that the phosphorus effectiveness in naturally restored (PF, IC) soils was lower, the mineralization rate was slower, and plant growth was restricted by phosphorus, which may be related to the fact that total phosphorus in Karst areas is not easy to convert into effective phosphorus [42]. N:P can reflect the decomposability of organic matter and determine the status of nutrient limitation in soils, and it is generally considered that N:P of less than 10 or greater than 20 is used as an indicator of the evaluation of vegetative productivity restricted by N or P limitation. Soil N:P (18.78) in the study area ranged from 10 to 20, indicating that the soil was limited by both nitrogen and phosphorus, natural restoration (PF, IC) soil N:P ratio > 20, the soil was limited by phosphorus and anthropogenic intervention (ZM, AP) soil N:P ratio ranged from 10 to 20, the soil was limited by both nitrogen and phosphorus.

The medium and primary productivity of most terrestrial ecosystems is constrained by TN or TP [43], and the stable functioning of these ecosystems relies on the availability of TN, TP to sustain soil SOC accumulation [7, 43, 44]. In agricultural systems, nitrogen and phosphorus deficiencies are usually addressed through the application of fertilizers as a way to maximize productivity. However, harvested agricultural products take away large amounts of nutrients, which need to be replenished through fertilizer application to prevent nutrient deficiencies over time. Unlike agriculture, large nutrient losses are not usually experienced under naturally restored vegetation because the amount of removal is small. As a result, soils under naturally restored vegetation are sufficient to maintain the nutrient levels they inherited from previous land uses. Nevertheless, the reduction of Ps in the naturally restored (PF, IC) substrate suggests that phosphorus may be a limiting factor for vegetation growth under naturally restored (PF, IC) conditions.

Factors Regulating the Ecological Stoichiometry of Different Vegetation Soils

Soil C content can be determined by the coefficients of humus and organic matter entering the soil, whereas N comes from biological nitrogen fixation or enters the soil through precipitation, and the main sources of P are

leaching and rock weathering [45, 46]. Similar to the results of many previous studies, SOC and TN had an extremely significant correlation because soil C and N are important components of soil organic matter, and SOC is the main source of N that affects the level of TN [47]. The significant correlation between SOC and TP was different from the results of Qiao et al. [48], which proved that SOC did not affect the spatial distribution. Notably, TN was also significantly correlated with TP ($P < 0.01$) and AP ($P < 0.05$). The C:P and N:P ratios were highly significantly positively correlated with a correlation coefficient of 0.57; similar to the relationship between SOC and TN, the C:P and C:N ratios were non-significantly correlated with a correlation coefficient of 0.23, as well as the C:N and N:P ratios were highly significantly negatively correlated with a correlation coefficient of 0.43. The correlation between C, N, and P stoichiometry values actually corresponds to the correlation between their nutrient contents.

Soil C, N, and P content, storage, and stoichiometry usually vary with vegetation type, soil depth, and soil properties [49]. Soil texture also affects the variation of C, N, and P content and their storage [49, 50]. In this study, soil SMC, Sand, and AN showed highly significant positive correlation ($P < 0.01$) with SOC and TN, and BD and Clay showed highly significant negative correlation ($P < 0.001$) with SOC, which was in agreement with the results of the previous study [46]. Soil capacity reflects the resistance, permeability, and aeration of the soil during root extension [51]. Soil water content can directly or indirectly affect soil nutrient content by altering ion transport, organic matter decomposition, and air heat transfer [52]. Soil C, N, and P stoichiometric ratios are affected by a combination of environmental factors such as soil-forming factors, soil water content, vegetation type, and human disturbance [53]. Therefore, the study of soil stoichiometric characteristics under different vegetation types under the same growing environmental conditions is of great significance in revealing the soil nutrient supply capacity under different vegetation types and the effects of different plant species on soil improvement.

Conclusions

The results of this study showed that vegetation type and soil depth had a significant effect on SOC and TP content and storage. SOC, TN, and TP content all decreased with increasing soil depth and had a significant surface aggregation effect. TN content and storage were less affected by vegetation type. Soil C:N values were relatively stable, while soil C:P and N:P values were correlated with changes in SOC and TN contents. Soil phosphorus effectiveness is lower, and vegetation restoration is more affected by phosphorus. Soil carbon and nitrogen contents are higher under the PF vegetation type than other vegetation types, so shrubs are more suitable for ecological restoration and environmental reconstruction in this area. Vegetation restoration in Karst areas of southwest China should first consider shrub vegetation types and pay attention to soil P

levels; soil texture, soil water content, and soil bulk weight should also be considered in the process of vegetation restoration with regard to the impacts of soil C, N, and P.

Acknowledgements

We are grateful to the National Natural Science Foundation of China (NSFC)(31460133).

Conflict of Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- BING K., SHI R.L., YUAN G.W., YUE J.Z., JIAN G.C. Population dynamics during succession of secondary natural forest in daqingshan, guangxi, china. *Journal of Plant Ecology*, **30** (6), 940, **2006**.
- LI Z., QIU X.R., SUN Y., LIU S., HU H., XIE J., CHEN G., XIAO Y., TANG Y., TU L. C:N:P stoichiometry responses to 10 years of nitrogen addition differ across soil components and plant organs in a subtropical *Pleuroblastus amarus* forest. *The Science of the total environment*, **796**, 148925, **2021**.
- AIHUA H., XIAN X., FEI P., QUANGANG Y., JIE L., HANCHEN D., CUIHUA H., SIYANG D. Different vegetation and soil degradation characteristics of a typical grassland in the Qinghai-Tibetan Plateau. *Acta Ecologica Sinica*, **40** (3), 975, **2022**.
- WANG S.Q., YU G.R. Ecological stoichiometry characteristics of ecosystem carbon, nitrogen and phosphorus elements. *Acta Ecologica Sinica*, **28** (8), 3947, **2008**.
- HAN X.G. Ecological stoichiometry:searching for unifying principles from individuals to ecosystems. *Chinese Journal of Plant Ecology*, **34** (1), 6, **2010** [In Chinese].
- DONGPING Z., LILING J., CONGSHENG Z., WEIQI W., CHUN W. Reviews on the ecological stoichiometry characteristics and its applications. *Acta Ecologica Sinica*, **33**, 5492, **2013**.
- ZHANG Y., XU X., LI Z., LIU M., XU C., ZHANG R., LUO W. Effects of vegetation restoration on soil quality in degraded Karst landscapes of southwest China. *The Science of the total environment*, **650**, Pt 2, 2665, **2019**.
- YANG Y., LUO Y. Carbon:nitrogen stoichiometry in forest ecosystems during stand development. *Global Ecology and Biogeography*, **20**, 361, **2011**.
- GREN G.I. The C:N:P stoichiometry of autotrophs—theory and observations. *Ecology Letters*, **7** (3), 191, **2004**.
- JOSHI R.K., GARTOKT S.C. Influence of vegetation types on soil physical and chemical properties, microbial biomass and stoichiometry in the central Himalaya. *CATENA*, **222**, 106835, **2023**.
- WANG L., ZHANG G., ZHU P., XING S., WANG C. Soil C, N and P contents and their stoichiometry as affected by typical plant communities on steep gully slopes of the Loess Plateau, China. *CATENA*, **208**, 105740, **2022**.
- WEI S., DING S., LIN H., LI Y., ZHANG E., LIU T., DUAN X. Microbial and enzymatic C:N:P stoichiometry are affected by soil C:N in the forest ecosystems in southwestern China. *Geoderma*, **443**, 116819, **2024**.
- TANG X., HU J., LU Y., QIU J., DONG Y., LI B. Soil C, N, P stocks and stoichiometry as related to land use types and erosion conditions in lateritic red soil region, south China. *CATENA*, **210**, 105888, **2022**.
- PANG D., CUI M., LIU Y., WANG G., CAO J., WANG X., DAN X., ZHOU J. Responses of soil labile organic carbon fractions and stocks to different vegetation restoration strategies in degraded Karst ecosystems of southwest China. *Ecological Engineering*, **138**, 402, **2019**.
- WEN L., LI D., XIAO K., TANG H., XIAO X., LI C. Difference in total N and its aggregate-associated N following cropland restoration in a Karst region, Southwest China. *Environmental Science and Pollution Research*, **30**, 50771, **2023**.
- SU L., DU H., ZENG F., PENG W., RIZWAN M., NUNEZ D.A., ZHOU Y., SONG T., WANG H. Soil and fine roots ecological stoichiometry in different vegetation restoration stages in a Karst area, southwest China. *Journal of environmental management*, **252**, 109694, **2019**.
- BAI Y., CHEN S., SHI S., QI M., LIU X., WANG H., WANG Y., JIANG C. Effects of different management approaches on the stoichiometric characteristics of soil C, N, and P in a mature Chinese fir plantation. *The Science of the total environment*, **723**, 137868, **2020**.
- KIM D., KIRSCHBAUM M.U., EICHLER L.B., GIFFORD R.M., LIANG L.L. The effect of land-use change on soil C, N, P, and their stoichiometries:A global synthesis. *Agriculture, Ecosystems Environment*, **348**, 108402, **2023**.
- CHEN M., YANG X., SHAO M., WEI X., LI T. Changes in soil C-N-P stoichiometry after 20 years of typical artificial vegetation restoration in semiarid continental climate zones. *The Science of the total environment*, **852**, 158380, **2022**.
- YAO X., HUI D., XING S., ZHANG Q., CHEN J., LI Z., XU Y., DENG Q. Mixed plantations with N-fixing tree species maintain ecosystem C:N:P stoichiometry:Implication for sustainable production. *Soil Biology and Biochemistry*, **191**, 109356, **2024**.
- CHEN G., YUAN J., WANG S., LIANG Y., WANG D., ZHU Y., WANG Y. Soil and microbial C:N:P stoichiometries play vital roles in regulating P transformation in agricultural ecosystems:A review. *Pedosphere*, **34** (1), 51, **2024**.
- ALAVAISHA E., MANZONI S., LINDBORG R. Different agricultural practices affect soil carbon, nitrogen and phosphorus in Kilombero -Tanzania. *Journal of environmental management*, **234**, 166, **2019**.
- CHENG R., WANG N., XIAO W., SHEN Y., LIU Z. Advances in Studies of Ecological Stoichiometry of Terrestrial Ecosystems. *entia silvae sinicae*, **54** (07), 136, **2018**.
- LI Y., ZHAO Y., BAO X., XIE H., LU X., FU Y., TANG S., GE C., LIANG C. Soil total and available C:N:P stoichiometry among different parent material soil profiles in rubber plantations of Hainan Island, China. *Geoderma Regional*, **36**, e00765, **2024**.
- CHEN X., FENG J., DING Z., TANG M., ZHU B. Changes in soil total, microbial and enzymatic C-N-P contents and stoichiometry with depth and latitude in forest ecosystems. *The Science of the total environment*, **816**, 151583, **2021**.
- XU H., WANG M., YOU C., TAN B., XU L., LI H., ZHANG L., WANG L., LIU S., HOU G. Warming effects on C:N:P stoichiometry and nutrient limitation in terrestrial ecosystems. *Soil and Tillage Research*, **235**, 105896, **2024**.

27. XIAO K., HE T., CHEN H., PENG W., SONG T., WANG K., LI D. Impacts of vegetation restoration strategies on soil organic carbon and nitrogen dynamics in a Karst area, southwest China. *Ecological Engineering*, **101**, 254, **2017**.
28. CAI H., SHU Y., WANG C., LIAO Y., LUO X., LONG H., LI X. Evolution Characteristics of Soil Active Organic Carbon and Carbon Pool Management Index Under Vegetation Restoration in Karst Area. *Huan jing ke xue = Huanjing kexue*, **44** (12), 6893, **2023** [In Chinese].
29. YANG T., ZHANG H., ZHENG C., WU X., ZHAO Y., LI X., LIU H., DONG L., LU Z., ZHOU J. Bacteria life-history strategies and the linkage of soil C-N-P stoichiometry to microbial resource limitation differed in Karst and non-Karst plantation forests in southwest China. *CATENA*, **231**, 107341, **2023**.
30. BAO S.D. Analytical Methods of soil and Agro-chemistry. Chinese Agriculture press, Beijing, China, 3, **2000** [In Chinese].
31. ZHANG G.L., GONG Z.T. Soil Survey Laboratory Methods. Science Press, Beijing, China, 25, **2012** [In Chinese].
32. DENG L., HAN Q., ZHANG C., TANG Z., SHANG-GUAN Z. Above-Ground and Below-Ground Ecosystem Biomass Accumulation and Carbon Sequestration with *Caragana korshinskii* Kom Plantation Development. *Land Degradation and Development*, **28**, 917, **2017**.
33. ZHANG P., LIU G., YU Y. Ecological Stoichiometry of Soil Carbon, Nitrogen and Phosphorus in Reclaimed Farmland in Coal Mining Subsidence Area. *Journal of Soil Science and Plant Nutrition*, **23**, 2511, **2023**.
34. TIAN H., CHEN G., ZHANG C., MELILLO J.M., HALL C.A.S. Pattern and variation of C:N:P ratios in China's soils: a synthesis of observational data. *Biogeochemistry*, **98** (3), 139, **2010**.
35. ZHENG X., SHENG M., ZHANG Y., GONG Z., WANG L. PhytOC sequestration characteristics and phytolith carbon sink potential of Karst Masson pine forest in southern China. *Science of The Total Environment*, **913**, 169688, **2024**.
36. ANGST Š., MUELLER C.W., CAJTHAML T., ANGST G., LHOTÁKOVÁ Z., BARTUŠKA M., ŠPALDOŇOVÁ A., FROUZ J. Stabilization of soil organic matter by earthworms is connected with physical protection rather than with chemical changes of organic matter. *Geoderma*, **289**, 29, **2017**.
37. WANG Y.D., WEI J.S., ZHOU M. Soil stoichiometric characteristics in the poplar and birch secondary forests in southern GreaterXing'an Mountains. *Chinese Journal of Soil Science*, **51** (5), 1056, **2020**.
38. RODRIGUES M., WITHERS P.J.A., SOLTANGHEISI A., VARGAS V., HOLZSCHUH M., PAVINATO P.S. Tillage systems and cover crops affecting soil phosphorus bioavailability in Brazilian Cerrado Oxisols. *Soil & Tillage Research*, **205** (1), 104, **2021**.
39. YANG Y., FANG J., JI C., DATTA A., LI P., MA W., MO-HAAMMAT A., SHEN H., HU H., KNAPP B.O., SMITH P. Stoichiometric shifts in surface soils over broad geographical scales: evidence from China's grasslands. *Global Ecology and Biogeography*, **23**, 955, **2014**.
40. CLEVELAND C.C., LIPTZIN D. C:N:P stoichiometry in soil: is there a "Redfield ratio" for the microbial biomass? *Biogeochemistry*, **85**, 252, **2007**.
41. SABINE GUSEWELL. N:P ratios in terrestrial plants: variation and functional significance. *New Phytologist*, **164** (2), 266, **2010**.
42. ÅGREN G.I. Stoichiometry and Nutrition of Plant Growth in Natural Communities. *Annual Review of Ecology, Evolution, and Systematics*, **39**, 153, **2008**.
43. DU C., WANG X., ZHANG M., JING J., GAO Y. Effects of elevated CO₂ on plant C-N-P stoichiometry in terrestrial ecosystems: A meta-analysis. *Science of The Total Environment*, **650**, 697, **2019**.
44. MA W., LI J., GAO Y., XING F., SUN S., ZHANG T., ZHU X., CHEN C., LI Z. Responses of soil extracellular enzyme activities and microbial community properties to interaction between nitrogen addition and increased precipitation in a semi-arid grassland ecosystem. *Science of The Total Environment*, **703**, 134691, **2020**.
45. XIAOHUI D., SHUZHENG L., JINWEI L., KUI L.I., GUOHUA L. Longitude gradient changes on plant community and soil stoichiometry characteristics of grassland in Hulunbeir. *Acta Ecologica Sinica*, **32** (11), 3467, **2012**.
46. WANG H., ZHANG G., LI N., ZHANG B., YANG H. Variation in soil erodibility under five typical land uses in a small watershed on the Loess Plateau, China. *CATENA*, **174**, 24, **2019**.
47. LIU J., WANG Y., LI Y., LIU X., JIANG Y., FU Y., JIN W., WU J. Ecosystem N:P stoichiometric ratios determine the catchment surface water N:P ratio through subsurface hydrological processes. *CATENA*, **194**, 104740, **2020**.
48. QIAO J., ZHU Y., JIA X., HUANG L., SHAO M. Vertical distribution of soil total nitrogen and soil total phosphorus in the critical zone on the Loess Plateau, China. *CATENA*, **166**, 310, **2018**.
49. JIANG F., WU X., XIANG W., FANG X., ZENG Y., OUYANG S., LEI P., DENG X., PENG C. Spatial variations in soil organic carbon, nitrogen and phosphorus concentrations related to stand characteristics in subtropical areas. *Plant and Soil*, **413** (1), 289, **2017**.
50. LIU S., ZHANG W., WANG K., PAN F., YANG S., SHU S. Factors controlling accumulation of soil organic carbon along vegetation succession in a typical Karst region in Southwest China. *Science of The Total Environment*, **521**, 52, **2015**.
51. ZHENG Y.S., CHEN L.G., HONG W. Study on productivity and soil properties of mixed forests of Chinese fir and *Phyllostachys heterocycla* cv. *Pubescens*, **34**, 16, **1998**.
52. LUO G.H., MA F.L., FU S.H. Effects of Waterlogging on Soil Nutrient on the Gentle Slope Zone in the Black Soil Region of Northeast China. *Journal of Anhui Agricultural Sciences*, **43** (15), 95, **2015**.
53. ACHAT D.L., BAKKER M.R., ZELLER B., DERRIEN D., NIKITICH P. Phosphorus status of soils from contrasting forested ecosystems in southwestern Siberia: effects of microbiological and physicochemical properties. *Biogeosciences*, **10** (2), 733, **2013**.