

Original Research

Study on Genotype × Environment Interaction in Wheat (*Triticum aestivum* L.) Varieties under the Changing Climate of Eastern Sub-Himalayan Plains

**Biplab Mitra¹, Bidusi Tripathy¹, Suwendu Kumar Roy², Srinivasan Vishnupriya²,
Saikat Das², Manoj Kanti Debnath³, Ahmed Gaber⁴, Akbar Hossain^{5*}**

¹Department of Agronomy, Uttar Bangla Krishi Viswavidyalaya, Pundibari, Coochbehar-736165, West Bengal, India

²Department of Genetics and Plant Breeding, Uttar Bangla Krishi Viswavidyalaya, Pundibari,
Coochbehar-736165, West Bengal, India

³Department of Agricultural Statistics, Uttar Bangla Krishi Viswavidyalaya, Pundibari,
Coochbehar-736165, West Bengal, India

⁴Department of Biology, College of Science, Taif University, P.O. Box 11099, Taif 21944, Saudi Arabia

⁵Division of Soil Science, Bangladesh Wheat and Maize Research Institute, Dinajpur 5200, Bangladesh

Received: 16 April 2024

Accepted: 12 June 2024

Abstract

The study of genotype x environment (G x E) interaction plays a crucial role in the context of changing climate scenarios. This field experiment was conducted during *rabi* (winter) seasons of two consecutive years, i.e., 2017-18 and 2018-19, to identify the potential wheat varieties having greater yield stability over varying sowing dates under eastern sub-Himalayan plains of India. The experiment was laid out in a split-plot design having four different dates of sowing (November 5, November 25, December 15, and January 5) in main plots with six wheat genotypes (HS 562, HD 2967, HD 3086, HI 1544, MACS 6222, and WR 522) in sub-plots, each replicated three times. Observations such as plant height (cm), days to physiological maturity, spikes m⁻², grains spike⁻¹, 1000 seed weight (g), biomass yield (q ha⁻¹), and grain yield (q ha⁻¹) were recorded. The genotypic coefficient of variation (GCV) and phenotypic coefficient of variation (PCV) values of the characters were low to moderate. The association among the characters indicated improvement in most of the attributing characters except 1000 seed weight, and the path analysis revealed that the highest direct effect on grain yield was exhibited by spike m⁻² (1.216). The ranking of the genotypes based on the rescaled index value of the seven characters showed that the variety HD 2967 was the best and highest performer with rank 1, followed by MACS 6222 (Rank 2) and HI 1544 (Rank 3). However, stability analysis revealed that HI 1544 and MACS 6222 were the most stable for grain yield compared to other varieties. Based on the AMMI stability value

(ASV), yield stability index (YSI), and the Eberhart and Russel model (1966), HI 1544 (Rank 1) was the most stable for grain yield over the eight environments. The AMMI analysis revealed that the most stable varieties were HI 1544 and MACS 6222, and the highest yielding variety, HD 2967 (Rank 3) had a better YSI ranking but with lower buffering capacity, leading to a higher response to environmental fluctuations arising out of the varying sowing dates under the eastern sub-Himalayan plains of India.

Keywords: AMMI, G×E, rescaled index, stability, wheat

Introduction

Commonly known as the ‘King of Cereals’, wheat (*Triticum aestivum* L.) occupies the prime position among the food crops in the world. In India, it is the second most important food crop next to rice and contributes to the total food grain production of the country to the extent of about 25%. Currently, India is the second largest producer of wheat in the world after China, with around a 12% share in global wheat production. Indian wheat production in 2019-20 has made another landmark achievement by producing 107.18 million metric tons with an average national productivity of 3508 kg ha⁻¹. The past year of production was also more than 100 million tons (103.60 million tons), and the current year of production has witnessed a change of 3.58 million tons (+3.46%). The positive production growth is attributed to the increased area of 4.21% despite a fall in the crop yield marginally by 0.72%. An increase in the support price also might have had a positive impact on the crop acreage (+1.24 million hectares). The average national productivity has crossed 3.5 t ha⁻¹ for the first time in the country [1].

Production variability in cereal crops has been related to the availability of precipitation and temperatures during the growing season [2]. Crop productivity is primarily determined by a combination of temperature and precipitation because temperatures have to be in the optimum range for plant growth and precipitation has to supply crop water requirements for a given environment. The investigations have found decreased grain yield with exposure to high maximum temperatures, particularly at the later part of the crop growth. Temperatures impact crop phenology, and each species has a specific base temperature, an optimum temperature value, and an upper-temperature limit [3]. An increase in temperature above the optimum has shown a negative impact on wheat yield, with a projected 5.3% yield reduction per 1°C temperature rise [4]. High temperature following anthesis is called terminal heat, and continual stress is experienced by the crop when the mean daily temperature during pollination increases, and it has a significant impact on yield [5, 6]. Heat stress affects wheat production by reducing the tiller number, grain filling period, and grain size.

It is believed that wheat cultivation in India is a pure gamble of temperature. It has been presumed that increasing temperatures will become more

significant in affecting wheat productivity in the coming years. However, some of this impact can be offset by ensuring these crops have an adequate soil water supply. Recent investigations [7, 8] suggest more attention to the effects of temperature on wheat productivity and suggest analyses and stimulation models be utilized for evaluating the potential growing regions and productivity for wheat under future climate scenarios.

Maintaining the appropriate sowing time is one of the most important agronomic practices for getting optimum plant growth and yield of wheat under heat-stressed environment [9], as in recent years we have experienced more fluctuations in temperature. Selection of suitable cultivars or genotypes can maintain higher productivity of wheat in any region. In general, late-sown wheat varieties face severe temperature stress, shortened heading, and maturity duration, and ultimately affect final yield and grain quality [10, 11]. Under the changing climate perspective, it is the need of the era to optimize the sowing window, which may have a great role in better performance of the crop. The low productivity of wheat in the eastern plains of India is due to a shorter favorable growing period, as in recent years, the temperatures began to rise from mid-February, and thus a short cool spell during its growing season with more fluctuation in temperature hampers the crop performance. Thus, the time of sowing is an important factor, and it is a non-cash input for achieving maximum yield. Advancing sowing time for a particular variety may not be always feasible in a high rainfall zone like the eastern sub-Himalayan plains of the country (India), where there is plenty of moisture in the soil even during the harvesting of the preceding rice crop in early November. Due to high residual moisture, the wheat crop can be grown successfully with a lower number of irrigations [12]. From an experiment conducted in the sub-Himalayan plains of India, it has been reported [13] that wheat sown on 25th November achieved maximum yield over the other dates of sowing; December 15 sowing recorded an estimated overall yield reduction of 40.7 kg ha⁻¹ day⁻¹ over November 25, sowing and at the same time, November 5 sowing also resulted in a yield decline by 42.7 kg ha⁻¹ day⁻¹, indicating the optimum sowing window in late November for eastern sub-Himalayan plains.

The deviation from the recommended sowing time of a variety for a particular set of environments results in failure to achieve its maximum production capacity.

At the same time, we have to assess the suitability of varieties under various dates of sowing. The varieties performing better under timely sown conditions may not perform well under late or very late sown situations. Similarly, there are certain varieties that are well-responsive to restricted irrigation facilities. However, the variety having greater flexibility over the sowing dates would be of high priority, considering the practical situation of farming. Although the wheat varieties differ in maturity duration, still in the present study they have been considered together intentionally due to shifting of the climatic conditions of the zones in which the wheat varieties are recommended, under the present changing climatic conditions, since extremely low yields are predicted due to adverse weather conditions [14]. An increase in genetic diversity at the field level allows a greater resilience to climate variability [15]. A study of genetic variability for the characters showing responses to environmental conditions is a prerequisite for the adaptation of a population to an environment [16]. This adaptation is ensured by the presence or absence of genotype \times environment (G \times E) interaction, which largely determines the average response of the genotypes [17]. Temporal (over time in a single location) stability has been used along with spatial (over different locations) stability to understand the adaptation of the genotypes [16]. Some promising wheat varieties under timely sown irrigated, late, and very late sown, as well as rainfed conditions for this sub-Himalayan plain [18-20] have been identified previously. Both the macro and micro environmental factors have an important role to play in the study of the stability of wheat varieties in grain yield over varying environmental conditions. The macro environmental factors are uncontrollable and include the broad external conditions that affect plant breeding and agricultural practices like weather patterns, geography, soil type, and quality, pest and disease pressure, etc. On the other hand, the micro environmental factors are controllable and can be managed to some extent, like agricultural practices (sowing date), field management, microclimate (windbreaks, shelterbreaks, canopy management, etc.), soil microbiome (microbial diversity, organic matter), etc. The impact of macro and micro environments on a crop is best judged by studying the G \times E interaction and selecting stable genotypes across different environmental conditions created due to changing climatic scenarios across the globe. Therefore, taking all these aspects into account and keeping in view the role of proper sowing time, which is a micro environmental and controllable factor in wheat production, the present experiment has been planned to study the genetic variability along with the G \times E interaction of the wheat varieties and identify the potential varieties for this sub-Himalayan plain in India that have greater yield stability over varying sowing dates.

Materials and Methods

Experimental Site, Method, and Materials

The field experiment was conducted at the Instructional Farm of Uttar Banga Krishi Viswavidyalaya, Pundibari, Cooch Behar, located at 26°24'02.2" N latitude, 89°23'21.7" E longitude, and at an elevation of 43 meters above mean sea level (msl). It was carried out during the *rabi* (winter) seasons of two consecutive years, i.e., 2017-18 and 2018-19. The experimental site was bestowed with a sub-tropical humid climate. The soil of the experimental site was sandy loam in texture, acidic in reaction (pH 5.81), with an organic carbon content of 0.85%. The experimental soil was low in available nitrogen (238.9 kg ha⁻¹), high in available phosphorus (32.6 kg ha⁻¹), and medium in available potassium (146.8 kg ha⁻¹). The meteorological monthly mean data pertaining to the period of experimentation are given in Table S1.

The experiment was laid out in a split-plot design with 24 treatment combinations, each replicated three times. Four different dates of sowing November 5th (D1), November 25th (D2), December 15th (D3), and January 5th (D4) were randomly allotted in each main plot, while six different wheat genotypes (HS 562, HD 2967, HD 3086, HI 1544, MACS 6222, WR 522) were randomly allocated in sub-plots. The important features of the varieties are mentioned in Table 1.

Agronomic Management

The seeds were sown manually in line with a seed rate of 100 kg ha⁻¹ for D1 and D2, and 125 kg ha⁻¹ for D3 and D4 at a row spacing of 20 cm. As the later sowing dates mostly result in decreased individual plant growth and tiller production, the seed rates under the later dates (D3 and D4) were 25% higher to maintain optimum stand count. The fertilizers were applied as per the recommendations of the All India Coordinated Wheat and Barley Improvement Project (AICW & BIP) under the Indian Council of Agricultural Research (ICAR), i.e., 150-60-40 kg ha⁻¹ N-P₂O₅-K₂O, respectively. The entire dose of P and K was applied as basal along with one-third of N. The rest N was applied in two equal splits, once at 21 days after sowing (DAS) and the next one at 42 DAS. Considering the deficiency of micronutrients (B and Zn in particular) in this tract, B was applied twice @ 0.20% with Soluble B (20% B), once at 35-40 DAS and the next at 55-60 DAS. Zinc (Zn) was applied at 0.10% with B in the second spray in the form of Chelated Zn. In both years, 4 irrigations were given at crown root initiation (CRI), active tillering, booting, and milking stages. Check basin method of irrigation was followed, keeping the depth of irrigation at 5 cm measured through volume basis. Harvesting of the crop was done manually and yield was estimated on a net plot basis excluding the border rows. After harvesting,

Table 1. Description of the wheat varieties taken in the experiment.

Variety with pedigree	Year of release	Duration (days)	Characteristics
HS 562 (OASIS/ SKAUZ//4*BCN/3/2*PASTPOR) IARI RS, Shimla	2015	130-140	Suitable for both irrigated and rainfed conditions; genetic yield potential of 62 q ha ⁻¹ under irrigated conditions; field resistance to leaf and stripe rusts; and possesses good chapatti and bread-making qualities; the average yield for the rainfed condition is 36 q ha ⁻¹ and for the irrigated condition is 52 q ha ⁻¹ .
HD 2967 ALD/COC//URES/HD2160M /HD2278 IARI, New Delhi	2010	130-140	Suitable for timely sown irrigated conditions; average seed yield is NWPZ: 50.4 q ha ⁻¹ ; NEPZ: 44.4 q ha ⁻¹ ; the variety has wide adaptability; it carries diversified genes other than 1B/1R; it possesses very high adult plant resistance against most prevalent leaf rust disease as well as 78S84 and 46S119, two most virulent races of yellow rust disease.
HD 3086 DBW14/HD 2733//HUW 468 IARI, New Delhi	2014	120-125	Suitable for timely sown irrigated conditions with an average yield of 54.6 q ha ⁻¹ , potential yield of 71 q ha ⁻¹ ; resistant to yellow rust and brown rust; superior quality parameters with protein content (12.5%), a sedimentation value (45 ml), best Glu-1 Score (10/10), and good extraction tare (70.5%) required for superior bread making.
HI 1544 HINDI62/BOBWHITE/CPAN2099 IARI RS, Indore	2008	110-115	It is an early maturing, semi-dwarf bread wheat variety for cultivation under high fertility and timely sown conditions; it combines early maturity, high stable yield, and tolerance to terminal heat stress along with high resistance to stem and leaf rusts; an average yield of 51-55 q ha ⁻¹ ; and good chapatti making quality.
MACS 6222 HD2189*2 //MACS 2496 IARI, Pune	2010	125-130	Suitable for irrigated, timely sown conditions; Medium early maturing, semi-dwarf, bold seeded; The potential yield is 47.1 q ha ⁻¹ ; leaf and yellow rust resistant.
WR 544 KALYANSONA/HD 2204/DW 38 IARI, New Delhi	2005	110-115	Suitable for irrigated late and very late sown conditions; average seed yield is 37.3 q ha ⁻¹ ; possesses genes for thermos tolerance and leaf rust resistance; fits well in intensive cropping systems

the produce was threshed and grains were dried to record yield at 14% moisture.

Data Collection

At harvest, the plant height was measured with a meter scale from the base to the tip of the plant. Five plants in each plot were randomly marked for this purpose except the border rows, and then the mean values were calculated and expressed in cm.

The days to maturity were counted from the date of seeding to the dates when the flag leaf and spike turned yellow. During the maturity stage, the number of spikes per square meter was randomly collected with the help of quadrates from five different spots in each plot, and accordingly, the mean values were calculated. The same procedure was followed for estimating the number of grains spike⁻¹. Thousand-grain samples were collected separately from each plot after threshing and drying under the sun. After proper sun drying, the samples were put into a dryer (65-70°C) for 2-3 days to take the final weights, and then the weight was expressed in grams for one thousand grains.

The entire produce from each net plot of twelve square meters was harvested (skipping the border rows) with the help of a sickle close to the ground level and then bundled separately and allowed to sun dry to reduce

the moisture level. After proper sun drying, weight was taken, averaged, and converted into quintals per hectare. After the separation of grains, the straw obtained from each plot over the net plot used for grain yield estimation was properly dried under the sun, and then the weight of the entire biomass was recorded and expressed in terms of quintals ha⁻¹ by considering both grain and straw.

Genetic Analysis for the Different Breeding Parameters and Statistical Analysis

The general statistical procedure was followed according to the standard method proposed [21]. The analysis of variance (ANOVA) and broad sense heritability (h^2_b) were estimated from the combined data over two years (2017-18 and 2018-19). The phenotypic coefficient of variation (PCV) and the genotypic coefficient of variation (GCV) were estimated according to the standard procedures [22]. The expected genetic advance (as a percentage of the mean) and the genotypic correlation were calculated [23]. The path analysis was carried out by the standard method [24]. The yield data was analyzed on an individual year basis and a combined basis, and the four different dates of sowing (D1, D2, D3, and D4) in the two years were treated to be a total of eight different environments (E1 to E8), considering the macro and micro environmental situations. So E1

was D1 in the first year of 2017-18, E2 was D2 in the first year of 2017-18, E3 was D3 in the first year of 2017-18, E4 was D4 in the first year of 2017-18, E5 was D1 in second year 2018-19, E6 was D2 in second year 2018-19, E7 was D3 in second year 2018-19, and E8 was D4 in second year 2018-19. The yield data was also used to estimate the different stability parameters as follows:

Regression-Based Stability Model

The stability model [25] deals with the regression approach to study the $G \times E$ interaction, and therein the interaction sum of squares is partitioned into two components. Among these two components, one component describes the heterogeneity of linear regression (b_i), while the other component illustrates combined deviations from individual regression lines (S^2_{di}). The b_i and the S^2_{di} are calculated as:

$$b_i = 1 + \frac{\sum (X_{ij} - \bar{X}_i - \bar{X}_j + \bar{X}_{..})(\bar{X}_j - \bar{X}_{..})}{\sum_{ij} (\bar{X}_j - \bar{X}_{..})^2}$$

$$S^2_{di} = \frac{1}{E-2} \left[\sum_i (X_{ij} - \bar{X}_i - \bar{X}_j + \bar{X}_{..}) - (b_i - 2)^2 \sum_j (\bar{X}_j - \bar{X}_{..})^2 \right]$$

Where X_{ij} is the performance of the i^{th} genotype in the j^{th} environment j , \bar{X}_i is the mean performance of the environment i , \bar{X}_j is the mean performance of the environment j , $\bar{X}_{..}$ is the grand mean, and E is the number of environments. Depending upon the value of b_i , the adaptability of the genotypes changes, i.e., when $b_i > 1$, the genotypes would be adapted to favorable environments, when $b_i < 1$, the genotypes would be adapted to unfavorable environmental conditions, and genotypes with $b_i = 1$ would have an average adaptation to all environments. Genotypes with $S^2_{di} = 0$ would be most stable, whereas if $S^2_{di} > 0$, it would indicate that the genotypes have lower stability across the environments. Overall, a genotype is said to be stable if the following conditions are fulfilled:

- i. The mean of the genotypes is greater than the population mean.
- ii. $b_i = 1$
- iii. $S^2_{di} = 0$

AMMI Stability Value (ASV)

In the AMMI model [26], the ASV is the difference between the coordinate point and the origin in a two-dimensional scatter diagram representing IPCA1 scores against IPCA2 scores. Because the IPCA1 score has more contribution to the $G \times E$ interaction sum of squares, a weighted value is needed. This weight is calculated for each genotype and each environment according to the relative contribution of IPCA1 to IPCA2 to the interaction sum of squares as follows:

$$ASV_i = \sqrt{\left[\frac{SS_{IPCA1}}{SS_{IPCA2}} (IPCA1Score) \right]^2 + (IPCA2Score)^2}$$

Where $\frac{SS_{IPCA1}}{SS_{IPCA2}}$ is the weight given to the IPCA1 value by dividing the IPCA1 sum of squares by the IPCA2 sum of squares. The larger the IPCA score, either negative or positive, the more specifically adapted a genotype is to certain environments. Smaller IPCA scores indicate a more stable genotype across environments.

Yield Stability Index (YSI)

For the selection of a genotype, stability should not be the only parameter, as the most stable genotype might not necessarily give the best yield performance [27, 28]. Therefore, there is a requirement for integrating both mean yield and stability into a single selection index. So, the various authors and scientists proposed different selection criteria for the simultaneous selection of yield and stability [29-31]. In this context, the rank of ASV and the rank of the mean performance of a character are considered. The lowest ASV value occupies rank one, while the highest mean value of a character occupies rank one, and then both ranks are summed into a single selection index of stability, called the yield stability index (YSI) and that is considered the most stable as well as the highest yielding genotype.

Statistical Analysis

The experimental data was subjected to statistical analysis using Windowstat (version-8.5), Cropstat (version 7.2), and OPSTAT.

Results

Variability Among the Wheat Genotypes over the Different Environments

The combined analysis of variance (Table 2) revealed that the six wheat genotypes differed significantly for grain yield and its attributing characters. The environments and the $G \times E$ interaction component also differed significantly for all the characters under study, indicating that the stability analysis of the wheat varieties can be further carried out in this study.

Genetic Variability

The study of genetic parameters (Table 3) showed wide variation in the range for all the characters. The grain yield ranged from 7.20 to 56.10 q ha⁻¹ among the genotypes, which showed a large variation. The GCV (4.89 to 13.88) and PCV (4.89 to 14.61) values of the characters were low to moderate. The heritability values of the characters were quite high. The genetic advance values ranged from low to high.

Table 2. Combined analysis of variance for the mean data of wheat genotypes over eight environments.

Sources of variation	DF	Mean sum of squares						
		Plant height (cm)	Days to physiological maturity	Spikes/m ²	Grains /spike	1000 seed weight (g)	Biomass yield (q/ha)	Grain yield (q/ha)
Genotypes (G)	5	145.426**	422.134**	3264.190**	38.144**	31.371**	578.539**	123.671**
Environment (E)	7	642.119**	609.328**	14254.996**	100.854**	96.851**	3492.826**	707.148**
G × E Interaction	35	13.160**	9.814**	891.784**	10.168**	6.008**	129.180**	24.013**
Combined Error	96	2.757	0.370	112.235	2.725	0.243	15.587	2.883

** Significant at 1% probability level

Table 3. Genetic parameters (combined over eight environments) for the different yield-attributing characteristics of wheat.

Characters	Mean	Range	GCV	PCV	Heritability (Broad Sense)	GA (as % of mean)
Plant height (cm)	86.643	61.10-110.20	4.891	4.982	0.964	9.893
Days to physiological maturity	111.003	84-130.00	6.541	6.550	0.997	13.458
Spikes/m ²	208.419	106.00-345.0	9.563	9.945	0.925	18.941
Grains /spike	36.889	21.40-51.90	5.663	6.397	0.784	10.328
1000 seed weight (g)	36.397	26.90-44.20	5.425	5.474	0.982	11.074
Biomass yield (q/ha)	72.750	24.90-131.20	11.374	12.295	0.856	21.675
Grain yield (q/ha)	27.832	7.20-56.10	13.882	14.611	0.903	27.170

Association between the Different Characters Indicating Their Influence on the Grain Yield

The genotypic correlation (Table S2) between grain yield and its attributing characters indicated that plant height was positively associated with days to physiological maturity (0.890**), spike per square meter (0.888**), grains per ear head (0.802*), biomass yield (0.765*), and grain yield (0.798*). The days to physiological maturity were positively associated with spike per square meter (0.872*), grains per ear head (0.943**), and grain yield (0.764*). Again, spike per square meter was positively associated with grains per spike (0.829*), biomass yield (0.994**), and grain yield (0.992**). Grains per spike were also positively associated with biomass yield (0.773*) and grain yield (0.840*). The character's 1000-seed weight was not associated with any of the characters. Biomass yield was found to be positively associated with most of the characteristics like plant height (0.765*), spikes/m² (0.992**), grains/spike (0.840*), and grain yield (0.993**). In totality, grain yield was positively correlated with all the characters except 1000 seed weight.

Path Analysis for the Relationship between the Different Characters and Grain Yield

The path analysis (Table 4) measures the depth of the relationship between the different characters, which

revealed that the highest direct effect on yield was exhibited by spike per square meter (1.216), which was closely followed by biomass yield (0.998) and grains per spike (0.633). These characteristics were also positively associated with grain yield, which indicated that any improvement in these three traits would result in a direct improvement in grain yield.

The correlation between plant height and grain yield (0.798) was significant and much higher than the direct effect of plant height (0.109) due to a negative indirect effect of plant height via days to physiological maturity. The correlation between days to physiological maturity and grain yield (0.764) was much higher than the direct effect of days to physiological maturity due to the indirect negative effect of days to physiological maturity via plant height. The correlation between spike per square meter and grain yield (0.992) was significant but lower than the direct effect of spike per square meter due to the negative indirect effect of spike per meter via plant height, days to physiological maturity, and 1000-seed weight. The correlation between grains per spike and grain yield (0.840) was higher than the direct effect of grains per spike (0.633), due to the greater negative indirect effect of grains per spike via days to physiological maturity (-1.547) and plant height (-0.088).

Table 4. Direct (diagonal) and indirect (off-diagonal) effects (combined over eight environments) of different yield components on grain yield in wheat.

Characters	Plant height (cm)	Days to physiological maturity	Spikes/m ²	Grains/spike	1000 seed weight (g)	Biomass yield (q/ha)	Correlation with grain yield
Plant height (cm)	-0.109	-1.459	1.080	0.508	0.015	0.764	0.798*
Days to physiological maturity	-0.097	-1.639	1.060	0.597	0.122	0.721	0.764*
Spikes/m ²	-0.097	-1.429	1.216	0.525	-0.214	0.992	0.992**
Grains/spike	-0.088	-1.547	1.008	0.633	0.062	0.771	0.840*
1000 seed weight (g)	0.002	0.289	0.376	-0.057	-0.693	0.577	0.495
Biomass yield (q/ha)	-0.084	-1.184	1.195	0.469	-0.401	0.998	0.993**

* Significant at 5% probability level, ** Significant at 1% probability level, residual effect = 0.2853

Table 5. ANOVA for stability of the character grain yield of wheat over eight environments as per Eberhart and Russel (1966).

Sources of variation	Degrees of freedom	Mean sum of squares
Replications within Environment	16	7.123
Genotypes (G)	5	123.674**
Environments (E) + (G×E)	42	137.869**
Environments (E)	7	707.150**
Genotype × Environment (G×E)	35	24.013**
Environments (Linear)	1	4950.051**
G×E (Linear)	5	18.158**
Combined Deviation	36	20.824**
Combined Error	80	2.883
Total	47	136.359

** Significant at 1% probability level

Ranking Based on Rescaled Index Value to Classify the Wheat Genotypes

The ranking of the genotypes (Table S3) based on the rescaled index value of the seven characters showed that the variety HD 2967 was the best performer with rank 1, followed by MACS 6222 (Rank 2), HI 1544 (Rank 3), HD 3086 (Rank 4), HS 562 (Rank 5), and WR 544 (Rank 6).

Stability Study Using Genotype × Environment Interaction

The ANOVA for stability (Table 5) as per Eberhart and Russel [25] for grain yield showed significance for genotypes and different components of the environment like [E+(G×E)], E, E (linear)] and the combined deviation, indicating a substantial difference between the varieties under study and the different environments created due to different dates of sowings and years.

The linear component of G×E differed significantly, which indicated that the performance of the wheat varieties could be predicted across the environments, created by the different dates of sowing and years, and the use of the stability parameters, i.e., b_i (regression coefficient) and S_{di}^2 (mean square deviation from linear regression), would be justified.

Eberhart and Russel (1966), AMMI and GGE Biplot Analysis

Based on mean grain yield over the environments (E1 to E8), regression coefficient (b_i), and mean square deviation from linear regression (S_{di}^2) as per Eberhart and Russel [25] as mentioned in Table 6, it is clear that only HI 1544 showed non-significant b_i and S_{di}^2 , but its mean (27.221q ha⁻¹) was lower than the population mean (27.831q ha⁻¹). The other five varieties exhibited significant S_{di}^2 along with non-significant b_i , which did not give a clear indication regarding the stability of the varieties.

Table 6. Mean grain yield of wheat genotypes in the individual eight environments (E₁ to E₈) and the stability parameters as per Eberhart and Russel (1966).

Genotypes	E1	E2	E3	E4	E5	E6	E7	E8	Mean	b _i	S ² _{di}
HS 562	28.033 ^{ab}	39.433 ^b	21.333 ^b	11.833 ^{ab}	38.400 ^{ab}	36.400 ^{bc}	26.767 ^b	15.033 ^{ab}	27.154	0.884 ⁴	18.844 ^{4**}
HD 2967	29.833 ^a	51.700 ^a	28.400 ^a	15.867 ^{ab}	44.067 ^a	47.933 ^a	35.800 ^a	18.800 ^{ab}	34.050	1.108 ³	31.154 ^{5**}
HD 3086	19.467 ^{cd}	24.700 ^d	19.300 ^{bc}	9.800 ^b	40.467 ^{ab}	45.667 ^a	36.933 ^a	14.500 ^{ab}	26.354	1.162 ⁵	10.415 ^{3**}
HI1544	22.133 ^{bc}	27.233 ^{cd}	21.300 ^b	10.300 ^{ab}	44.200 ^a	41.400 ^{ab}	34.667 ^a	16.533 ^{ab}	27.221	1.082 ²	3.191 ¹
MACS 6222	28.767 ^{ab}	34.200 ^{bc}	25.233 ^{ab}	17.300 ^b	42.867 ^a	44.900 ^a	32.700 ^{ab}	13.667 ^b	29.954	0.992 ¹	5.185 ^{2*}
WR 544	12.833 ^d	17.633 ^e	13.200 ^c	11.533 ^{ab}	34.600 ^b	34.000 ^c	33.300 ^{ab}	20.933 ^a	22.254	0.772 ⁶	34.614 ^{6**}
Population Mean	23.511	32.483	21.461	12.772	40.767	41.717	33.361	16.578	27.831		
SEm (±)	1.233	1.309	1.252	0.618	1.959	2.486	1.825	2.132	1.725		
Environmental Index	-4.320	4.652	-6.370	-15.059	12.935	13.885	5.530	-11.253			
CV (%)	9.082	6.981	10.100	8.376	8.322	10.321	9.473	22.266			
CD (P≤0.05)	3.885	4.125	3.943	1.946	6.172	7.833	5.749	-			

Four different dates of sowing; November 5th (D1), November 25th (D2), December 15th (D3), and January 5th (D4). These four different dates of sowing (D1, D2, D3, and D4) in the two years were treated to be a total of eight different environments (E1 to E8), considering the macro and micro-environmental situations. So E1 was D1 in the first year of 2017-18, E2 was D2 in the first year of 2017-18, E3 was D3 in the first year of 2017-18, E4 was D4 in the first year of 2017-18, E5 was D1 in second year 2018-19, E6 was D2 in second year 2018-19, E7 was D3 in second year 2018-19, and E8 was D4 in second year 2018-19. *Significant at 5% probability level, ** Significant at 1% probability level; Values in superscript in the format 1-6 indicate the respective ranks; Values in superscript in the format of a, b, ab, etc. indicate the DMRT values.

The heat map (Fig. 1) indicated the variable performance of the six wheat varieties over the eight environments created by the different dates of sowing over two years. The lighter shades in the heat map indicated higher yield and dark shades indicated lower yield. The stability analysis of the wheat varieties was further extended to AMMI and GGE Biplot analysis because the latter two techniques have additional advantages in providing more information and allowing greater discrimination among the varieties and their relationships between the environments [32]. The GGE biplot for discrimination and representativeness (Fig. 2) represents all the varieties in five environments, such as E3, E4, E5, E7, and E8, whereas the environments E1, E2, and E6 are highly differing and do not have much interaction with genotypes. The AMMI analysis of variance (Table S4) revealed that the replications, genotypes (varieties), environments, and G×E interactions component differed significantly for grain yield, which partially matched with the ANOVA of Eberhart and Russel model [25].

The AMMI analysis of variance (Table S4) for grain yield of the six wheat varieties showed highly significant differences among the genotypes and environments. The AMMI Stability Value (ASV) is calculated based on AMMI model IPCA1 and IPCA2 (interaction principal component 1 and 2, respectively) scores for each genotype. The varieties having the lowest ASV are considered widely adopted. The YSI (Yield Stability Index) is a more efficient indicator of stability as it

combines high-yield performance along with stability. Lower YSI values indicate high yield along with greater stability (28, 30, and 31). Based on the ASV in the present study (Table 7), HI 1544 (Rank 1) was the most stable over the eight environments, followed by HS 562 (Rank 2) and HD 3086 (Rank 3). Based on YSI, it was found that HI 1544 (Rank 1) was the most stable and high-yielding variety, followed by HS 562 (Rank 2) and HD 2967 (Rank 3).

Ranking of Wheat Genotypes Based on Grain Yield Deviation in Different Environments

The study was further extended to the ranking of the wheat varieties based on the negative grain yield deviation from the mean in the eight environments (Table 8). An interesting observation here was that the most stable variety, HI 1544 (-38.618), was ranked fourth, and the other stable variety, MACS 6222 (-34.849), was ranked second in the list of total negative deviations of the six wheat varieties.

The GGE-biplot for the two principal components (Fig. 3) represented 91.23% of the variability. The environment and varieties were placed in all four quadrants (I, II, III, and IV). In quadrant I, the best performers were HD 2967 and MACS 6222, which were good in E3, E4, E5, and E6. In quadrant II, the best performers were HI 1544 and HD 3086 for E7. In quadrant III, the best performer was WR544 for E8 with less interaction. In quadrant IV, the highest yielder




Fig. 1. Heat map for the performance of the different wheat genotypes for grain yield in the different environments. Note: Four different dates of sowing: November 5th (D1), November 25th (D2), December 15th (D3), and January 5th (D4). These four different dates of sowing (D1, D2, D3, and D4) in the two years were treated to be a total of eight different environments (E1 to E8), considering the macro and micro environmental situations. So E1 was D1 in the first year of 2017-18, E2 was D2 in the first year of 2017-18, E3 was D3 in the first year of 2017-18, E4 was D4 in the first year of 2017-18, E5 was D1 in the second year 2018-19, E6 was D2 in the second year 2018-19, E7 was D3 in the second year 2018-19, and E8 was D4 in the second year 2018-19.




Fig. 2. Discrimination versus representativeness. Note: Four different dates of sowing: November 5th (D1), November 25th (D2), December 15th (D3), and January 5th (D4). These four different dates of sowing (D1, D2, D3, and D4) in the two years were treated to be a total of eight different environments (E1 to E8), considering the macro and micro environmental situations. So E1 was D1 in the first year of 2017-18, E2 was D2 in the first year of 2017-18, E3 was D3 in the first year of 2017-18, E4 was D4 in the first year of 2017-18, E5 was D1 in the second year 2018-19, E6 was D2 in the second year 2018-19, E7 was D3 in the second year 2018-19, and E8 was D4 in the second year 2018-19.

Table 7. Ranking of six wheat genotypes based on grain yield, AMMI stability value (ASV), and yield stability index (YSI).

Genotypes	Grain Yield (GY)	RANK (GY)	IPCA1	IPCA2	AMMI Stability Value (ASV)	RANK (ASV)	Yield Stability Index (YSI)	RANK (YSI)
HS562	27.154	4	0.392	3.610	3.792	2	6	2
HD2967	34.050	1	3.519	-0.299	10.445	6	7	3
HD3086	26.354	5	-1.679	-0.845	5.051	3	8	4
HI1544	27.221	3	-0.725	-0.843	2.310	1	4	1
MACS6222	29.954	2	1.710	-1.545	5.304	4	6	2
WR544	22.254	6	-3.219	-0.077	9.548	5	11	5

was HS562 for E1 and E2 with very little interaction. The environment closest to the origin was E4, followed by E8, E3, and E1, which implied that these environments were unfavorable for grain yield. The ranking of the wheat varieties by mean yield and stability is represented in Fig. 4. It revealed that the genotypes HD 2967 and MACS 6222 were placed closer to the origin, and it indicated a higher mean yield. The stability line (Y axis) indicated that HD1644, HD 3086,


and HS 562 were the stable varieties. Since none of the varieties were placed away from the origin, it can be concluded that there was no greater G×E interaction and reduced stability.

In the What-Won-Where biplot (Fig. 5), the four varieties HD 3086, WR 544, HS 562, and HD 2967 were placed at the vertices of a quadrilateral and the two environments (E4 and E8) in the three sectors. In sector III, the variety WR544 was the best performer for E8.

Table 8. Ranking of six wheat genotypes based on grain yield deviation from the mean in the eight environments.

Genotypes	Grain yield (GY) (q/ha)	E1	E2	E3	E4	E5	E6	E7	E8	Negative deviation (%)
HS562	27.154 ⁴	0.879	12.279	-5.821	-15.321	11.246	9.246	-0.387	-12.121	-33.650 ¹
HD2967	34.050 ¹	-4.217	17.650	-5.650	-18.183	10.017	13.883	1.750	-15.250	-43.300 ⁵
HD3086	26.354 ⁵	-6.887	-1.654	-7.054	-16.554	14.113	19.313	10.579	-11.854	-44.003 ⁶
HI1544	27.221 ³	-5.088	0.012	-5.921	-16.921	16.979	14.179	7.446	-10.688	-38.618 ⁴
MACS6222	29.954 ²	-1.187	4.246	-4.721	-12.654	12.913	14.946	2.746	-16.287	-34.849 ³
WR544	22.254 ⁶	-9.421	-4.621	-9.054	-10.721	12.346	11.746	11.046	-1.321	-35.138 ³
Environmental mean deviation		-25.921	27.912	-38.221	-90.354	77.614	83.313	33.180	-67.521	
Environmental Index		-4.320	4.652	-6.370	-15.059	12.936	13.886	5.530	-11.254	

Values in superscript indicate the rank within the varieties for the negative deviations in yield. Four different dates of sowing: November 5th (D1), November 25th (D2), December 15th (D3), and January 5th (D4). These four different dates of sowing (D1, D2, D3, and D4) in the two years were treated to be a total of eight different environments (E1 to E8), considering the macro and micro environmental situations. So E1 was D1 in the first year of 2017-18, E2 was D2 in the first year of 2017-18, E3 was D3 in the first year of 2017-18, E4 was D4 in the first year of 2017-18, E5 was D1 in the second year 2018-19, E6 was D2 in the second year 2018-19, E7 was D3 in the second year 2018-19, and E8 was D4 in the second year 2018-19.



GGE Biplot showing components 1 and 2 explaining 91.23% of the total variation using Column Metric Preserving SVP and Tester-Centered G+GE with no scaling

Fig. 3. GGE Biplot for PC1 and PC2. Note: Four different dates of sowing: November 5th (D1), November 25th (D2), December 15th (D3), and January 5th (D4). These four different dates of sowing (D1, D2, D3, and D4) in the two years were treated to be a total of eight different environments (E1 to E8), considering the macro and micro environmental situations. So E1 was D1 in the first year of 2017-18, E2 was D2 in the first year of 2017-18, E3 was D3 in the first year of 2017-18, E4 was D4 in the first year of 2017-18, E5 was D1 in the second year 2018-19, E6 was D2 in the second year 2018-19, E7 was D3 in the second year 2018-19, and E8 was D4 in the second year 2018-19.




Fig. 4. GGE for Mean versus Stability. Note: Four different dates of sowing: November 5th (D1), November 25th (D2), December 15th (D3), and January 5th (D4). These four different dates of sowing (D1, D2, D3, and D4) in the two years were treated to be a total of eight different environments (E1 to E8), considering the macro and micro environmental situations. So E1 was D1 in the first year of 2017-18, E2 was D2 in the first year of 2017-18, E3 was D3 in the first year of 2017-18, E4 was D4 in the first year of 2017-18, E5 was D1 in the second year 2018-19, E6 was D2 in the second year 2018-19, E7 was D3 in the second year 2018-19, and E8 was D4 in the second year 2018-19.

HD 2967 from Sector I and HS 562 from Sector II were the best performers for E4. Sector II did not include any environment, and it can be concluded that varieties present in Sector II have no specific environment for recommendation.

Discussion

Variability and Association Study between the Different Characters

A similar result regarding the variation in wheat genotypes was reported [33] in the evaluation of 11 wheat genotypes under eight locations in the North Hill zone of India. The same was also reported [34] in the evaluation of five potato clonal hybrids under six locations in Bangladesh. A similar report was shown [35] in a work on twenty wheat genotypes under nine test environments in Egypt, to determine the abiotic stress tolerance capacity. This indicated that there was sufficient variability in the genotypes and the environments and the interaction between them.

The significance of the G×E interaction component indicated that the multi-environment performance of the genotypes can be further studied to assess their genetic potentiality for stability.

The GCV values were closer to the PCV values, indicating a lower effect of the environment on these characters. Similar views were opined [36] in a work on wheat genotypes for morpho-physiological characters. All the characters had high heritability, indicating the lower effect of the environment on these characters under study, and the GA was low to high for the different characters. The characters' biomass yield and grain yield exhibited high heritability and genetic advance, which indicated that they are controlled by additive genes and would respond to selection.

In the linear regression analysis, a similar result of a positive correlation of biomass with grain yield was reported [37], while evaluating the effects of organic and inorganic fertilizers on wheat with five combinations of N, P, manure, and compost treatments, and it was concluded that grain yield was most strongly correlated with total biomass. This is also supported by another report [38], in a study in Iran extending for a period




Fig. 5. Description of which won where what. Note: Four different dates of sowing: November 5th (D1), November 25th (D2), December 15th (D3), and January 5th (D4). These four different dates of sowing (D1, D2, D3, and D4) in the two years were treated to be a total of eight different environments (E1 to E8), considering the macro and micro environmental situations. So E1 was D1 in the first year of 2017-18, E2 was D2 in the first year of 2017-18, E3 was D3 in the first year of 2017-18, E4 was D4 in the first year of 2017-18, E5 was D1 in the second year 2018-19, E6 was D2 in the second year 2018-19, E7 was D3 in the second year 2018-19, and E8 was D4 in the second year 2018-19.

of several decades (from 1968 to 2018), involving 20 released spring wheat cultivars, which revealed that the wheat grain yield increased due to emphasis on traits like biomass, having a positive correlation with grain yield in combination with other agronomic traits. The association among the characters indicated that any improvement in most of the attributing characters except 1000-seed weight would result in an improvement of grain yield among the wheat genotypes under the present study.

Due to the correlation between plant height and grain yield being significant and higher than the direct effect of plant height, therefore, direct selection for plant height will not improve the grain yield. In this case, direct selection for a higher spike per square meter would result in improved grain yield due to its higher direct effect. Also, direct selection for higher days to physiological maturity will not improve grain yield due to its very high negative direct effect on grain yield. In the case of biomass yield, its correlation with grain yield (0.993) was nearly similar to the direct effect of biomass yield, which was due to the internal cancellation of the positive and negative indirect effect of biomass yield

via the other grain yield attributing characters. An interesting observation here was that plant height and days to physiological maturity created a negative indirect effect for the other grain yield attributing characters. On the other hand, spike per square meter and biomass yield created a positive indirect effect for the other grain yield attributing characters. The presence of a high residual effect in a population indicated the role of other possible independent variables that are not included in the study. In the present study, the residual effect of 0.2853 indicated that the characters presently studied fairly accounted for the total variability. The inclusion of a few more yield-attributing characters might have been better.

The inclusion of all the characters in the ranking (Table S3) of the varieties gave more clarity regarding their performance, not just based on grain yield, but also based on the grain yield attributing characters, and it helped to classify the wheat genotypes with HD 2967 with rank 1, followed by MACS 6222 (rank 2), and others. Interpretation of the results taking into consideration the ranking of the genotypes along with the stability parameters would throw more light on the

adaptation of the genotypes across the environments created by different dates of sowing and years.

Stability of the Wheat Genotypes across the Environments

Genotype \times environment (G \times E) interaction is the result of the differences in sensitivities of the genotypes to the prevailing conditions in the target environment [39]. This is very pertinent under the present prevailing climate change conditions and the changes in the dates of sowing in a crop like wheat where the source-sink relationship keeps shifting, depending on the climatic conditions. As per Eberhart and Russel [25], a variety is said to be stable only if $b_i=1$ and $S^2_{di}=0$ for that variety. Therefore, as per Table 6, strong conclusions could not be drawn regarding the stability of the varieties.

In the heat map (Fig. 1), the greater number of dark shades of HD 2967 in comparison to HI 1544 and MACS 6222 showed that HD 2967 suffered a greater degree of yield reduction with delay in sowing in the E4 and E8 (Table 8). However, HD 2967 was grouped along with the stable yielder MACS 6222 due to their higher average yields at rank 1 and 2, respectively, across the environments. Among the environments, it was found that those that were favorable (E2, E7, E5, and E6) with a positive environmental index were grouped into one, and those with a negative environmental index (E1, E3, E4, and E8) were grouped into another, thereby classifying the two distinct types of environments based on the sowing dates (Table 8). However, no specific trend in the yielding performance of the wheat varieties could be found, probably due to the variation in the responses of the varieties towards environmental fluctuations created by the changing dates of sowing over two years in combination with the changes in the weather parameter over the two years giving rise to complex interaction. Quite often the results of the Eberhart and Russel model [25] for stability do not match with those of AMMI and Biplot analysis due to the fact that AMMI and Biplot methods of stability analysis capture a much greater proportion of the genotype \times environment (G \times E) variance than the Eberhart and Russel method [40].

The AMMI model fits some of the several multiplicative forms instead of multiplicative forms in the assessment of varietal performance in varying environments [41]. The AMMI analysis (Table S4) determines the stability of the varieties across different environments using the PCA (Principal Component Analysis). Similar results of obtaining the lowest ASV with more stability were also reported [42] in fifteen rice genotypes under three different locations where the rice genotype SAHEL 108 (ASV = 0.05) was considered to be stable and possessed superior grain yield when compared to SAHEL 177 (ASV = 0.27). Similarly, in another evaluation of nineteen barley genotypes [43] under eight diverse locations, it was found that five barley genotypes KB 1405 (ASV = 0.33), BH 1013 (ASV = 0.57), BH 902 (ASV = 0.77), DWRB 150 (ASV = 1.14),

and DWRB 101 (ASV = 1.32) with low ASV are stable across the environments when compared to RD 2941 (ASV = 3.63). An investigation [34] with five advanced potato genotypes evaluated under six locations in Bangladesh also showed that the potato genotype clone 9.125 (ASV = 0.50), followed by BARI Alu-28 (ASV = 0.56) and clones 9.91 (ASV = 0.67) were the most stable when compared to the clone 9.35 (ASV = 4.61), which is undesirable. In contrast to the result, it was found that the best-yielding genotype was the least stable, and the most stable genotype was placed in the lowest rank in spring wheat and cotton, respectively, which indicates that highly stable genotypes need not be high-yielding genotypes [44, 45].

A similar result of obtaining low YSI (Table 7) with high grain yield was also reported [46] in an experiment involving twenty-two advanced breeding lines of rice in twelve different salt-affected environments, where the rice genotypes STBN 22 (YSI = 0.989) and STBN 24 (YSI = 0.997) exhibited the highest mean grain yield with low yield stability index. Similarly, it has been reported [47] that the wheat genotypes with low YSI exhibit high grain yield while investigating 20 winter wheat genotypes under 24 environments in Iran. The Eberhart and Russell model [25] and the AMMI analysis revealed that the most stable varieties were HI 1544 and MACS 6222. The highest-yielding variety, HD 2967 (Rank 3) performed substantially better for stability as per the YSI ranking.

The two stable varieties, namely HI 1544 and MACS 6222, had exhibited low G \times E interaction (Table 8) or a lower response to the environmental fluctuations arising out of different dates of sowing, starting from early (5th Nov.) to timely (25th Nov.) and then to late (15th Dec.) and very late (5th January) sowing, indicating a greater buffering capacity in them. On the other hand, the highest average yielding wheat variety, HD 2967 (34.05 q/ha), ranked 5th with a very high total negative deviation (-43.300), indicating a greater response to environmental fluctuations arising out of low buffering capacity.

The importance of the present study of well-known wheat varieties is enhanced by the fact that the best and the highest yielding variety HD 2967 in the first year exhibited the highest drop in grain yield from 51.700 q/ha in E2 (Nov. 25th sowing), which is a favorable environment, to 28.400 q ha⁻¹ in E3 (Dec. 15th sowing) and the lowest 15.867 q ha⁻¹ in E4 (Jan 5th sowing). Similarly, in the 2nd year, HD 2967 showed a steep drop from 47.933 q ha⁻¹ in E6 (25th Nov. sowing) to 18.800 q ha⁻¹ in E8 (5th Jan sowing), where E6 was the favorable environment of timely sown condition and E8 was the unfavorable late sown condition. Similar observations in the stable varieties HI 1544 and MACS 6222 did show a drop in the grain yield in E4 and E8 in the first and second years, but they were comparatively lower than the highest-yielding HD 2967. Another important observation was that in the most unfavorable environments E4 (Environmental

index = -15.059) and E8 (Environmental index = -11.254), the negative deviation in yield was very high in the highest-yielding HD 2967 and quite low in the stable varieties HI 1544 and MACS 6222. This finding is in contrast to the finding from Table S3, where HD 2967 ranked first, followed by MACS 6222 (rank 2) and HI 1544 (rank 3), due to the masking effect of yield depreciation under unfavorable environments by the extremely high yield of HD 2967 under favorable environments. Hence, blind use of high-yielding wheat varieties over a large area with changing environmental and climatic conditions is not desirable. In wheat cultivation, any change in the sowing date or sowing window period for the sowing can have a devastating effect on grain yield as the yield reduction is to about 2/3rd with a realization of only 1/3rd of the potential yield in very high-yielding varieties with low buffering capacity, showing higher responsiveness to environmental fluctuations. This is extremely relevant to the present climate change scenario across the globe, where the climatic factors fluctuate every year as observed in the present study (Table S1) for temperature, relative humidity, total rainfall, and the number of rainy days.

The findings of some of the environments being unfavorable for grain yield (Fig. 3) are supported by the environmental index of the respective environments, E4 (-15.059), E8 (-11.254), E3 (-6.370), and E1 (-4320), which clearly states that the delayed sowing on January 5th of both years (E4 and E8), delayed sowing on Dec. 15th (E3), and early sowing (E1) in the first year caused a greater reduction in grain yield. According to Eberhart and Russel [25], a lower and negative environmental index for any environment means an unfavorable environment. On the other hand, the environments E6, E7, and E5 were away from the origin, indicating that they were favorable environments for grain yield. This is also supported by the higher environmental index in E6 (November 25th sowing in 2nd year), E7 (December 15th sowing in 2nd year), and E5 (November 5th sowing in 2nd year) with considerably high environmental mean deviation (Table 8).

Conclusions

The present two-year trial was successful in properly discriminating between the wheat varieties on the basis of their variable performance due to a change in the date of sowing, which is a micro environmental and controllable factor, by identifying the G×E interaction component under the different environments created. The use of different dates of sowing over two years and their treatment as different environments is fully justified by the wide range of environmental indexes of the different environments created, each of which also behaved differently in a favorable and unfavorable way with respect to grain yield. The present study revealed that the varieties HD 2967 and MACS 6222 were the superior performing ones for grain yield, out

of which HD 2967 was a very high-yielding variety and performed well in E3, E4, and E5 environments but was also highly responsive to environmental fluctuations arising due to changes in dates of sowing. On the other hand, HI 1544 and MAC 6222 were identified as highly stable varieties performing well in almost all eight environments, with the least deviation in yield under unfavorable environments. Therefore, exploiting these varieties under the sub-Himalayan plains of India would ensure a high and stable grain yield.

Author Contribution

Conception: Biplab Mitra, Bidusi Tripathy, Suwendu Kumar Roy, S. Vishnupriya, Saikat Das, Manoj Kanti Debnath; Methodology: Biplab Mitra, Bidusi Tripathy, Suwendu Kumar Roy, S. Vishnupriya, Saikat Das, Manoj Kanti Debnath; Software, data curation and analysis: Biplab Mitra, Ahmed Gaber and Akbar Hossain; Preparation of all Tables and Figures: Biplab Mitra, Ahmed Gaber and Akbar Hossain; Preparation of the article: Biplab Mitra, Bidusi Tripathy, Suwendu Kumar Roy, S. Vishnupriya, Saikat Das, Manoj Kanti Debnath; Review and editing: Ahmed Gaber and Akbar Hossain; Supervision: Biplab Mitra; Project administrator: Biplab Mitra, Ahmed Gaber; Funding: Biplab Mitra, Ahmed Gaber, Akbar Hossain. All authors approved to submit the review in the journal.

Funding

The study was financially supported by Uttar Banga Krishi Viswavidyalaya, West Bengal, India. This research was also funded by Taif University, Saudi Arabia, Project No. (TU-DSPP-2024-07).

Acknowledgments

All authors are thankful to Uttar Banga Krishi Viswavidyalaya for giving support during drafting and finalization of the interesting review article. The authors extend their appreciation to Taif University, Saudi Arabia, for supporting this work through project number (TU-DSPP-2024-07).

Conflict of Interest

The authors have declared that no competing interests exist.

Informed Consent

Informed consent was obtained from all individual participants included in the study. Additional informed

consent was obtained from all individual participants for whom identifying information is included in this article.

Ethical Approval

This article does not contain any studies with human participants or animals performed by any of the authors.

Data Availability Statement

All data are fully available without restriction. All relevant data are within the manuscript and its Supporting Information files.

References

1. Directors Report, ICAR-AICW&BIP. Published by Indian Institute of Wheat and Barley Research (Indian Council of Agricultural Research), Karnal, Haryana, India, **2022**.
2. YU Q., LI L., LUO Q., EAMUS D., XU S., CHEN C., WANG E., LIU J., NIELSON D.C. Year patterns of climate impact on wheat yields. *International Journal of Climatology*, **34**, 518, **2014**.
3. HATFIELD J.L., BOOTE K.J., KIMBALL B.A., ZISKA L.H., IZAURRALDE R.C., ORT D., THOMSON A.M., WOLFE D. Climate impacts on agriculture: Implications for crop production. *Agronomy Journal*, **103**, 351, **2011**.
4. INNES P.J., TAN D.K.Y., VAN F., AMTHOR J.S. Effects of high temperature episodes on wheat yields in New South Wales, Australia. *Agricultural and Forest Meteorology*, **208**, 95, **2015**.
5. PRASAD P.V.V., DJANAGUIRAMAN M. Response of floret fertility and individual grain weight of wheat to high temperature stress: sensitive stages and thresholds for temperature and duration. *Functional Plant Biology*, **41**, 1261, **2014**.
6. REZAEI E.E., SEIBERT S., EWERT F. Intensity of heat stress in winter wheat-phenology compensates for the adverse effect of global warming. *Environmental Research Letters*, **10**, 024012, **2015**.
7. KARIMI T., STOCKLE C.O., HIGGINGS S.S., NELSON T.L., HUGGINGS D. Projected dryland cropping system shifts in the Pacific Northwest in response to climate change. *Frontiers in Ecology and Evolution*, **5**, 20, **2017**.
8. KAUR R., KAUR S., SANDHU S.S. Effect of terminal heat stress in wheat and its management with chemicals in North-West India. *Agricultural Research Journal*, **54**, 114, **2017**.
9. KAJLA M., YADAV V.K., KHOKHAR J., SINGH S., CHHOKAR R.S., MEENA R.P., SHARMA R.K. Increase in wheat production through management of abiotic stresses: a review. *Journal of Applied and Natural Science*, **7** (2), 1070, **2015**.
10. HOSSAIN A., TEIXEIRA DA SILVA J.A. Phenology, Growth and Yield of Three Wheat (*Triticum aestivum* L.) Varieties as Affected by High Temperature Stress. *Notulae Scientia Biologicae*, **4** (3), 97, **2012**.
11. HAKIM M.A., HOSSAIN A., TEIXEIRA DA SILVA J.A., ZVOLINSKY V.P., KHAN M.M. Yield, protein and starch content of twenty wheat (*Triticum aestivum* L.) genotypes exposed to high temperature under late sowing conditions. *Journal of Scientific Research* **4** (2), 477, **2012**.
12. MITRA B., DAS S. Economization of irrigation schedule in wheat (*Triticum aestivum* L.) cultivation under sub-Himalayan plains of West Bengal. *Journal of Crop and Weed*, **11** (2), 122, **2015**.
13. TRIPATHY B., SINGHA ROY A.K., MITRA B. Influence of dates of sowing on phenology, growth and yield of diverse wheat varieties under changing climate in Eastern sub-Himalayan Indian plains. *Journal of Cereal Research*, **12**, 69, **2020**.
14. LIU B., MARTRE P., EWERT P., PORTER J.R., CHALLINOR A.J., MULLER C., RUANA A.C., WAHA K., THORBURN P.J., AGGARWAL P.K. Global wheat production with 1.5 and 2.0°C above pre-industrial warming. *Global Change Biology*, **25**, 1428, **2018**.
15. ALTIERI M.A., NICHOLLS C.I. The adaptation and mitigation potential of traditional agriculture in a changing climate. *Climate Change*, **140**, 33, **2013**.
16. FRANK G.V., RIVIERE P., PIN S., BALTASSAT R., BERTHELLOT J.F., CAIZERGUES F., DALMASSO C., GASCUEL J.S., HYACINTHE A., MERCIER F., MONTAZ H., RONO T. B., GOLDRINGER I. Genetic diversity and stability of performance of wheat population varieties developed by participatory breeding. *Sustainability*, **12**, 384, **2020**.
17. PANDEY V., SINGH S.K., KORADA M., SINGH D.K., KHAIRE A.R., HABDE S., MAJHI P.K. *Indian Journal of Agricultural Research*, **54**, 689, **2020**.
18. DAS S., MITRA B. Performance of different wheat genotypes under various levels of nitrogen in rainfed condition of terai region of West Bengal. *Journal of Crop and Weed*, **7**, 23, **2011**.
19. MITRA B., SAMAJDAR T. Evaluation of promising wheat varieties in Terai region of West Bengal through on farm trial. *Research on Crops*, **14**, 33, **2013**.
20. DAS S., MITRA B. Performance of wheat genotypes under late and very late sowing in sub-Himalayan plains of West Bengal. *Crop Research*, **46**, 32, **2013**.
21. STEEL R.G.D., TORRIE J.H. *Principles and Procedures of Statistics*. 2nd ed. New York: McGraw-Hill, **1980**.
22. BURTON G.W. Quantitative inheritance in grasses. *Proc. 6th Int. Grassland Congress*, **1**, 277, **1952**.
23. JOHNSON H.W., ROBINSON H.F., COMSTOCK R.E. Genotypic and phenotypic correlations in soybeans and their implications in selection. *Agronomy Journal*, **47**, 477, **1955**.
24. DEWEY D.R., LU K.H. A correlation and path coefficient analysis of components of crested wheat grass seed production. *Agronomy Journal*, **51**, 515, **1959**.
25. EBERHART S.A., RUSSELL W.A. Stability parameters for comparing varieties. *Crop Science*, **6**, 36, **1966**.
26. PURCHASE J.L., HATTING H., VAN DENVENTER C.S. Genotype × environment interaction of winter wheat (*Triticum aestivum* L.) in South Africa: II. Stability analysis of yield performance. *South African Journal of Plant and Soil*, **17**, 101, **2000**.
27. MOHAMMADI R., ABDULAH A., HAGHPARAST R., ARMION M. Interpreting genotype-environment interactions for durum wheat grain yields using non-parametric methods. *Euphytica*, **157**, 239, **2007**.
28. MOHAMMADI R., AMRI A. Comparison of parametric and non-parametric methods for selecting stable and adapted durum wheat genotypes in variable environments. *Euphytica*, **159**, 419, **2008**.

29. BAJPAI P.K., PRABHAKARAN V.T. A new procedure of simultaneous selection for high yielding and stable crop genotypes. *Indian Journal of Genetics and Plant Breeding*, **60**, 141, **2000**.
30. RAO A.R., PRABHAKARAN V.T. Use of AMMI in simultaneous selection of genotypes for yield and stability. *Journal of Indian Society of Agricultural Statistics*, **59**, 76, **2005**.
31. BABARMANZOOR A., TARIQ M.S., GHULAM A., MUHAMMAD A. Genotype × environment interaction for seed yield in Kabuli Chickpea (*Cicer arietinum* L.) genotypes developed through mutation breeding. *Pakistan Journal of Botany*, **41**, 1883, **2009**.
32. ALWALA S., KWOLEK T., MCPHERSON M., PELLOW J., MEYER D. A comprehensive comparison between Eberhart and Russell joint regression and GGE biplot analyses to identify stable and high yielding maize hybrids. *Field Crops Research*, **119**, 225, **2010**.
33. JEBERSON M.S., KANT L., KISHORE N., RANA V., WALIA D.P., SINGH, D. AMMI and GGE biplot analysis of yield stability and adaptability of elite genotypes of bread wheat (*Triticum aestivum* L.) for Northern Hill Zone of India. *International Journal of Bioresource and Stress Management*, **8**, 635, **2017**.
34. KUNDU B.C., KAWOCHAR M.A., NAZMIN S., AHMED N.U., HALDER S.C., MOSTOFA M., DELOWAR H.K.M. Stability analysis for yield of advanced potato genotypes for commercial cultivation in Bangladesh, *SAARC Journal of Agriculture*, **18**, 73, **2020**.
35. AL-ASHKAR I., SALLAM M., ALMUTAIRI K.F., SHADY M., IBRAHIM A., ALGHAMDI S.S. Detection of high-performance wheat genotypes and genetic stability to determine complex interplay between genotypes and environments. *Agronomy*, **13**, 585, **2023**.
36. SINGH S.K., SINGH M.K., CHOUDHARY V.K., BARMAN M. Genetic variability analysis in bread wheat (*Triticum aestivum* L.) genotypes for morpho-physiological characters and grain micronutrient content. *International Research Journal of Pure and Applied Chemistry*, **21**, 1, **2022**.
37. AGE NEHU G., VANBEEK C., BIRD M.I. Influence of integrated soil fertility management in Wheat and tef productivity and soil chemical properties in the highland tropical environment. *Journal of Soil Science and Plant Nutrition*, **14**, 532, **2014**.
38. KHODARAHMI M., SOUGHI H., SHAHBAZI K., JAFARBY J., KHAVARINEJAD M.S. Trends in important agronomic traits, grain yield and its components in bread wheat cultivars released in northern warm and humid climate of Iran, 1968-2018. *Cereal Research Communications*, **51** (9), 1, **2023**.
39. FALCONER D.S., MACKAY T.F.C. *Introduction to Quantitative Genetics*. 4th Edition, Addison Wesley Longman, Harlow, **1996**.
40. NAMORATO H., MIRANDA G.V., VAGNODE SOUZA L., OLIVEIRA L.R., DELIMA R.O., MANTOVANI E.E. Comparing biplot multivariate analysis with Eberhart and Russell method for genotype × environment interaction. *Crop Breeding and Applied Biotechnology*, **9**, 299, **2009**.
41. BERNARDO A.B. Extending Hope Theory: Internal and External Locus of Trait Hope. *Personality and Individual Differences*, **49**, 944, **2010**.
42. ROY D., GAUR A., PANDEY I.D. Estimation of G × E interaction by AMMI model in 'Antenna Panel' genotypes of rice [*Oryza sativa* L]. *Brazilian Archives of Biology and Technology*, **65**, e22220202, **2022**.
43. KUMAR V., KHARUB A.S., SINGH G.P. Additive main effects and multiplicative interaction and yield stability index for genotype by environment analysis and wider adaptability in barley. *Cereal Research Communications*, **46**, 365, **2018**.
44. JĘDZURA S., BOCIANOWSKI J., MATYSIK P. The AMMI model application to analyze the genotype–environmental interaction of spring wheat grain yield for the breeding program purposes. *Cereal Research Communications*, **51**, 197, **2023**.
45. ABRO S., RIZWAN M., RAJPUT T.M., SIAL M.A., DEHO Z.A. Evaluation of upland cotton genotypes for stability over different locations using AMMI and GGE biplot analysis. *Pakistan Journal of Botany*, **54**, 1733, **2022**.
46. KRISHNAMURTHY S.L., SHARMA S.K., SHARMA D.K., SHARMA P.C., SINGH Y.P., MISHRA V.K., BURMAN D., MAJI B., BANDYOPADHYAY B.K., MANDAL S., SARANGI S.K., GAUTAM R.K., SINGH P.K., MANOHARA K.K., MARANDI B.C., SINGH D.P., PADMAVATHI G., VANVE P.B., PATIL K.D., THIRUMENI S., VERMA O.P., KHAN A.H., TIWARI S., SHAKILA M., ISMAIL A.M., GREGORIO G.B., SINGH R.K. Analysis of stability and G×E interaction of rice genotypes across saline and alkaline environments in India. *Cereal Research Communications*, **44**, 349, **2016**.
47. ROOSTAEIA M., MOHAMMADIB R., AMRI A. Rank Correlation among different statistical models in ranking of winter wheat genotypes. *Crop Journal*, **2**, 154, **2014**.

SUPPLEMENTARY MATERIAL

Table S1. Meteorological monthly mean data pertaining to the periods of experimentation.

Year	Month	Temperature (°C)		Relative Humidity (%)		Total Rainfall (mm)	No. of rainy days
		Maximum	Minimum	Maximum	Minimum		
2017-18	Nov.ember	29.55	16.31	95.03	57.39	1.0	1
	Dec.ember	27.27	13.41	97.08	56.45	0	0
	January	22.33	9.41	96.03	59.71	0	0
	February	25.1	11.19	86.96	53.50	0	0
	March	29.43	15.92	73.14	48.61	36	6
2018-19	April	30.67	18.84	72.07	58.42	10.18	4
	Nov.ember	28.56	15.2	88.67	54.67	0	0
	Dec.ember	26.36	11.52	93.86	56.45	1.16	1
	January	26.08	9.00	96.03	59.71	0	0
	February	26	11.55	87.39	53.5	2.23	1
	March	29.13	14.65	73.14	48.6	9.62	4
	April	30.94	19.8	73.25	59.74	34.34	8

Source: Gramin Krishi Mousam Seva Kendra, Pundibari, Coochbehar, West Bengal, India

Table S2. Genotypic correlation (pooled over eight environments) between grain yield and its attributing characters in wheat.

Characters	Days to physiological maturity	Spikes/m ²	Grains/spike	1000 seed weight (g)	Biomass yield (q/ha)	Grain yield (q/ha)
Plant height (cm)	0.890**	0.888**	0.802*	-0.022	0.765*	0.798*
Days to physiological maturity		0.872*	0.943**	-0.176	0.722	0.764*
Spikes/m ²			0.829*	0.309	0.994**	0.992**
Grains /spike				-0.090	0.773*	0.840*
1000 seed weight (g)					0.579	0.495
Biomass yield (q/ha)						0.993**

* Significant at 5% probability level, ** Significant at 1% probability level

Table S3. Mean performance of the wheat genotypes averaged over eight environments and their ranking on the basis of rescaled index value [(as suggested by Iyengar and Sudarshan (1982)] of plant height, days to physiological maturity, spikes/m², Grains/spike and 1000-seed weight.

Genotypes	Plant height (cm)		Days to physiological maturity		Spikes/m ²		Grains/spike		1000-seed weight (g)		Rescaled Index value (G) = $\frac{(X-\text{Min})}{(\text{Max}-\text{Min})}$	Total of the rescaled values (I) = (A + B + C + D + E + F + G)	Ranking on the basis of the total rescaled index value (I) *		
	X	Rescaled Index value (A) = $\frac{(X-\text{Min})}{(\text{Max}-\text{Min})}$	X	Rescaled Index value (B) = $\frac{(\text{Max}-X)}{(\text{Max}-\text{Min})}$	X	Rescaled Index value (C) = $\frac{(X-\text{Min})}{(\text{Max}-\text{Min})}$	X	Rescaled Index value (D) = $\frac{(X-\text{Min})}{(\text{Max}-\text{Min})}$	X	Rescaled Index value (E) = $\frac{(X-\text{Min})}{(\text{Max}-\text{Min})}$				X	Rescaled Index value (F) = $\frac{(X-\text{Min})}{(\text{Max}-\text{Min})}$
HS 562	90.360	0.773	118.127	0.048	216.500	0.661	37.647	0.598	33.497	0.000	70.517	0.422	27.157	2.919	5
HD 2967	92.997	1.000	118.920	0.000	236.837	1.000	40.037	1.000	37.580	0.770	85.360	1.000	34.050	5.770	1
HD 3086	85.930	0.391	109.297	0.579	199.047	0.369	35.840	0.295	36.853	0.633	70.777	0.432	26.353	3.069	4
HI1544	84.227	0.244	103.170	0.947	204.377	0.458	34.087	0.000	38.797	1.000	72.663	0.506	27.220	3.613	3
MACS 6222	84.943	0.306	114.213	0.283	216.837	0.666	38.343	0.715	37.090	0.678	77.510	0.694	29.957	4.007	2
WR 544	81.400	0.000	102.293	1.000	176.920	0.000	35.380	0.217	34.567	0.202	59.673	0.000	22.253	1.459	6

X = Mean performance of the character; *Ranking has been done in descending order taking the highest rescaled value as 1 (first rank) followed by other lower rescaled values from rank 2 onwards. The formulas for the index value (rescaled value) are the same for all the characters except in case of “Days to physiological maturity”, where lower values were more desirable and hence the rescaling formula was different.

Table S4. AMMI analysis of variance for grain yield of the six wheat varieties.

Sources of variation	Degrees of freedom	Mean sum of squares
Replication (within E)	16	21.37**
Genotypes (G)	5	371.02**
Environments (E)	7	2121.45**
G × E Interaction	35	72.04**
Error	80	8.65

** Significant at 1% probability level