

Original Research

# Micromorphological and Ultrastructural Exploration of *Sorbus foliolosa* (Rosaceae) Populations from the Himalayan Region of Kashmir

Muhammad Manzoor<sup>1\*</sup>, Mushtaq Ahmad<sup>1</sup>, Syed Waseem Gillani<sup>1\*\*</sup>,  
Muhammad Zafar<sup>1</sup>, Javeed Hussain<sup>2</sup>, Mohd Kafeel Ahmad Ansari<sup>3</sup>,  
Humaira Rizwana<sup>4</sup>, Islem Abid<sup>5</sup>, Muhammad Imran<sup>6</sup>, Shaista Jabeen<sup>1</sup>

<sup>1</sup>Department of Plant Sciences, Quaid-i-Azam University, Islamabad, Pakistan

<sup>2</sup>Department of Botany, University of Kotli, Azad Jammu and Kashmir, Pakistan

<sup>3</sup>Department of Biology, Faculty of Natural Science, University of Guyana, South America

<sup>4</sup>Department of Botany and Microbiology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

<sup>5</sup>Centre of Excellence in Biotechnology Research, King Saud University, P.O Box 2455, Riyadh 11495, Saudi Arabia

<sup>6</sup>Department of Botany, PMAS-Arid Agriculture University, Murree Road Rawalpindi, Pakistan

Received: 14 March 2024

Accepted: 28 June 2024

## Abstract

*Sorbus foliolosa* is a threatened high-altitude flowering (Rosaceae) species that is endemic to the Himalayan region. Due to ongoing habitat destruction, the population is decreasing at an alarming rate and has a narrow distribution range. So, there is an urgent need to evaluate the population size, distribution range, and conservation status in order to establish feasible conservation strategies. Recent field expeditions in the western Himalayan region of Kashmir at 4250 m elevation recorded *Sorbus foliolosa* populations. This study provides a detailed description of the morphology and micro-anatomy of the taxon under investigation, with a focus on seeds and leaf surfaces examined through a scanning electron microscope. On the adaxial leaf surface, images from SEM show raised and thickened periclinal walls and straight and thickened anticlinal walls. Irregular or striate patterns on the abaxial surface with anomocytic stomata. Reticulate sculpturing, raised and thickened periclinal walls, buttressed anticlinal walls, and irregular epidermal cell arrangements characterize seed surfaces. The species illustration, distribution map, and updated description are presented. *Sorbus foliolosa* is declared a threatened species in the Kashmir region because of its small population size, following the criteria of the IUCN Red List. We urge that policymakers and the forest department implement conservation strategies

\*e-mail: manzoorbaloach@gmail.com

\*\*e-mail: sgillani@bs.qau.edu.pk

and the involvement of local communities to conserve this critically endangered species in the Kashmir Himalayas.

**Keywords:** conservation, IUCN, Kashmir Himalayas, population size, SEM

## Introduction

*Sorbus* L. is a widely distributed genus. It inhabits temperate regions of Asia, North America, and Europe and comprises approximately 250 species when considered in its broadest sense. The highest diversity centers of the *Sorbus* genus are found in East Asia, particularly the western Himalayan region [1-3]. The morphological information [4] and molecular evidence [5] indicated that *Sorbus* has a polyphyletic nature. Generally, taxonomists classify them into five genera: *Chamaemespilus* [6], *Aria* [7], *Micromeles* [8], *Torminalis*, and *Sorbus* [6]. The *Sorbus* genus encompasses 80 species, each with distinct characteristics, such as pinnately compound leaves. The mountains of southern China, particularly the Hengduan mountains, along with the surrounding regions of Nepal, Myanmar, and the eastern Himalayas, have the greatest diversity of *Sorbus* species, with approximately 60 species found in these areas [3, 9-11]. This region is widely acknowledged as a worldwide biodiversity hotspot, and the genus *Sorbus* also thrives there [12, 13].

*Pyrus foliolosa* (Wall.) Spach, originally recognized as a *Numen nudum* by [14] and referred to as “677 *Pyrus foliolosa*” Wallich in 1824. *Pyrus foliolosa*, a synonym of *Sorbus foliolosa*, an endemic species found in the Central and Western Himalayan range, was given the name “*Napalia* 1821” in the original description [15]. However, Wallich 677 contains samples that were collected from two locations in Nepal, namely Gossain Than and Sheopore. In accordance with Art. 8.2 footnote 2 [16], it consists of two different collections of specimens of *P. foliolosa* and *P. wallichii* [17]. Hooker observed that the original description of *P. foliolosa* is a mixture of the abovementioned collections, which represent two separate taxa. He attempted to lectotypify *P. foliolosa* Wallich by limiting *P. foliolosa* to material collected from Gossain Than (lectotype), which has 9–11 pairs of leaflets with rusty brown hairs on the abaxial side of the leaflet, petiole, inflorescence, and also indicated *P. ursina* (Don, 1834) as its synonym. He named the Sheopore specimens (the remaining syntypes) as *P. wallichii* Hooker (1878), a new species that differs from *P. foliolosa* in having seven or eight pairs of leaflets as well as white wool on the petiole, inflorescence, and abaxial side of the leaflet.

*Sorbus* is one of the most difficult genera to classify in the Rosaceae family, owing to the complexity caused by polyploidy or genome doubling, which is frequently accompanied by gametophytic apomixis and hybridization [18]. The systematic position of related *Sorbus* species cannot be determined via traditional

morphological taxonomy, requiring molecular approaches. Some *Sorbus* species have been successfully studied using a variety of molecular systematic methods. The Rosaceae family is among the most significant plant families due to its vast diversity. There are 3000 species and 100 genera in the Rosaceae family. Herbaceous species are found in a variety of habitats, including temperate forests, high altitude zones, and roadside. Although woody taxa are less prevalent in mixed deciduous forests, even woody species are pioneers and are important in the early stages of forest succession [13]. Recent studies have reclassified the Rosaceae family based on a variety of features, including shape, structure, and genetic information. Maloideae is now included in the Pyreae tribe and Pyrinae subtribe according to this new classification [19]. The primary causes of the intricate taxonomy of the Maloideae are introgression and hybridization [20]. The subfamily Maloideae comprises the genus *Sorbus*, which is distinguished by its distinctive, pinnately complex leaves and fragile seed coat. Most *Sorbus* species have a significant degree of resemblance in floral structure and have inconspicuous floral colors, making it challenging to differentiate between them. Although there are few significant morphological features in the flowers, the majority of non-reproductive organ traits, such as pairs of leaflets, leaflet size, stipule form, serration location, shape and size of buds, fruit color, and indumentum, demonstrate the importance and distinctiveness of the genus *Sorbus* at the species level, particularly in a specific local region [21, 22].

In the Rosaceae and other angiosperm families, seed and pollen morphology have long been considered essential for taxonomic identification [23, 24]. Seed morphology and coat texture have been demonstrated to be useful traits for identifying *Sorbus* species [25]. Several palynologists believe that palynomorphological traits are significant for taxonomic classification and delineation within the Rosaceae family [23, 26]. Exine sculpture is extremely important, with prominent characteristics including the size and number of perforations and the variety and length of striae and grooves [27, 28]. Other key features for identifying Rosaceae include polar axis, endoaperture, pollen shape (P/E), and operculum structure [28, 29]. Previous research identified exine ornamentation (e.g. groove width, length, and orientation, as well as perforation number and diameter) and the P and P/E ratio, as well as aperture structure, as distinguishing properties of *Sorbus* pollen grains [30].

Due to its small restricted habitat niche, small population size, high grazing intensity, and global change impact, it is most vulnerable to extinction

for its population at higher elevations. In addition, due to some drawbacks, like inaccessibility due to the remote mountainous region, little is known about the distribution patterns, ecology, and conservation of threatened species. Our main objectives were as follows: 1) to assess the population size and distribution of species across the mountainous region of western Himalaya 2) to provide ecological insight and determine factors influencing the *Sorbus foliolosa* population, and 3) to update the morphological and anatomical description while suggesting reasonable conservation strategies for *Sorbus foliolosa*.

## Materials and Methods

This species grows in rocky and sloppy habitats, with a preference for the northern aspect. The climate of Phullawae Valley is alpine, with mostly hilly terrain and steep natural slopes with gradients of up to 60°. The alpine zone experiences extreme cold during the winter season, with heavy snowfall and freezing temperatures dropping as low as -10°C from November

to April [31-35]. The average summer temperature, from June to August, is around 10°C, while the summer season is cold and short. The area receives approximately 1000 mm of precipitation annually, most of which falls as snow in winter (Fig. 1) [36, 37].

## Specimen Collection

During the floristic exploration in the Western Himalayan region of Kashmir, Phullawae Neelum Valley Hills, fresh plant specimens were collected between June and August 2021 at an altitude of 4250 m. The collected specimens were subsequently dried, pressed, and mounted on the herbarium sheets by following standard protocols [38, 39]. The dried plant sample was then submitted to the Herbarium of Pakistan (ISL) and assigned Accession Number ISL-1333386. The plant name was validated by Plants of the World Online (<https://powo.science.kew.org/>). After conducting a thorough survey of existing literature, no previous record of *Sorbus foliolosa* was found in the Kashmir Himalayan region.




Fig. 1. A map illustrates the position and location of *Sorbus foliolosa* sampling sites in the Himalayan region of Kashmir.

### Morphological Investigation

The morphological characteristics of the samples were evaluated using a hand lens with 30-40x magnification. Before proceeding with further analysis, fresh leaf samples were sliced and treated with a solution of 70% alcohol and 30% acetic acid in a 1:3 ratio for 48 hours.

### Anatomical and SEM Examination

A foliar anatomy slide was prepared using the protocols of Clarke (1960). Small pieces of leaves were placed in the test tube with the addition of one part of lactic acid (25%) and three parts of nitric acid (75%) for the decolorization of the plant material. After heating for 5-10 minutes at 100°C, the solution was placed in a petri dish and given to two water washes for the removal of debris. A prepared piece of leaf was placed on a slide, coated with a cover slip, and observed under LM. The foliar anatomy and pollen slides were examined using an LM (OPTIKA Microscope, Italy), and photographs were taken using the Meiji Affinity DK-5000 digital camera.

Before SEM examination, dried seeds and leaves were extracted and washed with ethanol to remove dust. After cleaning, the seeds and leaves were adhered to stubs by using sticky tape, and a thin layer of gold-palladium was applied to the specimens. The SEM analysis was conducted using a model JEOL-5910, and the Polaroid P/N 665 film was used for the photomicrographs. The leaf and seed micromorphological features were investigated using the techniques outlined by [40] and [41]. The fertility of pollens (fertile and sterile pollens) was examined by following the protocols of Ahmad et al. [42].

### Map Generation

Russian maps at 1:10,000 scale were utilized to generate the GIS database. and Arc GIS software was used to map the location of the *Sorbus foliolosa* (Arc-Map, 10.8 Version). The data was exported in Geo-TIFF format and tiled using Arc View GIS software. After developing the species inventory map, the AJK district's boundaries were digitized using Arc Map software [39, 43]. A digital illustration was created




Fig. 2. Digital illustration of *Sorbus foliolosa* representing stamens, pollen, fruit, seed, and petiole.



using Adobe Illustrator (Version 2019) (Fig. 2).

## Results

### Taxonomic Description

**Habit:** Shrubs. 1-4.5 m tall. Branchlets are greyish brown to brownish, terete or round. **Leaf macro-morphology:** Young leaves have densely rusty hairs, sometimes hairless (Fig. 3); lenticels are oblong; buds are ovoid with a blunt apex and a size of  $(3.7-6.2 = 4.34 \pm 1.92)$  mm; several scales have the color brown and are puberulous apically. Leaves are imparipinnate, joined with rachis  $(7.2-12.1 = 2.0.3 \pm 1.42)$  cm; petiole  $(1.4-1.9 = 2.70 \pm 0.58)$  cm; stipules cauducous, linear to lanceolate,  $(4.4-8 = 1.54 \pm 0.21)$  mm, margin serrate; rachis sulcate, slightly rusty, narrow winged; leaflets blades 5-9 pairs, sub-leathery, at intervals of  $(0.1-1.3 = 3.01 \pm 0.87)$  cm. **Leaf micro-morphology by LM:** Adaxial surface is dark green, length of epidermal cells  $(33.08-47.03 = 39.70 \pm 2.37)$  mm, width of epidermal cells  $(19.77-25.68 = 22.61 \pm 1.07)$  mm, number of epidermal cells  $(98-115 = 107.50 \pm 3.06)$ , oblong or oblong-elliptic  $(2.2-3.3 = 1.02 \pm 0.03 \times 0.7-1.6 = 2.43 \pm 0.68)$  cm, abaxial surface is slightly hairy, length of epidermal cells  $(25.57-42.12 = 31.36 \pm 2.92)$  mm, width of epidermal cells  $(14.62-17.80 = 15.88 \pm 0.52)$  mm, number of epidermal cells  $(70-93 = 84.50 \pm 3.98)$ , anomocytic stomata and rarely actinocytic stomata,

elliptic stomata shape, length of guard cells  $(7.23-7.83 = 7.49 \pm 0.13)$  mm, number of stomata  $(38-53 = 43.50 \pm 2.57)$ , length of stomata  $(37.00-42.80 = 40.10 \pm 1.01)$  mm, width of stomata  $(28.08-30.58 = 29.35 \pm 0.48)$  mm, length of subsidiary cells  $(26.13-36.35 = 31.57 \pm 1.74)$  mm, width of subsidiary cells  $(9.95-13.13 = 11.02 \pm 0.56)$  mm, ovate stomatal pore shape, length of stomatal pore  $(24.88-27.03 = 25.77 \pm 0.42)$  mm, width of stomatal pore  $(12.13-13.63 = 12.68 \pm 0.26)$  mm, Stomatal index  $(66.52\%)$  mm (Fig. 4).

**Inflorescence:** Inflorescence are Compound corymbs with diameter of  $(4.7-9.8 = 5.20 \pm 2.01)$  cm, many flowers; pedicels having rusty hairs; lenticels are oblong; bracts cauducous  $(1.3-3.1 = 4.64 \pm 1.65)$  mm. Pedicel  $(1.2-2.3 = 1.47 \pm 0.45)$  mm. The flower's diameter is  $(4.7-9.8 = 32.22 \pm 1.79)$  mm, Hypanthium obconic, slightly pubescent or glabrous. Sepals are obconic, triangular, glabrous, and apex-acute. Petal's color is white, ovate-broadly obovate  $(2.1-2.8 = 1.54 \pm 0.88)$  mm, base clawed. Stamens are 15-20 times smaller than petals. Styles 2 or 4, free at base, hairless. Fruit color is red, ovoid or spherical, with a diameter of  $(4.3-6.2 = 3.21 \pm 2.01)$  mm and small lenticels; sepals are persistent.

**Pollen:** Pollen are tricolpate; axis length  $(29.75-36.25 = 33.85 \pm 1.13)$  mm, equatorial diameter  $(30.25-37.75 = 33.80 \pm 1.32)$  mm, number of colpi 3, length of colpi  $(6.75-7.75 = 7.30 \pm 0.20)$  mm, width of colpi  $(2.5-3.50 = 3.00 \pm 0.18)$  mm, exine thickness  $(2.25-3.00 = 2.57 \pm 0.14)$  mm, fertile pollen 116 (fertility 85.92%) and sterile pollen 19 (sterility 14.08%), mesocolpium



Fig. 3. Field photographs of *Sorbus foliolosa* (A) Stem and branchlets, (B) Fruit, (C) Leaves, (D) Habitat.



(32-34.00 = 33.00±0.40) mm (Fig. 4). Seed examination by Stereomicroscope: Seeds are brownish (0.9-2.2 = 2.25±0.14) mm, seed length (3.85-4.10=3.95±0.04) mm, seed width (1.51-2.10 = 1.87±0.11) mm, length/width (1.86-2.72 = 2.15±0.16) mm (Fig. 4). Leaves and seed micromorphology by SEM: Both adaxial and abaxial leaf surfaces were observed through SEM. The adaxial and abaxial leaf surfaces showed noticeable differences, as the adaxial leaf surface has no stomata, while the abaxial leaf surface contains an anomocytic stomatal type. The anticlinal walls of the adaxial surface are straight and thickened, while the periclinal walls are raised and thickened. On the abaxial surface, irregular/striate can be observed, along with anomocytic stomata

that have an elliptic shape and an ovate stomatal pore (Fig. 4). The seed shape of *S. foliolosa* is obovate, with an obtuse seed base and a light brown color. The seed outline appears smooth, with ribbed-reticulate surface sculpturing. The shape of the epidermal cell is irregular, with an unevenly thickened wall. The anticlinal wall is raised, straight to curved, whereas the periclinal wall is slightly concave. (Fig. 4). Phenology: Flowering time from June to August and fruiting July to September.

### Distribution

*Sorbus foliolosa* is an endemic species found in the Himalayan region. This species was observed and




Fig. 4. (a, b) Light microscopy of the foliar epidermal anatomy of *Sorbus foliolosa* (40x); (c, d) SEM micrographs of adaxial surface; (e, f) abaxial surface (10x) showing anomocytic stomata with irregular arrangements of cells (40x); (g, h) SEM micrographs of the abaxial surface; (i) Stereomicroscope photographs of seed morphology representing the brownish colors of achenes; (j) showing the general surface of the seed; (k, l) outline is smooth with a ribbed-reticulate surface; the anticlinal wall is straight to curved, and the periclinal wall is straight and depressed; (m, n) light microscopy; (LM) of pollen micrographs with polar view showing tricolpate pollen shape; (o, p) equatorial view of pollen.

collected at an elevation of 4250 m in the alpine zone of the Phullawae region in the Kashmir Himalayas.

### Conservation Status

*Sorbus foliolosa* is confined to the alpine zone of the Phullawae region in the Kashmir Himalayas. According to IUCN guidelines, which classify species with less than 250 individuals in a 100 km<sup>2</sup> area as critically endangered, only 3-4 populations of this species have been recorded in the Phullawae region [44].

### Associated Flora

*Allium victorialis* L., *Gentiana kurroo* Royle, *Geranium himalayense* Klotzsch, *Bergenia stracheyi* (Hook.f. & Thomson) Engl., *Anaphalis nepalensis* (Spreng.) Hand.-Mazz., *Artemisia roxburghiana* Besser, *Rheum webbianum* Royle, *Melanoseris brunoniana* (Wall. ex DC.) N. Kilian & Ze H.Wang, *Rhodiola fastigiata* (Hook.f. & Thomson) S.H. Fu, *Poa bulbosa* Steud., *Swertia alata* (D.Don ex G.Don) Royle ex C.B. Clarke, and *Lagotis cashmeriana* (Royle ex Benth.) Rupr.

### Discussion

*Sorbus foliolosa* is well distributed in the western Himalayan region of Kashmir, but its previous description was lacking in detail, and field records were incomplete. Furthermore, due to its restricted habitat in the Kashmir Himalayas, only a limited number of individuals have been recorded. The specimen closely resembles the original material in all aspects, which has allowed for a thorough description of the species. The *Sorbus* genus especially exhibits a higher degree of resemblance in terms of color and structure of inflorescence. Furthermore, there is a significant amount of uniformity among the groups in fruit color and leaflet number [13]. *Sorbus* fruits exhibit a variety of colors, such as white, pink, red, and orange-red. Additionally, the number of leaflet pairs in the *Sorbus* genus ranges from 2-21 pairs [3, 22]. The number of reliable floral morphological features for identification is limited. Identification of the species is according to its vegetative leaf morphology, including leaflet pairs, size, serration location, stipule shape, and fruit color. These characteristics have often been utilized to distinguish *Sorbus* species. Most notably, *Sorbus foliolosa* can be distinguished from other species of *Sorbus* by key features, including petiole length of (1.4-1.9 = 2.70±0.58) cm, cauducous stipules, linear-lanceolate leaves with serrate margins, 5-9 pairs of leaflet blades, absence of stomata on the adaxial surface, slightly pubescent abaxial surface with a large number of stomata, Compound corymbs Inflorescence, white petals, persistent sepals, 15-20 stamens, rounded red fruit, and brownish seed color [21, 45]. A valid

identification method that can assist with accurate taxonomic identification and authenticity is anatomical characters [46].

Previous research provided some details about the primary morphological characteristics of leaves. Adaxial surfaces lack stomata, whereas abaxial surfaces exhibit anomocytic stomata and rarely actinocytic stomata. The upper surface displays isodiametric epidermal cells, while the lower surface features irregular epidermal cells with a zigzag or wavy cell wall pattern [47]. Variations in epidermal size were also noted among Rosaceae species. Stomata exhibited the anomocytic type; a type of stomata was also reported in the *Rosa* genus [48], while other types have been found in the Rosaceae family [49-51].

The study demonstrates how seed shape, particularly seed micro sculpture, can be used to identify diverse species within the *Sorbus* genus. The shape of seeds is important for phylogenetic studies and species identification in the Rosaceae family [52]. Furthermore, several studies have shown that *Sorbus* exhibits significant variability in seed features such as size, shape, and surface patterns, which provide valuable taxonomic information [53]. Seeds are brownish, (0.9-2.2 = 2.25±0.14) mm, with a seed length of (3.85-4.10 = 3.95±0.04) mm and a seed width of (1.5-2.10 = 1.87±0.11) mm. *Sorbus foliolosa* seed outline appears smooth, with reticulate surface sculpturing, while other species of *Sorbus* like *S. meliosmifolia* and *S. megalocarpa* have light ribbed-reticulate patterns of seed surface, whereas *S. flogneri* and *S. alnifolia* can be distinguished by their seed surfaces having a polygonal to irregular cell pattern. These seed results are strongly supported by [30].

Pollens of *Sorbus foliolosa* are tricolpate, with an axis diameter (29.75-36.25 = 33.85±1.13) mm and an equatorial diameter (30.25-37.75 = 33.80±1.32) mm. This result is consistent with those of [52]. *S. scalaris*, *S. prattii*, *S. sargentiana*, *S. meliosmifolia*, *S. megalocarpa*, *S. koehneana*, *S. folgnieri*, *S. amabilis*, and *S. alnifolia* all show trizonocolporate pollen grains [30]. The pollen size of Chinese *Sorbus* taxa was found to be medium-sized, with the polar axis diameter ranging from 25.21-40.41 mm [54]. In contrast, some Polish *Sorbus* taxa had smaller pollen sizes as compared to *Sorbus foliolosa*, with a polar axis diameter ranging from 22.91-33.2 mm [55]. *Sorbus foliolosa* was collected from alpine pastures with harsh environmental conditions such as low temperatures, short growing seasons, and heavy snowfall. Due to the aforementioned reasons, the differences observed among various *Sorbus* species in pollen size can be attributed to increased temperatures and solar radiation in particular environments. According to research, plants that live in areas with higher temperatures and potential evapotranspiration produce fewer but larger pollen grains [56]. Current findings help in understanding the population size and distribution range of *Sorbus foliolosa* in the western Himalayan region of Kashmir. Since its first report had



only a little morphological description, this study offers detailed insights on the macro-morphological features by using LM and SEM. The study also identifies the factors that influence the population size of *Sorbus foliolosa*, evaluates conservation on a regional scale, and provides insights on future sustainable management and conservation of threatened species in the era of global change.

### Future Management

To preserve this species, we suggest creating a nature protection site rather than a nature reserve. Anthropogenic activities have substantially restricted the distribution regions of *Sorbus foliolosa*, with limited populations in mountainous areas being vulnerable to disturbances [57-61]. To preserve *Sorbus foliolosa* and its habitat, we urge a complete ban on activities like branch picking, cutting, and habitat trampling where this species is found. Specifically, more rigorous management practices should be implanted in these hotspot areas. Our goal is to reduce human disturbance and safeguard suitable habitats to foster population growth and expand distribution ranges. The government, stakeholders, and forest department should develop public awareness and outreach programs in areas where *Sorbus foliolosa* populations are present. Involving organizations and local communities is crucial for the long-term conservation of this threatened species and should be encouraged from the outset.

### Conclusions

*Sorbus foliolosa* is well distributed in the alpine zone of the Kashmir Himalayas, but there is limited documentation. In this study, we provide for the first time the micro-morphological features (leaves, pollen) and SEM analysis (leaf surfaces, seeds, pollen) of this species. Furthermore, this study provides diagnostic features for *Sorbus foliolosa* that facilitate its identification from other *Sorbus* species. A newly discovered population of *Sorbus foliolosa* with a small number of individuals needs to be protected by the forest department. In the era of global change and the decline of species, conservation strategies should be implemented to protect habitat as well as genetic diversity. Moreover, seed propagation and tissue culture should be used to support species conservation efforts.

### Ethical Approval and Consent to Participate

All the experiments were performed in accordance with relevant guidelines and regulations.

### Author Contributions

Muhammad Manzoor: Field Collection, Writing-Original draft preparation, Mushtaq Ahmad & Muhammad Zafar: Supervision and Review Editing, Muhammad Manzoor: Statistical analysis, Syed Waseem Gillani & Muhammad Imran: Field Collection, Methodology and Data analysis, Investigation, Syed Waseem Gillani, Shaista Jabeen & Muhammad Manzoor: Interpretation of Results. Humaira Rizwana & Islem Abid: Funding and data curation. Javeed Hussain and Mohd Kafeel Ahmad Ansari: Review, editing, Validation.

### Acknowledgment

The authors extend their appreciation to the Researchers Supporting Project number (RSPD2025R745), King Saud University, Riyadh, Saudi Arabia.

### Conflict of Interest

There is no declared conflict of interest between the authors in the publishing of this research work.

### Data Availability Statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

### References

1. ALDASORO J.J., AEDO C., NAVARRO C., GARMENDIA F.M. The genus *Sorbus* (Maloideae, Rosaceae) in Europe and in North Africa: morphological analysis and systematics. *Systematic botany*. **189**, 1998.
2. SENNIKOV A.N., KURTO A. A phylogenetic checklist of *Sorbus* sl (Rosaceae) in Europe. *Memoranda - Societatis pro Fauna et Flora Fennica*. **93**, 1, 2017.
3. LU L., SPONGBERG S. *Photinia lindley*. *Flora of China*. **9** 46, 2003.
4. ZHENG D., ZHANG M. A Cladistic and Phenetic Analysis of the Infrageneric Relationships of *Sorbus* s. l. (Maloideae, Rosaceae) Based on the Morphological Characters. *Acta Horticulturae Sinica*. **34** (3), 723, 2007.
5. CAMPBELL C., EVANS R., MORGAN D., DICKINSON T., ARSENAULT M. Phylogeny of subtribe *Pyrinae* (formerly the Maloideae, Rosaceae): limited resolution of a complex evolutionary history. *Plant systematics and evolution*. **266** (1), 119, 2007.
6. MEDICUS F.C. *Philosophische Botanik mit kritischen Bemerkungen*. Hof-und Akademischen. **1789**.
7. PERSON C.H. *Synopsis Plantarum: seu Enchiridium botanicum, complectens enumerationem systematicam specierum hucusque cognitarum*. Apud CF Cramerum. **1805**.



8. DECAISNE J. Mémoire sur la famille des Pomacées. Volume 10 di Archives, Muséum national d'histoire naturelle. France. **1874**.
9. LONG D., GRIERSON A., LONG D. Sorbus. 1, Part 3). Royal Botanic Garden Edinburgh, Edinburgh. **592, 1987**.
10. MCALLISTER H. The genus Sorbus: mountain ash and other rowans. Royal Botanic Gardens Kew. **2005**.
11. WATSON M., MANANDHAR V., WATSON M., AKIYAMA S., IKEDA H., PENDRY C., RAJBHANDARI K., SHRESTHA K. Sorbus L. Flora of Nepal. Royal Botanic Garden Edinburgh, Edinburgh. **25, 2012**.
12. MYERS N., MITTERMEIER R.A., MITTERMEIER C.G., DA FONSECA G.A., KENT J. Biodiversity hotspots for conservation priorities. Nature. **403** (6772), 853, **2000**.
13. LI M., OHI-TOMA T., GAO Y.-D., XU B., ZHU Z.-M., JU W.-B., GAO X.-F. Molecular phylogenetics and historical biogeography of Sorbus sensu stricto (Rosaceae). Molecular Phylogenetics and Evolution. **111** 76, **2017**.
14. WALLICH N. numerical list of dried specimens of plants in the East India Company's Museum. **1828**.
15. WALLICH N. Plantae Asiaticae Rariores; or Descriptions and Figures of a Select Number of Unpublished East Indian Plants, vol. 2. Treuttel and Würtz, London, 200 pp., **1831**.
16. LAHIRI S., DASH S.S. Lectotypification of three names in the genus Primula L. (Primulaceae). Kew Bulletin. **78** (4), 647, **2023**.
17. TURLAND N.J., WIERSEMA J.H., BARRIE F.R., GREUTER W., HAWKSWORTH D.L., HERENDEEN P.S., KNAPP S., KUSBER W.-H., LI D.-Z., MARHOLD K. International Code of Nomenclature for algae, fungi, and plants (Shenzhen Code) adopted by the Nineteenth International Botanical Congress Shenzhen, China, July 2017. Koeltz botanical books. **2018**.
18. VELEBIL J., LEPSÍ M., NOSKOVÁ J., LEPSÍ P. Taxonomic assessment of Sorbus subgenus Aria in the Malé Karpaty Mountains. Preslia. **94**, 305, **2022**.
19. WANG Q., NIU Z., LI J., ZHU K., CHEN X. The complete chloroplast genome sequence of the Chinese endemic species Sorbus setschwanensis (Rosaceae) and its phylogenetic analysis. Nordic Journal of Botany. **38** (2), **2020**.
20. WAGNER I., MAURER W., LEMMEN P., SCHMITT H., WAGNER M., BINDER M., PATZAK P. Hybridization and genetic diversity in wild apple (Malus sylvestris (L.) Mill.) from various regions in Germany and from Luxembourg. Silvae Genetica. **63** (1-6), 81, **2014**.
21. RUSHFORTH K. A preliminary revision of the Sino-Himalayan whitebeams (Sorbus section Thibeticae: Rosaceae). Yearbook of the International Dendrology Society. **2010**.
22. LU L., SPONGBERG S. Sorbus L. In: Wu ZY, Raven PH, Hong DY (Eds) Flora of China 9. Science Press, Beijing & Missouri Botanical Garden Press, Saint Louis. 144, **2003**.
23. ULLAH F., GAO Y.-D., ZAMAN W., GAO X.-F. Pollen morphology of Rosa sericea complex and their taxonomic contribution. Diversity. **14** (9), 705, **2022**.
24. BASER B., SAGIROGLU M., DOGAN G., DUMAN H. Morphology of pollen in Ferula genus (Apiaceae). PhytoKeys. **179**, 111, **2021**.
25. BEDNORZ L. Morphological variability of fruits and seeds of Sorbus torminalis in Poland. Dendrobiology. **57** (3), **2007**.
26. SONG J.-H., MOON H.-K., HONG S.-P. Pollen morphology of the tribe Sorbarieae (Rosaceae). Plant systematics and evolution. **302**, 853, **2016**.
27. LECHOWICZ K., BOCIANOWSKI J., WRÓŃSKA-PILAREK D. Pollen morphology and variability of species from the genus Rubus L. (Rosaceae) alien and invasive in Poland. Webbia. **76** (1), 109, **2021**.
28. WRÓŃSKA-PILAREK D., SOWELO M., ANTKOWIAK W., BOCIANOWSKI J., LECHOWICZ K. Pollen morphology and variability of native and alien, including invasive, species of the genus Spiraea L. (Rosaceae) in Poland. PloS One. **17** (8), e0273743, **2022**.
29. PATHAK M.L., IDREES M., XU B., GAO X.-F. Pollen morphology of subfamily Maloideae (Rosaceae) with special focus on the genus Photinia. Phytotaxa. **404** (5), 171, **2019**.
30. QIU Z., ZHAO Y., QIAN Y., TAN C., TANG C., CHEN X. Seed and Pollen Morphology in Nine Native Chinese Species of Sorbus (Rosaceae). International Journal of Fruit Science. **23** (1), 87, **2023**.
31. MANZOOR M., AHMAD M., ZAFAR M., GILLANI S.W., SHAHEEN H., PIERONI A., AL-GHAMDI A.A., ELSHIKH M.S., SAQIB S., MAKHKAMOV T. The local medicinal plant knowledge in Kashmir Western Himalaya: a way to foster ecological transition via community-centred health seeking strategies. Journal of Ethnobiology and Ethnomedicine. **19** (1), 56, **2023**.
32. SHAHEEN H., AZIZ S., NASAR S., WAHEED M., MANZOOR M., SIDDIQUI M.H., ALAMRI S., HAQ S.M., BUSSMANN R.W. Distribution patterns of alpine flora for long-term monitoring of global change along a wide elevational gradient in the Western Himalayas. Global Ecology and Conservation. **48**, e02702, **2023**.
33. GILLANI S.W., AHMAD M., ZAFAR M., HAQ S.M., WAHEED M., MANZOOR M., SHAHEEN H., SULTANA S., REHMAN F.U., MAKHKAMOV T. An Insight into Indigenous Ethnobotanical Knowledge of Medicinal and Aromatic Plants from Kashmir Himalayan Region. Ethnobotany Research and Applications. **28** (1), **2024**.
34. MANZOOR M., AHMAD M., ZAFAR M., HAQ S.M., SHAHEEN H., WAHEED M., GILLANI S.W., SULTANA S., MAKHKAMOV T. Unveiling the Indigenous Ethnomedicinal knowledge of Genus Nepeta from Division Muzaffarabad, Azad Jammu & Kashmir, Pakistan. Ethnobotany Research and Applications. **26** (1), **2023**.
35. MIRZAMAN Z., KAYANI S., MANZOOR M., JAMEEL M.A., WAHEED M., GILLANI S.W., BABAR C.M., BUSSMANN R.W. Ethnobotanical study of Makra Hills district Muzaffarabad, Azad Jammu and Kashmir, Pakistan. Ethnobotany Research and Applications. **26**, 1, **2023**.
36. PAK-MET. The normals of climatic data of Azad Jammu & Kashmir. Pakistan Meteorological Department, Regional Centre Muzaffarabad, Pakistan. **2018**.
37. KHAN S., SHAHEEN H., AZIZ S., NASAR S. Diversity and distribution of Genus Primula in Kashmir region: an indicator genus of the western Himalayan mountain wetlands and glacial forelands. Biodiversity and Conservation. **30** (6), 1673, **2021**.
38. SMITH B., CHINNAPPA C. Plant collection, identification, and herbarium procedures. Plant microtechniques and protocols. **541**, **2015**.
39. SHAHEEN H., ATTIQUE A., RIAZ M.T., MANZOOR M., KHAN R.W.A., RIAZ M.T. From biodiversity hotspot to conservation hotspot: assessing distribution, population structure, associated flora and habitat geography of threatened Himalayan Yew in temperate forest ecosystems of Kashmir. Biodiversity and Conservation. **1**, **2024**.

40. BAHADUR S., AHMAD M., ZAFAR M., GUL S., AYAZ A., ASHFAQ S., RUBAB S., SHUAIB M., AHMAD S. Taxonomic study of one generic and two new species record to the flora of Pakistan using multiple microscopic techniques. *Microscopy Research and Technique*. **83** (4), 345, **2020**.
41. JABEEN S., ZAFAR M., AHMAD M., ALTHOBAITI A.T., OZDEMIR F., KUTLU M., MAKHKAMOV T.K., SULTANA S., AMEEN M., MAJEED S. Ultra-sculpturing of seed morphotypes in selected species of genus *Salvia* L. and their taxonomic significance. *Plant Biology*. **25** (1), 96, **2023**.
42. AHMAD M., BANO A., ZAFAR M., KHAN M.A., CHAUDHRY M.J.I., SULTANA S. Pollen morphology of some species of the family asteraceae from the alpine zone, Deosai Plateau, northern Pakistan. *Palynology*. **37** (2), 189, **2013**.
43. ALAM N.M., SHAHEEN H., MANZOOR M., TINGHONG T., ARFAN M., IDREES M. Spatial Distribution and Population Structure of Himalayan Fir (*Abies pindrow* (Royle ex D. Don) Royle) in Moist Temperate Forests of the Kashmir Region. *Forests*. **14** (3), 482, **2023**.
44. IUCN <https://www.iucnredlist.org/resources/categories-and-criteria>. **2022**.
45. CHEN X. Taxonomic and nomenclatural notes on *Sorbus foliolosa* and related taxa (Rosaceae). *Phytotaxa*. **382** (3), 297, **2018**.
46. SHAHEEN S., SHARIFI-RAD J., ALI M., SHAMIM Z., RASOOL B., ASHFAQ M., MUKHTAR H., HARUN N., HANIF U., SIDDIQUE R. Light and scanning electron microscopy-based foliar morpho-anatomical comparison of selected family Rosaceae members distributed in District Lahore, Punjab, Pakistan. *Microscopy Research and Technique*. **85** (4), 1597, **2022**.
47. BURDA S., MARCELA P.I., TOMESCU C. Aspects Regarding Morpho-Anatomy, Bio-Ecology And Chorology Of *Sorbus intermedia* (Ehrh.) Pers.(Rosaceae) From Romania. **2006**.
48. SHAMSO E., SADEK A., HOSNI H.A. Morphological and anatomical characteristics of endemic *Rosa arabica* (Rosoideae, Rosaceae) from Sinai, Egypt. *Taeckholmia*. **39** (1), 34, **2019**.
49. AL-DOSKEY Z.A.K.S. Taxonomic Value of Leaf Epidermal Characters of the Genera *Rosa* L. and *Rubus* L. (Rosaceae) in Iraq. IOP Publishing. **2023**.
50. BABOSHA A., RYABCHENKO A., KUMACHOVA T., KOMAROVA G., YATSENKO I. Micromorphology of the leaf surface in some species of Dryadoideae (Rosaceae). *Micron*. **167**, 103428, **2023**.
51. KUMACHOVA T., BABOSHA A., RYABCHENKO A., IVANOVA T., VORONKOV A. Leaf Epidermis in Rosaceae: diversity of the cuticular folding and microstructure. *Proceedings of the National Academy of Sciences, India Section B: Biological Sciences*. **91** (2), 455, **2021**.
52. CHUNG K.-S., ELISENS W.J., SKVARLA J.J. Pollen morphology and its phylogenetic significance in tribe Sanguisorbeae (Rosaceae). *Plant systematics and evolution*. **285**, 139, **2010**.
53. BEDNORZ L., WALKOWIAK R., MACIEJEWSKA-RUTKOWSKA I., MOLINSKI K. Seed variability of the Polish species of the genus *Sorbus* (Rosaceae). *Dendrobiology*. **55** (3), **2006**.
54. YANG L., WU Y., PEI X., GUAN X., ZHENG J. Pollen morphological characteristics and cluster analysis on some species in *Sorbus* Linn. *Journal of Plant Resources and Environment*. **28**, 84, **2019**.
55. BEDNORZ L., MACIEJEWSKA-RUTKOWSKA I., WRONSKA-PILAREK D., FUJIKI T. Pollen morphology of the Polish species of the genus *Sorbus* L. *Acta Societatis Botanicorum Poloniae*. **74** (4), **2005**.
56. EJSMOND M.J., WRONSKA-PILAREK D., EJSMOND A., DRAGOSZ-KLUSKA D., KARPIŃSKA-KOŁACZEK M., KOŁACZEK P., KOZŁOWSKI J. Does climate affect pollen morphology? Optimal size and shape of pollen grains under various desiccation intensity. *Ecosphere*. **2** (10), 1, **2011**.
57. NOOR W., ZAFAR M., AHMAD M., SADIA B., GILLANI S.W., MANZOOR M. AMEEN M. Taxonomic significance of palyno-morphic markers for the delimitation of some Brassicaceous taxa in Balochistan Province (Pakistan). *Genetic Resources and Crop Evolution*. **1**, **2024**.
58. SADIA B., AHMAD M., ZAFAR M., NOOR W., MANZOOR M., GILLANI S.W., SULTANA S. Taxonomic implications of petiole microanatomical traits among Asteraceae tribes from arid-semiarid regions of Baluchistan, Pakistan. *Genetic Resources and Crop Evolution*. **1**, **2024**.
59. KAYANI S., AHMAD M., GILLANI S.W., MANZOOR M., REHMAN F.U., JABEEN S., SHAH S.A.H. Ethnomedicinal appraisal of the medicinal flora among the sub-alpine and alpine indigenous communities of Palas Valley Kohistan, Northern Pakistan. *Ethnobotany Research and Applications*. **28**, 1, **2024**.
60. MANZOOR M., AHMAD M., ZAFAR M., GILLANI S.W., SHAH G.M., SHAHEEN H., KHISHLATOVNA K.K. Exploration of traditional Ethno-gynaecological knowledge: advances to ethnobotanical studies from indigenous communities of Neelum Valley in the Himalayan Region. *Plant Science Today*. **11** (spl), **2024**.
61. GILLANI S.W., AHMAD M., ZAFAR M., MANZOOR M., SHAH G.M., SHAHEEN H., KHISHLATOVNA K.K. Ethnobotanical Exploration of Traditional Medicinal Plants Among the Rural Inhabitants of District Muzaffarabad, Kashmir Himalayan Region. *Plant Science Today*. **11** (spl), **2024**.