

Original Research

Characteristics of Bacterial Communities in Rhizosphere and Non-Rhizosphere Soil of the Relict Plant *Diplandrorchis sinica* S. C. Chen

Lijie Zhang^{1,2}, Jingwen Wu^{1,2}, Haiyan Li³, Baiting Qin⁴, Qiang Zhou⁵, Yudi Zhao¹,
Bo Qu⁶, Xuhui Chen^{6*}

¹College of Forestry, Shenyang Agricultural University, Shenyang 110866, Liaoning, China

²Key Laboratory of Forest Tree Genetics, Breeding and Cultivation of Liaoning Province, Shenyang 110866, Liaoning, China

³Liaoning Laotudingzi national nature reserve fushun administration, Xinbin 113208, Liaoning, China

⁴College of Life Science and Bioengineering, Shenyang Institute of Technology, Shenyang 113122, Liaoning, China

⁵Liaoning Provincial Forestry and Grassland Bureau, Shenyang 110001, Liaoning, China

⁶College of Bioscience and Biotechnology, Shenyang Agricultural University, Shenyang 110866, Liaoning, China

Received: 24 August 2023

Accepted: 30 June 2024

Abstract

Diplandrorchis sinica is a monotypic orchid relict plant. Soil habitat and soil microorganisms are the main factors for the growth and development of orchid species. In order to find out the soil physical and chemical properties and soil microbial community characteristics in the growing area of *D. sinica* and expand the population of *D. sinica*, the rhizosphere and non-rhizosphere soils in the Rare and Endangered Species Reserve of Fushun, Laotudingzi Nature Reserve, Liaoning Province, were collected for Illumina Miseq high-flux sequencing. The Alpha and Beta diversity and LEfSe of the sequencing data were analyzed, combined with the physical and chemical properties of the soil in the growth area of *D. sinica* to study the changes of soil physical and chemical properties, soil bacterial composition and diversity in the growing area of *D. sinica*. The results showed that a total of 26,690 valid bacterial sequences were obtained by sequencing. After cluster analysis, 9,556 OTUs were classified into 34 phyla, 108 classes, 316 families, and 472 genera. Soil bacterial diversity is rich in *D. sinica*, and the main dominant bacteria in rhizosphere and non-rhizosphere soil were Proteobacteria, Bacteroidetes, Acidobacteria and Verrucomicrobia; at the level of order classification, Rhizobiales and Gaiellales were mainly rhizobiales; at genus classification level, *Hyphomicrobium* and *Rhizobium* were the main genera. The values of total nitrogen (STN), available phosphorus (AP), hydrolyzed nitrogen (HN), total phosphorus (TP), total potassium (TK), and pH in rhizosphere soil of *D. sinica* were significantly different from those in non-rhizosphere soil. Moreover, it was significantly correlated with Proteobacteria, Actinobacteria, and Verrucomicrobia in soil microorganisms. The soil physical and chemical properties affected the microbial and bacterial richness in rhizosphere soil of Diactylodes. LEfSE analysis showed that 13 indicator species were selected when the alpha level

*e-mail: xhchen@syau.edu.cn

was $\text{LDA} > 2$ and $p < 0.05$. The indicator species of Rhizo were *Yersinia*, Thermogemmatissporaceae, Thermogemmatissporales, *Spirosoma*, etc. Bulk is mainly Merhylophilaceae, Methylophilales, Chromatiales, etc. This study revealed for the first time the diversity of rhizosphere and non-rhizosphere soil bacterial communities of the endangered species *D. sinica*, laying a foundation for the research on the mechanism of the endangered species.

Keywords: *Diplandrorchis sinica*, Rhizosphere soil microorganisms, High-throughput sequencing, Bacterial diversity, Orchidaceae

Introduction

Diplandrorchis sinica, a saprophytic small herb of *Diplandrorchis* in Orchidaceae, is a unique monotypic orchid relict plant in China [1] and is a national Class II rare and endangered plant under protection [2]. The botanist Chen Xinqi (1979) discovered and proposed that *Diplandrorchis* is a very primitive and phylogenetically important new genus of Orchidaceae [2]. *D. sinica* is named for its two fertile stamens located in the apical ventral and dorsal direction of the stamen column. During the growth process, *D. sinica* may display flowers and no leaves. The plant has 13~17 pale green or greenish-white flowers. Petals are similar to labellum, anthers broadly ovate-oblong, stigma terminal, stamen beak absent, perianth nearly actinomorphic; the characteristics of nearly upright and rarely torsional flowers, upright stigma, and no stamen beak indicate that it should belong to the Bird nest orchid, one of the most primitive groups in Orchidaceae [3]. *D. sinica* lives saprophytic life. Years of investigation have found that it does not grow every year, and the number varies year to year, with a maximum of 14 plants found in one year. *D. sinica* is very demanding on the growing environment and has a short life cycle of only 15–20 days [4, 5]. Because the distribution area is extremely narrow, the population number of *D. sinica* is greatly limited. Therefore, how to save the small population and expand the population has become an urgent problem for orchid species conservation.

The germplasm resources of Chinese orchids are very rich, and the history of cultivation can be traced back to two thousand years ago [6]. Studies have shown that most orchids are rare and endangered species and have become the group of greatest concern in biodiversity conservation. In recent years, due to climate and environmental changes and the destruction of plant habitats, the number of endangered species of orchids has been decreasing, and the resources of orchids have been highly valued by researchers at home and abroad. At present, due to the limitation of the population and growth period of *D. sinica*, their pollination and reproduction methods are still unknown, which significantly restricts large-scale artificial breeding.

Soil microorganisms have been found to promote the growth and development of orchid roots, so the characteristics of their rhizosphere microbial community and rhizosphere ecological function have

become hot spots for research at home and abroad. Studies have shown that orchids (Orchidaceae) are typical mycorrhizal plants, with small and large seeds without endosperm, which need to be symbiotic with suitable mycorrhizal fungi to germinate under natural conditions [7]. Seed germination, plant growth, and development of *Cypripedium macranthum* are closely related to mycorrhizal fungi. Isolated a strain of *Epulorhiza* sp from the fibrous roots of wild *C. macranthum* and confirmed that the fungus had a promotional effect on the seed germination of *C. macranthum* [8]. Most orchid plants grow in symbiosis with soil fungi [7, 8], and this mycorrhizal symbiosis accompanies almost the entire life history of orchids from seed germination to flowering and fruiting [6]. Endophytic fungi and bacteria in roots, stems, and other tissues of some orchid species play a very important role in their growth and development [9], seed germination [7, 10], and stress tolerance [6]. For example, *Fusarium* is a non-mycorrhizal fungus in the roots of orchid plants, which is effective in promoting the germination of orchid seeds, although it can induce plant diseases under certain conditions [11]. In one study, the dominant endophytic fungi of Euphorbiaceae were found to show a gradient of decreasing fungal abundance at three different horizontal distances from the root, inter-root soil, and root-enclosed soil during different reproductive growth periods, and its microhabitat fungal community was characterized by obvious seasonal changes in community structure and composition [12]. Inter-root microorganisms can coexist with the plant root system, colonize and sustain in the root system, and play an important role in promoting plant growth and development [13]. Among these microorganisms, the utilization and sensitivity of bacteria to root secretions is much higher than that of fungi, and bacteria are the most active and dominant microorganisms in the inter-root [14].

In recent years, the studies on *D. sinica* mainly focus on distribution area and phenological observation [15, 16], habitat characteristics [17], habitat investigation [18], and isolation of mycorrhizal fungi [19]. As a rare and endangered species of Orchidaceae, the study of rhizosphere and non-rhizosphere soil bacterial diversity of *D. sinica* has not been reported at home and abroad. Therefore, in order to characterize the bacterial community in the soil microorganisms of *D. sinica*, this study took rhizosphere and non-rhizosphere soil as the research object, and sequenced the V3~V4 region fragments of 16S rRNA




Fig. 1. Protection zone and habitat conditions of *Diplandrorchis sinica*. (a) *D. sinica* protection group area; (b) *D. sinica* habitat; (c) *D. sinica* plant; (d) fleshy root.

gene of rhizosphere and non-rhizosphere soil bacteria based on Illumina Miseq high-throughput sequencing technology. The bacterial community structure and diversity characteristics of rhizosphere and non-rhizosphere soil of *D. sinica* were discussed at the molecular level, in order to provide scientific reference for the effective protection of endangered plant resources, and to lay a foundation for further discussion on artificial reproduction, habitat-mimicking cultivation and endangered mechanism of the *D. sinica* species.

Material and Methods

Soil Sample Collection

The growing site of *D. sinica* is located at Dadonggou Ranger Fire Station, Fushun Administration, Laohuadingzi Nature Reserve, Liaoning Province, 124°41'~125°05'E, 41°11'~41°21'N. The Rare and Endangered Conservation Area of *Biruilan* (Fig. 1-A) belongs to 2 subcompartments of 3 forest classes, and the survey area is 0.2 hectares. *Acer mono*, *Quercus mongolica*, *Populus davidiana*, *Maackia amurensis*, *Juglans mandshurica*, *Tilia amurensis*, *Carpinus cordata*, *Amygdalus davidiana*, *Fraxinus rhynchophylla*, *Phellodendron amurense*, *Ulmus Rubra* and other tree species grow in the reserve. The soil samples were collected from the Hongyan Forest Fire Station of Fushun Administration, Laotuding Nature Reserve, Liaoning Province (E124°41'13"–125°5'15", N41°11'11"–41°21'34") on August 10, 2016. In the habitat territory of *D. sinica* (Fig. 1-B) at an altitude of 632 m, soil with a depth of 0~20 cm was taken from the root zone of *D. sinica* (Fig. 1-C), the large soil chunks attached to the fleshy roots were shaken off, and then the rhizosphere soil attached to the fleshy roots was collected with small tweezers (Fig. 1-D). The soil was collected as rhizosphere soil, and 4 replicates were labeled as T1, T2, T3, and T4, respectively. At the same time, 10 collection sites were randomly set along the S-shaped path in its protection area, and the surface soil was removed. The sterilized ring knife was used to collect the soil at each collection site, and the soil was mixed and put into the sterilized

encapsulation bag, which was labeled as non-rhizosphere soil samples. The four repeated sites were labeled as T5, T6, T7, and T8. Soil samples were placed in sterile sealed plastic bags, placed in ice boxes, and brought back to the laboratory. Some soil samples were naturally air-dried, ground and screened indoors, and stored in a refrigerator at 4°C for the determination of basic physical and chemical properties of soil. The other part of the samples were stored in the refrigerator at -80°C for soil microbial index analysis.

Determination of Soil Physical and Chemical Properties

The samples from rhizosphere and non rhizosphere soils of *D. sinica* were used for physicochemical properties determination. The soil pH was measured using a pH meter. Total nitrogen (TN) was measured using the Kjeldahl method, in which the TN in soil samples was converted into NH_3 by boiling with concentrated H_2SO_4 and the catalytic agent, and then distilling with NaOH. The NH_3 was evaporated into a boric acid indicator and then titrated using a standard solution of HCl. Hydrolyzed nitrogen (HN) was measured using the alkaline hydrolysis diffusion method, in which the soil samples were hydrolyzed with NaOH solution in a sealed diffusion dish, and the HN was converted into NH_3 and constantly spread and escaped. The NH_3 was absorbed by the boric acid indicator and then titrated using a standard solution of HCL, the same as described above.

Total potassium (TK) and fast-acting potassium (AK) were measured using the flame photometric method. TK was determined using the melting method with NaOH in a silver crucible, in which the TK was converted into soluble potassium and then the TK concentration in the solution was determined by referring to the potassium standard curve after being measured using a flame photometer. The AK was determined by CH_3COOK immersion extraction. The soil samples were added into a 10-fold volume of CH_3COOK and then the extracted solutions were used for AK concentration determination, the same as described above.

Total phosphorus (TP) and available phosphorus (AP) were measured by the molybdenum blue colorimetric method. Total phosphorus extraction is the same as total potassium, which converts the total phosphorus into

soluble phosphorus and then adds dinitrophenol indicator and molybdenum antimony antichromogen. The solution was used for TP concentration determination by referring to the phosphorus standard curve after being measured using a spectrophotometer. The phosphorus standard curve was drawn by the concentrations of the standard phosphorus solution that reacted with the molybdenum antimony antichromogen and their absorption value. AP was extracted by NaHCO_3 immersion extraction, of which the available phosphorus in soil samples was dissolved in NaHCO_3 . The available phosphorus concentration in the solution was determined as the same as above. The total phosphorus and available phosphorus contents were then calculated.

The soil organic carbon (SOC) was measured by the potassium dichromate ($\text{K}_2\text{Cr}_2\text{O}_7$) external heating method. The SOC in the soil samples was converted into CO_2 after boiling in a $\text{K}_2\text{Cr}_2\text{O}_7$ solution, and the dichromate ion was reduced to a trivalent chromium ion. The remaining $\text{K}_2\text{Cr}_2\text{O}_7$ was titrated with the FeSO_4 standard solution, and then the SOC content was calculated according to the difference between the used FeSO_4 for soil samples and the blank control. Soil organic matter (SOM) content was calculated as $1.724 \times \text{SOC}$, and the carbon-nitrogen ratio in soil organic matter was calculated as SOC/TN .

Extraction of Total DNA from Soil Genome

DNA extraction was extracted using MO BIO Power Soil DNA Extraction Kit (MO BIO Laboratories, Carlsbad, CA, USA). DNA concentration and quality were checked using a NanoDrop Spectrophotometer. Extracted DNA was diluted to 10 ng/ μL and stored at -40°C for downstream use.

PCR Amplification

Universal primer for bacteria 16S: F515: 5'-GTGC-CAGCMGCCGCGG-3' and R909: 5'-CCCCGYCAATTC-MTTTTRAGT-3' with 12 nt unique barcode was used to amplify the V4 hypervariable region of 16S rRNA gene for pyrosequencing using Miseq sequencer [20, 21]. The PCR mixture (25 μL) contained 1x PCR buffer, 1.5 mM MgCl_2 , each deoxynucleoside triphosphate at 0.4 μM , each primer at 1.0 μM and 0.5U of ExTaq (TaKaRa, Dalian) and 10ng soil genomic DNA. The PCR amplification program included initial denaturation at 94°C for 3 min, followed by 30 cycles of 94°C for 40 s, 56°C for 60 s, and 72°C for 60 s, and a final extension at 72°C for 10 min. Conduct two PCR reactions for each sample, and combine them together after PCR amplification. PCR products were subjected to electrophoresis using 1.0% agarose gel. The band with the correct size was excised and purified using SanPrep DNA Gel Extraction Kit (Sangon Biotech, China, Cat# SK8132) and quantified with Nanodrop. All samples were pooled together with an equal molar amount from each sample. The sequencing samples were prepared using a TruSeq DNA kit according to the manufacturer's instructions. The purified library was diluted, denatured, re-diluted, mixed with PhiX (equal to 30% of the final DNA amount) as described in the Illumina library preparation

protocols, and then applied to an Illumina Miseq system for sequencing with the Reagent Kit v2 2 \times 250 bp as described in the manufacture manual.

Data Analysis

The sequence data were processed using QIIME Pipeline-Version 1.7.0 (<http://qiime.org/>). All sequence reads were trimmed and assigned to each sample based on their barcodes. The sequences with high quality (length > 150 bp, without ambiguous base 'N', and average base quality score > 30) were used for downstream analysis. Sequences were clustered into operational taxonomic units (OTUs) at a 97% identity threshold. The aligned ITS gene sequences were used for chimera check using the Uchime algorithm [22]. All the samples were randomly resampled to 28,894 reads. We calculated alpha-diversity (phylogenetic distance whole tree, chao1 estimator of richness, observed species, and Shannon's diversity index) and beta-diversity (PCoA, UniFrac) analyses, for which the rarefaction curves were generated from the observed species. Taxonomy was assigned using the Ribosomal Database Project classifier [23].

Results

Determination of Physical and Chemical Properties of Soil in *D. sinica* Habitat

From Table 1, we can see that the pH of the rhizosphere soil of *D. sinica* is higher than that of the non-rhizosphere soil, but there is no significant difference between the two, which is a neutral acidic soil; the contents of total nitrogen (TN), total potassium (TK), total phosphorus (TP), available phosphorus (AP), soil organic matter (SOM), hydrolyzed nitrogen (HN) and fast-acting potassium (AK) in the rhizosphere soil are significantly higher than that of the non-rhizosphere soil ($p < 0.01$); while the soil organic carbon (SOC) and soil carbon to nitrogen ratio of the rhizosphere soil are lower than that of the non-rhizosphere soil ($p < 0.01$). These results indicate that the growth of *D. sinica* requires a high SOM, HN, and AK, and neutral to acidic pH.

Bacterial Alpha Diversity in Rhizosphere and Non-Rhizosphere Soil of *D. sinica*

High-throughput sequencing of 16S rRNA V3-V4 regions of rhizosphere and non-rhizosphere soil bacteria of *D. sinica* was performed, and a total of 26,690 raw sequences were obtained from high-throughput sequencing of bacterial 16S rRNA genes. 26,471 valid sequences were obtained after splicing and filtering, and 9,556 OTUs belonging to 34 phyla, 113 orders, 361 families, and 551 genera were classified after clustering analysis. The Shannon index and Chao1 index of the bacterial community in rhizosphere soil were 10.42 and 6728.07, respectively, while the Shannon index and Chao1 index

Table 1. Analysis of physicochemical properties of rhizosphere and non-rhizosphere root soils of *Diplandrorchis sinica*.

Indicators	Rhizosphere soil	Non-rhizosphere soil
pH	6.93±0.08 A	6.79±0.38 A
Total nitrogen (g/kg)	3.08±0.04 A	0.29±0.05 B
Hydrolyzed nitrogen (mg/kg)	417.20±3.08 A	235.79±2.46 B
Total potassium (g/kg)	17.18±0.40 A	6.29±0.11 B
Fast-acting potassium (mg/kg)	241.21±5.40 A	189.56±1.84 B
Total phosphorus (g/kg)	7.73±0.08 A	2.00±0.08 B
Available phosphorus (μg/kg)	21.59±0.45 A	13.26±0.84 B
Soil organic carbon (g/kg)	24.39±0.46 B	5.69±0.235 A
Soil organic matter content (g/kg)	42.05±0.76 A	9.81±0.26 B
Carbon to nitrogen ratio in organic matter	7.92±0.06 B	20.16±2.86 A

Note: The different capital letters represented the values in the rhizosphere and non-rhizosphere soil differed significantly at $\alpha=0.01$ level.




Fig. 2. Rhizosphere (R) and non-rhizosphere (NR) soil bacterial diversity of *Diplandrorchis sinica*.

Table 2. Statistics of rhizosphere and non-rhizosphere soil bacterial species of *Diplandrorchis sinica*.

Soil style	Phylum	Class	Order	Family	Genus
R	32	106	204	312	452
NR	30	100	202	314	449
Both in R and NR	29	98	196	302	429

Note: R and NR meant rhizosphere and non-rhizosphere.




Fig. 3. Rhizosphere and non-rhizosphere soil bacterial diversity of *Diplandrorchis sinica*.

of the non-rhizosphere soil were 10.84 and 7298.39, respectively. It can be seen from Fig. 2 that the Shannon, Chao1, and Simpson indices of the rhizosphere soil were lower than those of the non-rhizosphere soil, with significant differences ($p < 0.05$).

Analysis of Bacterial Community Composition and Abundance in Rhizosphere and Non-Rhizosphere Soils of *D. sinica*

After filtering the original sequences and counting the number of each species in each sample, a statistical table of species of rhizosphere and non-rhizosphere soil bacteria of *D. sinica* can be obtained. There are no significant differences in the numbers of phylum, class, order, family, and genus between rhizosphere and non-rhizosphere soil of *D. sinica*, and most of the phylum, class, order, family, and genus are common both in rhizosphere and non-rhizosphere soil (Table 2).

At the phylum taxonomic level (Fig. 3-A), Proteobacteria dominated the rhizosphere and non-rhizosphere soil bacterial taxa of *D. sinica*, accounting for 41.87% and 37.97% ($p > 0.05$), respectively, followed by Actinobacteria, accounting for 19.93% and 27.45% ($p < 0.01$), respectively, Bacteroidetes, accounting for 9.57% and 8.11% ($p > 0.05$), respectively, and Acidobacteria, accounting for 9.01% and 8.11% ($p < 0.05$), respectively, Verrucomicrobia (7.67% and 3.23%, $p < 0.05$), and Planctomycetes (3.81% and 4.40%, $p > 0.05$), respectively.

At the level of bacterial class (Fig. 3-B), the dominant bacterial taxa in the rhizosphere and non-rhizosphere soils of *D. sinica* were mainly α -Alphaproteobacteria (20.39% and 22.55%, $p > 0.05$), Thermoleophilia (9.76% and 11.61%, $p > 0.05$), β -Alphaproteobacteria (9.27% and 9.67%, $p > 0.05$), Gammaproteobacteria (8.13% and 2.36%, $p > 0.05$), Spartobacteria (7.43% and 2.99%, $p < 0.05$), Saprospirae (5.67% and 7.14%, $p > 0.05$) and Actinobacteria (5.99% and 9.26%, $p < 0.01$).

At the level of bacterial orders (Fig. 3-C), the dominant bacterial taxa in the rhizosphere and non-rhizosphere soil of *D. sinica* were mainly Rhizobiales (16.68% and 18.22%, $p > 0.05$), Gaiellales (7.01% and 6.45%, $p > 0.05$), Chthoniobacterales (7.43% and 2.99%, $p < 0.05$), Saprospirales (5.67% and 7.14%, $p > 0.05$), and Actinomycetales (5.84% and 8.91%, $p < 0.05$), Sphingobacteriales (2.35% and 0.16%, $p < 0.05$), Burkholderiales (4.11% and 2.59%, $p > 0.05$), Xanthomonadales (2.11% and 1.70%, $p > 0.05$), RB41 (2.88% and 3.22%, $p < 0.05$), Syntrophobacterales (2.65% and 2.11%, $p < 0.05$), Solirubrobacterales (2.75% and 5.14%, $p < 0.05$).

At the bacterial family level (Fig. 3-D), the main dominant bacteria in the rhizosphere and non-rhizosphere soil of *D. sinica* were: Hyphomicrobiaceae (7.62% and 7.87%, $p > 0.05$), Gaiellaceae (6.72% and 5.96%, $p > 0.05$), Chthoniobacteraceae (7.43% and 2.99%, $p < 0.05$), Chitinophagaceae (5.62% and 7.09%, $p > 0.05$), Bradyrhizobiaceae (5.11% and 4.97%, $p > 0.05$), Sphingobacteriaceae (2.22% and 0.06%,

$p < 0.05$), Syntrophobacteraceae (2.65% and 2.11%, $p > 0.05$), Pseudomonadaceae (4.63% and 0.14%, $p < 0.05$), Nitrososphaeraceae (1.90% and 4.27%, $p < 0.05$), Xanthomonadaceae (1.24% and 0.39%, $p < 0.05$), Oxalobacteraceae (2.67% and 0.06%, $p < 0.05$) and Rhodospirillaceae (1.80% and 1.92%, $p > 0.05$).

At the bacterial genus level (Fig. 3-E), the main dominant bacteria in the rhizosphere and non-rhizosphere soil of *D. sinica* were: *Rhodoplanes* (5.08% and 5.04%, $p > 0.05$), DA101 (6.20% and 2.23%, $p > 0.05$), *Bradyrhizobium* (4.69% and 4.44%, $p > 0.05$), *Sphingobacterium* (1.85% and 0.001%, $p < 0.05$), *Pseudomonas* (4.62% and 0.14%, $p < 0.05$) and *Candidatus Nitrososphaera* (1.89% and 4.26%, $p > 0.05$), respectively. In addition, there are some unclassified Gaiellaceae, unclassified Chitinophagaceae, unclassified Syntrophobacteraceae, unclassified RB41, β -Amorphic (unclassified Betaproteobacteria), Rhizobiales (unclassified Rhizobia), Solirubrobacterales (unclassified Solirubrobacterales), and Rhodospirillaceae (unclassified Rhodospirillaceae), together accounting for 25.89% and 28.90%, respectively.

Relationship Between the Dominant Bacterial Taxa and Soil Physicochemical Properties in *D. sinica* Habitats

The redundancy analysis of rhizosphere and non-rhizosphere soil microorganisms and their soil physicochemical properties of *D. sinica* was performed, and the results can be seen in Fig. 4: for the two axes of the soil bacterial community RDA contributions were 68.66% and 24.28%, respectively, and the four samples of rhizosphere soil of *D. sinica* were clustered on one side, but T1, T2, T4, and T3 were far away from each other, indicating differences between samples, while the four samples of non-rhizosphere soil were clustered together. A total of 92% of the total characteristic values were explained by the five selected basic soil physicochemical properties, in which soil values of available phosphorus (AP), hydrolyzed nitrogen (HN), total phosphorus (TP), total potassium (TK), and pH of rhizosphere soils were significantly different from those of non-rhizosphere soils. Soil microorganisms of the phylum Proteobacteria, Actinobacteria, and Verrucomicrobia were significantly and positively correlated with each other and influenced the rhizosphere soil microbial abundance of *D. sinica*. Specifically, Actinobacteria, Chloroflexi, and Crenarchaeota were highly significantly negatively correlated with soil TK, TP, AP, soil organic matter content (SOM), HN, AK, and highly significantly positively correlated with TN, SOC (Soil organic carbon), and C/N ($p < 0.01$) (Fig. 5). The Verrucomicrobia showed highly significant positive correlations with TK and SOM content ($p < 0.01$), significant positive correlations with TP, AP, HN, AK ($p < 0.05$), and significant negative correlations with TN, SOC, and C/N ($p < 0.05$). Nitrospirae showed significant positive correlations with soil TK, TP, AP, SOM, HN, and AK ($p < 0.05$), significant negative correlation with




Fig. 4. RDA redundancy analysis of dominant rhizosphere (R) and non-rhizosphere (NR) bacteria and soil physical and chemical properties. T1-T4 represented the 4 duplicates of rhizosphere soil samples, and T5-T8 represented the 4 duplicates of non-rhizosphere soil samples. pH, potential of hydrogen; TN, total nitrogen; TK, total potassium; TP, total phosphorus; AP, available phosphorus; HN, hydrolyzed nitrogen.




Fig. 5. Correlation analysis between dominant bacterial phyla groups and soil physical and chemical properties. pH, potential of hydrogen; TK, total potassium; TP, total phosphorus; AP, available phosphorus; SOM, soil organic matter; HN, hydrolyzed nitrogen; AK, fast-acting potassium; TN, total nitrogen; SOC, Soil organic carbon; C/N, Carbon to nitrogen ratio.




Fig. 6. Branches of bacteria evolution in rhizosphere and non-rhizosphere soil of *Diplandrorchis sinica*.

TN and SOC ($p < 0.05$), and highly significant positive correlation with soil C/N ($p < 0.01$, Fig. 5).

Microbial Markers of Rhizosphere and Non-Rhizosphere Soil Bacteria in *D. sinica*

The classification level of the tree diagram in Fig. 6 proceeds from the phylum to the genus. Node size indicates the relative abundance of each sample grouping. Nodes of different colors indicate significantly enriched microorganisms in the corresponding groups, which had a significant effect on the differences between groups, while yellow nodes indicate that microorganisms were not significantly different in any of the different groups.

LefSe was used to analyze the rhizosphere and non-rhizosphere soil microorganisms of *D. sinica* separately to find the indicator species for each treatment. Linear discriminant analysis (LDA) was performed on samples according to different grouping conditions to identify bacteria that had a significant differential effect on sample delineation. The results showed that rhizosphere soil bacterial

microbial diversity was rich, with significant differences ($p < 0.05$) between rhizosphere and non-rhizosphere soil bacterial microorganisms (as in Fig. 5); when $LDA > 2$, $p < 0.05$, the main rhizosphere soil bacterial biomarkers of *D. sinica* were *Yersinia*, Thermogemmatissporales, *Spirosoma*, NRP_I, *Cellvibrio*, Alteromonadaceae, MBGA and TM1; non-rhizosphere soil bacterial biomarkers are mainly Methylophilaceae, Methylophilales, Chromatiales, and ML635J_21 (e.g., Fig. 7).

Discussion

The pH of the rhizosphere soil of the endangered species of *D. sinica* was slightly higher than the pH of the non-rhizosphere soil, which may be due to the secretion produced by the fleshy roots of *D. sinica* into the rhizosphere soil, thus causing the high pH of the rhizosphere soil. The contents of total nitrogen, total potassium, total phosphorus, available phosphorus, soil organic matter content, hydrolyzed nitrogen, and available potassium




Fig. 7. LDA effect size of differentially abundant groups in rhizosphere and non-rhizosphere soil.

in the rhizosphere soil of *D. sinica* were significantly higher than those in the non-rhizosphere soil ($p < 0.05$), while the contents of total nitrogen, soil organic carbon, and soil carbon to nitrogen ratio in the rhizosphere soil of *D. sinica* were lower than those in the non-rhizosphere soil, and the differences between them were highly significant ($p < 0.01$). This indicates that the growth of *D. sinica* can increase the rhizosphere soil pH, and the organic matter secreted by the rhizosphere during the growth process, enters the rhizosphere soil, which leads to the difference between the rhizosphere soil and non-rhizosphere soil in terms of physicochemical properties. This phenomenon of nutrient enrichment in the soil in the rhizosphere region occurs in *Solidago canadensis*, *Larix principis-rupprechtii*, *Paris polyphylla*, and *Choerospondias axillaris* among other species are consistent with the findings [24–27].

Soil microorganisms are important components of soil, the analysis of soil Illumina Miseq of the rare and endangered plant indicated that the dominant taxa of rhizosphere and non-rhizosphere soil bacteria of *D. sinica* were mainly Proteobacteria, Alphaproteobacteria, Rhizobiales, Hyphomicrobiaceae, and unclassified Gaiellaceae genus. However, the inter-rhizosphere soil microbial bacterial α -diversity in *D. sinica* was lower than in non-rhizosphere soils, and the two were significantly different ($p < 0.05$), and this finding is consistent with the study of inter-rhizosphere microorganisms of the endangered plant *Cypripedium macranthos* [28]. Similar results were also reported in the research about the soil bacterial community structure diversity in vineyards [29] and the natural secondary forest soil microbial community characteristics of *Quercus mongolica* [30]. Currently, *Sphingosine* bacteria and *Branchomycetes* have been isolated from the roots of *Dendrobium moschatum*, and these two strains of bacteria were inoculated with *D. moschatum* seeds, and the results

showed that the species could promote seed germination [28]. Many domestic scholars have also identified *Pseudomonas* and *Bacillus* as the dominant orchid species from endophytic bacterial isolation of orchid species such as *Cymbidium*, *Dendrobium*, and *Goodyera* [31–33].

Soil inter-rhizosphere is a special ecosystem with significant differences in chemical and biological properties between rhizosphere and non-rhizosphere soils. Rhizosphere soil is susceptible to plant roots and is a tiny area where complex interactions between beneficial and harmful microorganisms and host plants occur. Root secretions and the microenvironment also influence soil physical and chemical properties and soil microbial community distribution [34, 35]. Therefore, plant roots, soil environment, and soil microorganisms interact with each other to form a stable and reciprocal relationship that promotes plant growth and development [36–37]. Plants also have active selectivity on the rhizosphere soil bacterial community structure [38]. Plants change the inter-root environment through root activities, thus selectively increasing or decreasing the abundance or diversity of certain rhizosphere soil bacterial taxa, so that the abundance of bacterial species beneficial to the plant's own growth increases, while bacterial species unfavorable to its own growth decreases or even disappears, thus forming an inter-root bacterial community structure beneficial to itself. This results in different plants having different rhizosphere bacterial composition structures [39]. The results of this study revealed that soil values of total nitrogen (TN), active phosphorus (AP), hydrolyzed nitrogen (HN), total phosphorus (TP), total potassium (TK), and pH in the rhizosphere soil of *D. sinica* were significantly different from those of non-rhizosphere soil; and were significantly correlated with soil microorganisms of Proteobacteria, Actinobacteria, and Verrucomicrobia.

These dominant groups influence the abundance of microbial bacteria in the rhizosphere soil of *D. sinica*. *Bradyrhizobium*, *Sphingobacterium*, and *Pseudomonas* are the main genera in the rhizosphere soil of *D. sinica*, which play important roles in increasing the organic matter content in the rhizosphere soil [40], promoting nitrogen fixation by rhizosphere microorganisms [41] and suppressing fungal diseases [42]. Meanwhile, we used LEfSe analysis in order to explore the indicator species among soil microbial bacteria affecting the endangerment of *P. bifidum*, and found that hundreds of indicator species could be found when the alpha level was 0.05 (catalog lefse 0.05), but no indicator species could be found when the alpha level was 0.01, so we chose 13 indicator species at the condition of $LDA > 2$, $p < 0.05$, of which the main inter-rhizosphere soil bacterial biomarkers of *D. sinica* were *Yersinia*, Thermogemmatissporaceae, Thermogemmatissporales, *Spirosoma*, NRP_I, *Cellvibrio*, Alteromonadaceae, MBGA, and TM1, indicating that the fleshy roots of *D. sinica* are significantly enriched for these groups during growth; while the non-rhizosphere soil bacterial biomarkers are mainly Methylophilaceae, Methylophilales, Chromatiales and ML635J_21.

The discovery of these indicator species has laid the foundation for the study of the endangerment mechanism of *D. sinica*. In addition, from the analysis of the high-throughput sequencing results of the rhizosphere and non-rhizosphere soils in this study, there is still a large number of unidentified groups of soil bacteria in the inter-rhizosphere soil of *D. sinica*, among which the key bacteria affecting the endangerment of *D. sinica* may be unearthed, which will be analyzed in depth in the next step.

Conclusions

In this study, we analyzed the physical and chemical properties and microbial community characteristics of rhizosphere and non-rhizosphere soil in the *D. sinica* population. The results showed a significant difference in total nitrogen (TN), available phosphorus (AP), hydrolyzed nitrogen (HN), total phosphorus (TP), total potassium (TK), and pH in rhizosphere and non-rhizosphere soil of the *D. sinica* population. The soil physical and chemical properties affected the microbial and bacterial richness in rhizosphere soil of the *D. sinica* population. Soil bacterial diversity is rich, with the main dominant bacteria Proteobacteria, Bacteroidetes, Acidobacteria, and Verrucomicrobia at the phylum level; α -Alphaproteobacteria, Thermoleophilia, β -Alphaproteobacteria, Gammaproteobacteria, Spartobacteria, Saprospirae, and Actinobacteria at the class level; Rhizobiales, Gaiellales, Chthoniobacteriales, Saprospirales, and Actinomycetales at the order level; Hyphomicrobiaceae, Gaiellaceae, Chthoniobacteraceae, Chitinophagaceae, and Bradyrhizobiaceae at the family level; *Rhodoplanes*, DA101, *Bradyrhizobium*, and *Pseudomonas* at the genus level. LEfSe analysis showed indicator bacteria in rhizosphere and non-rhizosphere soil, which were *Yersinia*, Thermogemmatissporaceae,

Thermogemmatissporales, *Spirosoma*, NRP_I, *Cellvibrio*, Alteromonadaceae, MBGA, and TM1 in rhizosphere soil and Methylophilaceae, Methylophilales, Chromatiales, and ML635J_21 in non-rhizosphere soil.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (grant number 31630378) and the Conservation and Rescue Project of the State Forestry Bureau of China (2013028).

Conflict of Interest

The authors declare no conflict of interest.

References

1. CAO W., WU Y.Y., LI Y., CONG X.X. Priority conservation areas for threatened plants in Northeast China. *Journal of Applied Ecology*, **24** (2), 326, **2013**.
2. CHEN X.Q. *Dichondra*, a new genus of Orchidaceae that is very primitive and important in phylogeny. *Journal of Plant Taxonomy*, **17** (1), 1, **1979**.
3. Flora of China Editorial Committee. *Flora of China, Orchidaceae*. Beijing: Science Press, China, **25**, pp. 183, **2009**.
4. ZHU Y.P., LI S.Q., LIU Y.Q., JU W.P., SHI B.L., TIAN H.S. A preliminary report on the regional survey of the distribution of the population of *Spodoptera sinensis*. *Liaoning Forestry Science and Technology*, **1**, 23, **2004**.
5. ZHANG L.J., SHEN H.L., CUI J.G., ZHOU Q., JU W.P., LI H.Y. Rare and Endangered *Plant-Dichondra*, *Liaoning Forestry Science and Technology*, **6**, 50, **2008**.
6. YANG Q.Y. Study on Diversity of Orchidaceae Mycorrhizal Fungi and Its Influence on Orchid[D]. Dissertation for the Degree, Chinese Academy of Forestry, **6**, 1, **2018**.
7. ZHENG X.J., YE J., CUAN C.D., MA H.Y. Review on Recent Progress of Orchid Seeds Germination. *Northern Horticulture*, (19), 206, **2010**.
8. FU Y.J., ZHANG J.L., HOU X.Q. Comparative Analysis of Fungi Diversity in Rhizospheric and Non-rhizospheric Soil from *Cypripedium macranthum* Estimated via High-throughput Sequencing. *Acta Agriculturae Boreali-occidentalis Sinica*, **28** (02), 253, **2019**.
9. HUANG M., JIANG B., GAO D.Z., XIE B., WEI S.Y. Community characteristics of endophytic fungi and soil fungi in the *Calanthe sieboldii*. *Ecological Science*, **41** (4), 112, **2022**.
10. ZHANG J., HOU X.Q., FU Y.J. Rhizospheric Bacteria Diversity of *Cypripedium macranthum* Estimated via High Throughput Sequencing. *Southwest China Journal of Agricultural Sciences*, **30** (4), 811, **2017**.
11. JIANG Y.L., CHEN X.H., MIAO Q., QU B. Difference in fungal communities between in roots and in root-associated soil of nine orchids in Liaoning, China. *Chinese Journal of Plant Ecology*, **43**, 1079, **2019**.
12. HUANG M., JIANG B., GAO D.Z. Community characteristics of endophytic fungi and soil fungi in the *Calanthe sieboldii*. *Ecological Science*, **41** (4), 112, **2022**.

13. PRASHAR P., KAPOOR N., SACHDEVA S. Rhizosphere: its structure, bacterial diversity and significance. *Reviews in Environmental Science and Bio/Technology*, **13**, 63, **2014**.
14. LU Y., ROSENCANTZ D., LIESACK W. Structure and activity of bacterial community inhabiting rice roots and the rhizosphere. *Environmental microbiology*, **8** (8) 1351, **2006**.
15. ZHU Y.P., LI S.Q., LIU Y.Q., JU W.P., SHI B.L., TIAN H.S. A preliminary report on the regional survey of the distribution of the population of *Spodoptera sinensis*. *Liaoning Forestry Science and Technology*, **1**, 23, **2004**.
16. LV R. The Distribution of *Diplandrorchis sinica* S.C.Chen and the New Discovery of Phenological Observation. *Journal of Heilongjiang Vocational Institute of Ecological Engineering*, **30** (1) 16, **2017**.
17. ZHANG H., ZHANG H.S., FU J., HE H. Habitat characteristics of endangered plant *Diplandrorchis sinica* population in Mt. Laotudingzi of eastern Liaoning province. *Journal of Liaoning Normal University (Natural Science Edition)*, **37** (3), 389, **2014**.
18. ZHANG L.J., ZHOU Q., JU W.P., LI H.Y., WANG Y.J., ZHOU Y.B. Investigation on habitat of rare and endangered plants of *Diplandrorchis*. *Northern Horticulture*, (16), 111, **2010**.
19. QU B., JIANG Y.L., ZHANG Y.Z.H., ZHANG L.J. *Diplandrorchis Sinica* Mycorrhizal Fungi Optimization of Culture Conditions. *Hubei Agricultural Sciences*, **55** (22), 5798, **2016**.
20. CAPORASO J.G., LAUBER C.L., WALTERS W.A., BERG-LYONS D., LOZUPONE C.A., TURNBAUGH P.J., FIERER N., KNIGHT R. Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *Proceedings of the National Academy of Sciences*, **108** (Suppl 1), 4516, **2011**.
21. CAPORASO J.G., LAUBER C.L., WALTERS W.A., BERG-LYONS D., HUNTLEY J., FIERER N., OWENS S.M., BETLEY J., FRASER L., BAUER M. Ultra-high-throughput microbial community analysis on the IlluminaHiSeq and MiSeq platforms. *Isme Journal Multidisciplinary Journal of Microbial Ecology*, **6** (8), 1621, **2012**.
22. EDGAR R.C., HAAS B.J., CLEMENTE J.C., QUINCE C., KNIGHT R. UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics*, **27** (16), 2194, **2011**.
23. WANG Q., GARRITY G.M., TIEDJE J.M., COLE J.R. Naive bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Applied and environmental microbiology*, **73** (16), 5261, **2007**.
24. YANG H.J., WANG Q., WAN Z.X., ZHANG Z.Y., CHEN D.D., TAN J. Structure and diversity of microbial communities in the rhizosphere and non-rhizosphere soil in areas with invasive *Solidago canadensis* L. *Journal of Biosafety*, **30** (4), 235, **2021**.
25. GUO H., TANG W.P. Enzyme activity and microbial community diversity in rhizosphere and non-rhizosphere of *Larix principisrupprechtii*. *Ecology and Environmental Sciences*, **29** (11), 2163, **2020**.
26. ZHENG M.X., CHEN H., ZHU Y.J., SU H.L. Microbial diversity in rhizosphere and non-rhizosphere soils of *Paris polyphylla var. chinensis* Plants. *Fujian Journal of Agricultural Sciences*, **35** (12), 1357, **2020**.
27. TONG Q., CHEN M.T., LONG J.Q., TONG F.P., LI G., LIU Z.H., CHEN R., WU M. Research on soil nutrient characteristics at rhizosphere and non-rhizosphere for different age groups of *Choerospondias axillaris*. *Journal of Central South University of Forestry and Technology*, **39** (12), 108, **2019**.
28. ZHANG J., HOU X.Q., FU Y.J. Analysis of bacterial diversity in rhizosphere soil of *Cypripedium grandiflorum* based on high-throughput sequencing. *Journal of Southwest Agriculture*, **30** (4), 811, **2017**.
29. WANG X.W., HONG Z.H., LIU A.R., LUO L.X. Diversity analysis of soil bacterial community structure in vineyards based on fluorescence quantitative PCR and high-throughput sequencing technology. *Wine Technology*, **269** (11), 28, **2016**.
30. CHEN Y.X., DENG J.J., ZHOU Y.B., YIN Y., WEI Y.W., BAI X.J., ZHU W.X. Characteristics of soil microbial community in natural secondary forest of *Quercus mongolica* and its relationship with soil physical and chemical properties. *Journal of Shenyang Agricultural University*, **49** (4), 409, **2018**.
31. LIU L., SUN L., ZHANG R.Y., YAO N., LI L.B. Diversity of Indoleacetic Acid producing Endophytic Bacteria from the Roots of *Cymbidium album*. *Biodiversity Science*, **18** (2), 182, **2010**.
32. HOU T. Diversity of mycorrhizal fungi in dominant orchids in Huanglong Valley [D] Beijing Forestry University, 67, **2010**.
33. CHU X.L., YANG B., GAO L., LI H.L., HU L., YANG N. Species diversity of culturable bacteria isolated from the roots of *Cymbidium hybridum*. *Plant Science Journal*, **28** (2), 199, **2010**.
34. BHATTACHARJEE A., DUBEY S., SHARMA S. Storage of soil microbiome for application in sustainable agriculture: prospects and challenges [J]. *Environmental Science and Pollution Research International*, **29** (3), 3171, **2022**.
35. AHKAMI A.H., WHITE R.A., HANDAKUMBURA P.P. Enhancing sustainable plant ecosystem productivity[J]. *Rhizosphere engineering*, (3), 233, **2017**.
36. PENG J G, GONG J Y, FAN Y.H., ZHANG Y.F., BAI Q.Y., WANG Y.M., XIE L.J. Diversity of soil microbial communities in the inter-root and non-inter-root areas of *Rhododendron brevicornis*[J]. *Forestry Science*, **58** (2), 89, **2022**.
37. JING H. Mechanisms of nitrogen addition affecting greenhouse gas release from inter-root and non-inter-root soils of *Pinus sylvestris*[J]. *Doctoral Dissertation*, Northwest Agriculture and Forestry University, **2021**.
38. GAN D., FENG J., HAN, M., ZENG H., ZHU B. Rhizosphere effects of woody plants on soil biogeochemical processes: A meta-analysis. *Soil Biology and Biochemistry*, **160**, 108310, **2021**.
39. WU A.L., JIAO X.Y., WANG J.S., DONG E.W., GUO J., WANG L.G., SUN A.Q., HU H.W. Sorghum rhizosphere effects reduced soil bacterial diversity by recruiting specific bacterial species under low nitrogen stress. *Science of the Total Environment*, **770**, 144742, **2021**.
40. XIE F.C., ZHANG G.Y., ZHENG Q.J., LIU K.M., YIN X.J., SUN X.Y., SAUD S., SHI Z.J., YUAN R.L., DENG W.J., ZHANG L., CUI G., CHEN Y.J. Beneficial effects of mixing Kentucky bluegrass with red fescue via plant-soil interactions in black soil of Northeast China. *Frontiers in Microbiology*, **11**, 556118, **2020**.
41. HU H.Y., LI H., HAO M.M., REN Y.N., ZHANG M.K., LIU R.Y., ZHANG Y., LI G., CHEN J.S., NING T.Y., KUZYAKOV Y. Nitrogen fixation and crop productivity enhancements codriven by intercrop root exudates and key rhizosphere bacteria. *Journal of Applied Ecology*, **58** (10), 2243, **2021**.
42. YIN C.T., VARGAS J.M.C, SCHLATTER D.C., HAGERTY C.H., HULBERT S.H., PAULITZ T.C. Rhizosphere community selection reveals bacteria associated with reduced root disease. *Microbiome*, **9** (1), 86, **2021**.