

Original Research

New Findings of Efficient Low-Temperature Complex Cellulose-Degrading Bacteria: a Comprehensive Study

Haoran Ma[#], Baiting Zhang[#], Chunrui Liu[#], Haiyang You[#], Siyuan Li[#], Junyan Chu[#],
Lu Jiang[#], Xuesong Wang[#], Zhihua Liu^{*}, Yingjie Dai^{**}

College of Resources and Environment, Northeast Agricultural University,
No. 600 Changjiang Road, Xiangfang District, Harbin 150030, China

Received: 23 May 2024

Accepted: 22 July 2024

Abstract

Natural compost was used as a source of bacteria and cellulose was the only carbon source. After initial screening, secondary screening, and molecular biology identification, X11 belonged to the *Pseudomonas sp.*, Z3 and Z25 belonged to the *Cladosporium sp.*, and F74 belonged to the *Streptomyces sp.* Based on the mixed study of three types of cellulose-degrading bacteria, it was found that the filter paper degrading enzyme activity of the X11F74 was 0.327 IU/g, the carboxymethylcellulose (CMC) degrading enzyme activity was 1.312 IU/g, and the microcrystalline cellulose enzyme activity was 0.289 IU/g, significantly superior to other combinations, demonstrating synergistic effects. The optimal conditions for enzyme production by X11F74 CMC were determined by a one-way variable test to be an incubation time of 5 d, an initial pH of the medium of 6, an incubation temperature of 20°C, and an inoculum of 5%. Finally, the mechanism of cellulose degradation by X11F74 was analyzed.

Keywords: characterization, complex cellulose-degrading bacteria, degradation mechanisms, low temperature, screening

Introduction

Composting, recognized as an economical and effective organic waste treatment method, garners considerable attention for its reliance on microbial and enzymatic actions to decompose unstable solid organic matter into stable humus [1, 2]. Presently, composting techniques are categorized into two groups: natural composting and artificial composting

[3, 4]. Among them, natural composting is the more traditional and common method, which makes use of the decomposition of microorganisms in nature to convert organic wastes into organic fertilizers [5]. However, the natural composting process necessitates an extended fermentation period. Cow dung as a composting raw material [6] not only supplies a substantial quantity of nutrients and organic matter for composting [7, 8] but also enhances plant productivity [9]. Simultaneously, it efficiently mitigates emissions of GHGs, NH₃, VOCs, and other gas emissions [10, 11]. Straw as an auxiliary feedstock for composting can not only provide a source of compost organic matter carbon for composting,

[#]Equal contribution

^{*}e-mail: zhihua-liu@neau.edu.cn

^{**}e-mail: dai5188@hotmail.com

regulate the C/N ratio, enhance compost porosity [12], and regulate the structure of microbial communities [13], but also contribute to the reduction of environmental pollutants like CH_4 and N_2O [14], and increase the added value of agricultural waste in economic, environmental, or social terms [15]. The annual production of straw in China is about 7.06×10^8 t, and that of livestock manure is about 2×10^9 t [16]. Both straw and livestock manure contain a large amount of cellulose, hemicellulose, and lignin, representing abundant biomass resources in nature [17]. Cellulase can be used for the development of this biomass resource. Cellulase is a general term for a group of enzymes that can degrade cellulose into simple sugars, rather than referring to one enzyme alone. It primarily comprises endoglucanase, exoglucanase, and β -glucosidase. Cellulose-degrading bacteria can synthesize these enzymes and utilize them for cellulose degradation. However, the close arrangement of atoms in cellulose crystals renders it impermeable to enzymes and water molecules [18]. At the same time, the lignin in the plant cell wall is connected with cellulose, relying on hydrogen bonding, which hinders the contact between cellulose and enzymes, although lignin does not damage the enzymatic reaction of cellulose, which leads to a longer fermentation time in natural composting techniques [19]. Hence, we need more efficient cellulose-degrading bacteria to enhance degradation efficiency.

Composite fungicide generally refers to a system formed by mixing two or more single bacteria. Research has demonstrated that synergy exists among microorganisms during the composting process, with a single strain having a lower role in accelerating cellulose degradation than the combined effect of a composite microflora [20]. Moreover, the composite microflora demonstrated the capability to enhance the diversity of cellulose-degrading genes and communities within the compost [21], stimulate the potential functions and interactions of the flora, and augment the cellulose-degrading bacterial degradation capacity in the original compost [22]. For the current urgent situation, this approach can take full advantage of more efficient microbial degradation systems for cellulose. Utilizing this method is a more appropriate option given the long duration of current natural composting techniques, the difficulty of converting waste into an effective resource, and the urgency for production [19, 22].

In Heilongjiang, China, a major grain-producing province, the ambient temperature is around -16°C in winter. Low ambient temperatures are a challenge for natural composting because low temperatures result in reduced microbial abundance, simplified composition, and suppressed activity, which in turn leads to decreased composting efficiency [23, 24]. In order to ensure the normal operation of composting, some compost heating methods have been extensively studied, such as natural gas heating, biogas heating [25], electric heating, and solar heating [26, 27]. These heating techniques aim to enhance microbial activity in the compost at low temperatures [23]. However, these methods are

costly and energy-intensive, limiting their potential effectiveness in practical composting applications [28, 29]. Previous research indicates that cold-adapted microorganisms have their own adaptive ability to cope with cold stress and can maintain high activity in low-temperature environments [30–32]. Therefore, screening for cold-adapted microorganisms is of high practical value.

In this study, natural compost was used as the bacterial source and cellulose as the only carbon source, and the low-temperature efficient cellulose-degrading bacteria were initially screened, and antagonistic experiments and one-way variable experiments were used to further study the interaction system that exists among the composite strains under low-temperature conditions and to ultimately determine the mixed flora with the best cellulose-degrading effect and its optimal enzyme-producing conditions. These findings provide a theoretical basis and technical support for the optimization of the low-temperature composting process of straw and livestock manure at a later stage, so as to better solve the problems of pollution and resource wastage of the accumulation of straw and livestock manure under low-temperature conditions.

Materials and Methods

Sample Collection and Processing

The samples were taken from a natural compost pile in Xinyi Village, Harbin City, the pile (2.8 m long, 1 m wide, and 0.8 m high), with the main components of which are the mixed fermentation products of straw and animal manure. In the pre-composting period (2 d of fermentation), 2 points were randomly selected from the upper, middle, and lower layers of the compost (10, 30, and 60 cm) for sampling, totaling 6 samples, and the obtained compost fresh samples were packed into pre-inactivated bacteriological sealing bags and then retrieved for use after being wrapped. The six samples were mixed well to obtain the original sample to be tested.

Isolation and Purification of Cellulose-Degrading Bacterial Strains

A direct isolation method was employed to obtain strains with the capability to utilize cellulose as the sole carbon source [33]. As shown in Fig. 1a), 10 g of the original sample was placed into a 250 mL triangular flask containing small glass beads, 100 mL of sterilized distilled water was added, and the suspension was shaken at 20°C for 30 min at 120 r/min, so that the strains were completely free and evenly distributed in the water. Take 1 mL of the suspension and add it to a triangular flask containing 100 mL of Hutchison's liquid medium and incubate it at 20°C . After the filter paper disintegrated, the culture was shaken at 20°C




Fig. 1. Schematic illustration for the cellulose-degrading bacteria screening process via two steps: a) Isolation and purification of cellulose-degrading strains. b) Screening and conservation of these identified strains.

and 120 r/min for 30 min, and then transferred to fresh Hutchison's liquid medium, the inoculum amount was 10% each time, and so on for several generations, eliminating the cultures that had lost the ability to decompose and were unstable. The culture solution was taken from the culture, and under the condition of aseptic operation, the culture was picked from the broken filter paper strip, transferred to fresh CMC liquid medium, and incubated in constant temperature oscillation at 20°C and 120 r/min for 2-5 d. Under the condition of aseptic operation, the bacterial stock solution was diluted in accordance with the concentration gradient of 10⁻⁴-10⁻⁹, and the obtained dilution was spread on the CMC medium plate. Under the condition of aseptic operation, the bacterial stock solution was diluted according to different concentration gradients of 10⁻⁴-10⁻⁹, and the obtained dilutions were spread on CMC medium plates, and the plates were incubated at 20°C for 2-5 d to obtain single colonies. The single colonies were repeatedly streaked and purified until a pure strain was obtained.

Hutchison liquid medium: 1 g KH₂PO₄, 0.01 g FeCl₃, 0.3 g MgSO₄·7H₂O, 2.5 g NaNO₃, 0.1 g CaCl₂, 0.1 g NaCl, 1 000 mL H₂O, pH 7.0-7.4.

Carboxymethyl cellulose medium: 15 g CMC-Na, 1 g NH₄NO₃, 1 g yeast extract, 0.5 g MgSO₄·7H₂O, 1 g KH₂PO₄, 20 g agar, 1 000 mL H₂O.

Screening and Preservation of Cellulose-Degrading Bacterial Strains

The isolated and purified strains underwent initial screening using the Congo red staining method. As shown in Fig. 1b), the single bacterial spots obtained after isolation and purification were inoculated on CMC medium, and the colonies were grown after 2-5 d of incubation at 20°C. An appropriate amount of Congo Red staining solution was added to the plate and stained for 1 h. The staining solution was discarded and washed for 1 h with an appropriate amount of NaCl solution. The presence or absence of a hydrolysis circle was used to determine whether it was a cellulose-degrading bacteria. Based on the diameter of the hydrolysis circle, strains demonstrating superior cellulose degradation were initially selected. By regulating the temperature, the strains were screened for their ability to grow even under low temperature conditions. The strains obtained from the initial screening were streaked and cultured on CMC medium for 2-5 d. The temperature gradient of 3°C was used for cooling, and the growth of the strains and the number of survivors were observed after each cooling to finally obtain the lowest temperature for the growth of the strains. The total enzyme activity, endoglucanase activity, and exoglucanase activity were assessed through the dinitrosalicylic acid (DNS)

colorimetric assay. The filter paper degrading enzyme activity was employed to indicate the total enzyme activity, CMC degrading enzyme activity represented the endoglucanase activity, and microcrystalline cellulose enzyme activity denoted the exoglucanase activity [34-37]. Under the condition of aseptic operation, the strain obtained by re-screening was inoculated on the slant medium by using the scribing method, and after growing colonies for 2-5 d at 20°C, it was placed in the refrigerator at 4°C for storage. In order to maintain the activity of the strain, it was usually transferred once every 1-2 months.

Molecular Biology Identification

Extraction of strain DNA: DNA is extracted from the organism and subsequently processed through standard molecular biology experimental procedures.

PCR amplification and sequencing: the strains that have been isolated were sequenced with primers 27f: 5'-AGAGTTTGATCCTGGCTCAG-3' and 1492R: 5'-GGTTACCTTGTTACGACTT-3' (for bacteria and actinomycetes), NS1: 5'-GTAGTCATATGCTTGTCTC-3' and NS8: 5'-TCCGCAGGTTACCTACGGA-3' (for fungi) [38] and amplified by PCR. The reaction system was: 1 µL of primer 27f (10 µM), 1 µL of primer 1492R (10 µM), 5 µL of 10×EasyTaq Buffer, 4 µL of dNTPs (2.5 mM), 0.5 µL of EasyTaq DNA Polymerase, 1 µL of DNA buffer, and 37.5 µL of sterilized deionized water (for bacterial vs. actinobacteria). Primer NS1 (10 µM) 1 µL, primer NS8 (10 µM) 1 µL, 10×EasyTaq Buffer 5 µL, dNTPs (2.5 mM) 4 µL, EasyTaq DNA Polymerase 0.5 µL, DNA Buffer 1 µL, and sterilized deionized water 37.5 µL (for fungi). The PCR amplification procedure was 94. The PCR amplification program was as follows: pre-denaturation at 94°C for 10 min, start the cycle, denaturation at 94°C for 30 s, annealing at 50°C for 45 s, extension at 72°C for 100 s, 35 cycles, and stabilization at 72°C for 10 min (for bacteria and actinomycetes). Pre-denaturation at 94°C for 5 min, start the cycle, denaturation at 94°C for 40 s, annealing at 60°C for 60 s, extension at 72°C for 90 s, 30 cycles, and stabilization at 72°C for 10 min (for fungi). The PCR amplified products were sent to UW Genome Technology Co. for sequencing. The measured results were compared with the data on Genbank to find the homologous sequences and analyzed.

Cultivation Studies of Mixed Bacteria and Determination of Enzyme Activity

The antagonism test was carried out on the identified dominant strains. Two different bacteria were drawn parallel lines on the CMC plate medium and incubated at 20°C for 2-5 d to observe whether there was any antagonism phenomenon. The turbidimetric method was used to investigate the growth of a mixture of bacteria by taking advantage of light impermeability of the bacteria to light. The bacterial solution to be tested

was diluted 10-fold, and shaken well, and the culture solution without inoculum was used as a blank control, and the optical density value was determined using a spectrophotometer at a wavelength of 560 nm. The strains without antagonism were mixed, and the enzyme activity was determined in the same way as for single bacteria. The dominant mixed bacteria were screened by determining enzyme vigor.

Optimization of Enzyme Production Conditions for Mixed Bacteria

To investigate the effect of incubation time on cellulase enzyme activity, the screened dominant mixed bacteria were inoculated into the liquid medium, and the CMC degrading enzyme activity was measured every 24 h, and consecutively until the enzyme activity began to decline, to determine the period of maximum enzyme activity as well as the change of the enzyme activity during the whole growth process; to investigate the effect of initial pH on cellulase activity, on the basis of the period of maximum enzyme activity has been determined, set the initial pH range of 3.0-11.0, determine the CMC degrading enzyme activity of the mixed bacteria, and determine the optimal enzyme-producing initial pH; to investigate the effect of incubation temperature on cellulase activity, on the basis of the period of maximum enzyme activity has been determined and the optimal enzyme-producing initial pH, set the incubation temperature. The range was 5-15°C, and the CMC degrading enzyme activity of the mixed bacteria was determined to determine the optimal enzyme-producing temperature; the effect of inoculum amount on cellulase enzyme activity, based on the period of maximum enzyme activity, the optimal initial pH for enzyme production and the optimal enzyme-producing temperature had been determined, the range of inoculum amount was set from 0.5 to 7%, and the CMC degrading enzyme activity of the mixed bacteria was measured to determine the optimal inoculum amount for enzyme production.

Investigation of Degradation by Mixed Bacteria

The weight loss method was used to determine the degradation rate of the mixed bacteria. The test of degradation of filter paper by cellulose degrading bacteria was carried out by drying the filter paper to a constant weight and adding it to the culture medium, washing and drying the filter paper at the end of the test, recording the constant weight of the filter paper before and after the test, finding the difference in weights, calculating the rate of weight loss, and studying the degradation of cellulose by the mixed bacteria through the rate of weight loss of the filter paper. Under the optimal conditions for enzyme production, the mixed bacteria were inoculated into 100 mL of PPCS medium (containing 0.5 g of filter paper) for static incubation with three replicates. The culture solution underwent

centrifugation at 6000 r/min for 5 min, followed by removal and washing of the supernatant. Subsequently, it was centrifuged again at 6000 r/min for 5 min, washed with a hydrochloric acid and nitric acid mixture to eliminate bacterial residues, further washed with water, and centrifuged. The resulting sample was then dried at 105°C and weighed. The following formula was used to calculate the degradation rate [39]

$$\text{Degradation rate} = (\text{weight of filter paper before treatment} - \text{weight of filter paper after treatment}) / \text{weight of filter paper before treatment} \times 100\% \quad (1)$$

Results and Discussion

Enzyme Activity Analysis of Low-Temperature Cellulose-Degrading Bacteria

Through isolation, purification, as well as primary and secondary screening, 12 strains of low-temperature cellulose-degrading microorganisms were finally obtained, named X11, X36, X42 (bacteria), Z2, Z3, Z21,

Z25, Z27 (fungi), and F5, F14, F22, F74 (actinomycetes). The strains were inoculated into the corresponding liquid media according to class, and the effect of each strain on the enzyme activities of filter paper degrading enzyme, CMC degrading enzyme, and microcrystalline cellulase was different, as presented in Fig. 2. Among the three types of strains, the highest filter paper degrading enzyme activities were found in X11, Z3, and F74, with corresponding enzyme activities of 0.235 IU/g, 0.269 IU/g, and 0.321 IU/g. The highest CMC degrading enzyme activities were found in X11, Z3, and F74, with corresponding enzyme activities of 0.541 IU/g, 0.564 IU/g, and 0.747 IU/g. The highest microcrystalline cellulase enzyme activities were also found in X11, Z3, and F74, with corresponding enzyme activities of 0.273 IU/g, 0.271 IU/g, and 0.262 IU/g. Furthermore, the filter paper degrading enzyme activity of Z25 was 0.232 IU/g, the CMC degrading enzyme activity was 0.532 IU/g, and the microcrystalline cellulase enzyme activity was 0.261 IU/g, which was very similarly different from the three enzyme activities of Z3. Therefore, bacteria X11, fungi Z3 and Z25, and actinomycetes F74 were finally identified as candidate cryogenic strains for cellulose degradation. The four




Fig. 2. Results of filter paper enzyme, CMC enzyme, microcrystalline cellulase enzyme activity of strains, and differences in enzyme activity.




Fig. 3. Congo red staining results.

cellulose-degrading bacteria obtained by Congo red staining are shown in Fig. 3.

Molecular Biology Identification of Low-Temperature Cellulose-Degrading Bacteria

The sequence lengths of the amplified products of Bacteria X11, Fungi Z3, Fungi Z25, and Actinobacteria F74 were 1393 bp, 1686 bp, 568 bp, and 1408 bp, respectively, which were consistent with the expected size of the amplified fragments. The measured sequences were compared in the Genbank database to find their homologous sequences, and the results are shown in Table S1. Strain *Pseudomonas* sp. *sgn 46* was found to share up to 100% homology with bacterium X11, and strain *Cladosporium cladosporioides* strain *TXJ13021* was found to share up to 100% homology with fungus Z3. Strains of *Cladosporium* sp. *MUT 4306*, *Cladosporium cladosporioides* strain *M61*, *Cladosporium cladosporioides* strain *CFP14*, and *Cladosporium* sp. *CHTAE11* showed up to 100% homology with the fungus Z25 with up to 100% homology. Various strains of *Streptomyces* spp. documented in the GenBank database showed more than 99% homology with Actinomyces F74. The corresponding phylogenetic trees

based on this comparison are presented in Fig. 4. It was finally determined that Bacteria X11 belongs to the *Pseudomonas* sp., Fungi Z3 and Fungi Z25 belong to the *Cladosporium* sp., and Actinobacterium F74 belongs to the *Streptomyces* sp.

Mixed Antagonistic Effects of the Candidate Strains

Antagonism tests on the four strains identified revealed antagonistic reactions between strains Z3 and F74, but not between the other strains (Table 1). Seven groups of mixed bacteria were finally obtained as X11F74, Z25F74, X11Z3, X11Z25, Z3Z25, X11Z3Z25, and X11Z25F74 (Fig. S1). Among them, combination X11Z25F74 reached the peak of growth on day 4 and had the fastest growth among all the combinations, while combination X11Z25 was the slowest growth among all the combinations, where combination X11Z25Z3 had the largest growth OD of 0.421 and combination X11Z25F74 had the smallest growth OD of 0.298. In the later stages of incubation, due to nutrient deprivation, the bacteria begin to autolyse and the turbidity of the culture solution begins to decrease.




Fig. 4. Phylogenetic tree of strains (A) X11 bacteria (B) Z3 fungi (C) Z25 fungi (D) F74 actinomycete.

Table 1. The antagonism relationship between strains.

Strain	X11	Z3	Z25	F74
X11		-	-	-
Z3	-		-	+
Z25	-	-		-
F74	-	-	-	

Note: „+” indicates the presence of antagonism, „-” means the absence of antagonism.

Differences in Enzyme Activity of Mixed Bacteria

From Fig. 2, comparing the enzyme activities of different mixed bacteria, it can be found that X11F74 has the best promotion effect on filter paper degrading enzyme, CMC degrading enzyme, and microcrystalline cellulase, with the highest enzyme activity of 0.327 IU/g, 1.312 IU/g, and 0.289 IU/g, which indicated that the degradation of cellulose was optimally degraded by X11F74. All three enzyme activities of the mixed

bacterium X11F74 were greater than those of any of the single bacteria in its fractions, suggesting a synergistic effect of bacteria X11 and F74. It was inferred that the mixed culture of strains could improve the enzyme activity, and the advantage of mixed bacteria for cellulose degradation over single bacteria was obvious. Combined with the above analysis, the mixed bacterium X11F74 is the advantageous low-temperature bacterial group for cellulose degradation.

Study on the Optimal Enzyme Production Conditions of Candidate Colony X11F74

The effect of single factors on the optimal enzyme production conditions of X11F74 is shown in Fig. 5. With different incubation times, X11F74 produced the best enzyme at 5 d of incubation with an enzyme activity of 1.378 IU/g. With different pH, X11F74 produced the best enzyme at pH 6 with an enzyme activity of 1.462 IU/g. The CMC degrading enzyme activity was more stable in the range of 6.0 to 9.2 compared to pH less than 6, suggesting that X11F74 is acid-sensitive and somewhat alkali-resistant, which may be due to the resistance of proteases isolated from




Fig. 5. Effect of incubation time, initial pH, incubation temperature, and amount of inoculum on the enzyme-producing activity of the mixed bacterium X11F74. Samples 1-30 represent groups of treatments studied for changes in CMC enzyme activity.

Table 2. Degradation of filter paper by mixed bacteria.

Sample	Before processing (g)	After processing (g)	Weightlessness (g)	Weight loss rate (%)
1	0.5007	0.3860	0.1147	22.91
2	0.5010	0.3864	0.1146	22.87
3	0.5006	0.3868	0.1138	22.73

Streptomyces and *Pseudomonas* spp. to alkaline stress [40, 41]; with different temperatures, X11F74 produced the best enzyme at 20°C with an enzyme activity of 1.462 IU/g; the best enzyme production was achieved by X11F74 at an inoculum level of 5% with an enzyme activity of 1.477 IU/g at different inoculum levels. Due to the excessive addition of bacteria, the nutrients within the medium began to be scarce, affecting the normal growth of the bacteria, and their enzyme activity instead decreased. In summary, the highest enzyme activity of the mixed bacterium X11F74 was observed under the conditions of a 5 d incubation time, an initial pH of 6, an incubation temperature of 20°C, and an inoculum volume of 5%. Under the optimal enzyme-producing conditions, it was inoculated in the liquid medium with filter paper as the only carbon source to explore its practical application effect under the optimal conditions, and the results are shown in Table 2, and the degradation rate of filter paper could reach 22.84%. Fig. 6a) shows the correlation matrix between the enzyme-producing activity of the mixed bacterium X11F74 and the environmental factors, and the results show that the enzyme-producing activity of the mixed bacterium X11F74 has the strongest correlation with the time factor. Meanwhile, this study screened about 60 pieces of literature about low-temperature cellulose-degrading bacteria in recent years and investigated the correlation between enzyme activity and environmental

factors, and the results were shown in Fig. 6b), which found that the microbial enzyme-producing activity was most strongly correlated with the time factor, which verified the results of the correlation matrix study of this experiment. This provides a theoretical basis for the practical application of the mixed bacterium X11F74, which can still better improve the composting efficiency by reasonably applying the key factor of good incubation time when other conditions are difficult to fully satisfy.

Discussion

In this study, a large amount of literature on microbial enzyme production in recent years was reviewed, and 40 papers were selected and statistically analyzed through a box plot as shown in Fig. S2. Most of the optimal temperatures for microbial enzyme production were concentrated between 32°C and 43°C, and there were fewer studies related to low-temperature cellulose-degrading bacteria, which highlighted the urgency of studying the enzyme production of low-temperature degrading bacteria. Therefore, this study involved the isolation of four strains of low-temperature cellulose-degrading bacteria with high degradation ability from mixed compost derived from straw and livestock manure in cold regions. We hypothesize synergistic interactions among diverse strains during composting, collectively




Fig. 6. a) Pearson correlation matrix between enzyme-producing activity of the hybrid bacterium X11F74 and environmental factors. b) Pearson correlation matrix between low-temperature cellulose-degrading bacteria and environmental factors.

contributing to the degradation of cellulosic materials, and achieving efficient cellulose degradation. Following the initial screening of the four strains, an antagonism test was conducted, and the non-antagonistic single strains were mixed. The growth and enzyme activity of the resultant mixed bacteria were studied, revealing mixed bacteria X11F74 produced the best enzyme effect with more growth and faster growth. A comparison of the enzyme activities of the single bacteria with those of the mixed bacteria X11F74 revealed that all three enzyme activities of the mixed bacteria X11F74 were greater than those of any of the single bacteria in its components. The important effect of synergistic interactions between microorganisms in composting on cellulose degradation has been demonstrated in previous studies. In a study of experimental composting of corn stover and cattle manure by Meng et al. [42], it was found that the degradation rate of cellulose in compost was significantly increased by the synergistic interaction between microorganisms. Han et al. [43] found that β -glucosidase-producing bacteria and fungi synergistically drive cellulose degradation and increase the degradation rate during corn stover composting. Therefore, we conclude that the enzyme activity between both Bacteria X11 and Actinobacteria F74 is enhanced through synergistic action, which effectively improves cellulose degradation efficiency. The optimal conditions for enzyme production by the mixed bacterium X11F74 were found to be 5 d of incubation time, initial pH of the medium at 6, incubation temperature at 20°C, and an inoculum size of 5% by a one-factor variant test, and the degradation rate of the mixed bacterium under the optimal conditions was 22.8 %. In addition, based on the status of change in the CMC degrading enzyme activity of the mixed bacteria, it can be inferred that the mixed bacteria belong to the cold-resistant rather than

cold-loving bacteria, which are sensitive to acidity and have a certain degree of alkali resistance.

Taken together, the mechanism of cellulose degradation by the mixed bacteria X11F74 was investigated. Mixed bacterium X11F74 increased the activity of secreted cellulase through synergistic action, and cellulase was used to hydrolyze cellulose. The hydrolysis process of cellulase is a joint action of endoglucanase, exoglucanase, and β -glucosidase. Firstly, endoglucanase breaks down the β -1,4 glycosidic bond inside the non-crystalline region of cellulose molecule into short β -glucooligosaccharide, and then exoglucanase reacts with its non-reducing end to decompose the β -oligosaccharides into fibrous disaccharides in disaccharide units; then, endoglucanase and exoglucanase carry out synergistic action to decompose partially degraded cellulose to generate fibrous disaccharides, and fibrous trisaccharides, and other oligosaccharides; finally, it is broken down into glucose under the action of β -glucosidase to complete the hydrolysis process. The degradation mechanism is presented in Fig. 7.

The optimum growth temperature of *Bacillus agave* from soil screened by Malik et al. [44] was 37°C, and the optimum pH was 7.0, in comparison to which the mixed bacterium X11F74 is better suited for cellulose decomposition at lower temperatures and higher acidity. In comparison to an *Aspergillus sydowii* isolated by Matkar et al. [45] with an optimal inoculum quantity of 10% and an optimal incubation time of 6 d, the optimal inoculum quantity of the mixed bacterium X11F74 was smaller and the optimal incubation time was less. Therefore, the mixed bacterium X11F74 obtained in this experiment was superior. In addition, cellulose-degrading bacteria are of great interest for other applications. For example, cellulose-degrading bacteria




Fig. 7. Mechanism of cellulose degradation by cellulose-degrading bacteria.

and pure cultures for the preparation of microbial fuel cells are capable of generating electricity directly from cellulose as an electron donor. This contributes to the reduction of the use of fossil fuels and to the reduction of greenhouse gases [46]. Lactic acid production using cellulolytic flora in the gut of insects saves resources and costs and protects the environment. These studies indicate that cellulose-degrading bacteria have high utilization value, and we look forward to subsequently the further excavation of the potential functions of cellulose-degrading bacteria [47-49].

Conclusions

In this investigation, natural compost served as the bacterial source, and cellulose was employed as the only carbon source. Following a sequence of domestication, isolation, purification, primary screening, rescreening, and enzyme activity studies, four strains of low-temperature cellulose-dominant degrading bacteria were ultimately isolated. Molecular biology identified Bacterium X11 as a member of the *Pseudomonas* sp., while fungi Z3 and Z25 were assigned to the genus *Cladosporium* sp., and Actinomycetes F74 were attributed to the *Streptomyces* sp. Using an antagonism assay, the strains without antagonism were amalgamated and an examination of the mixed bacterial growth and enzyme activity ensued. The resultant analysis disclosed that the filter paper degrading enzyme activity of the mixed bacteria X11F74 measured 0.327 IU/g, CMC degrading enzyme activity reached 1.312 IU/g, and microcrystalline cellulose enzyme activity registered at 0.289 IU/g. The three enzyme activities were significantly better than the other combinations and significantly higher than any of the single bacteria in their constituents, indicating the synergistic effect of X11 and F74, thus identifying the mixed bacterium X11F74 as the best degrading bacterium. Furthermore, the optimal conditions for enzyme production by the mixed bacteria were determined as follows: an incubation time of 5 d, an initial medium pH of 6, an incubation temperature of 20°C, and an inoculum of 5%. The mixed bacterium X11F74 can indeed be applied to low-temperature compost as a low-temperature efficient cellulose degrading bacterium to promote cellulose degradation. However, the existing literature lacks comprehensive insights into the specific interaction mechanisms governing mixed bacteria. We look forward to the next step of further research on the mixed bacterium from the aspect of molecular cloning and so on.

Acknowledgments

This work was Supported by the Strategic Priority Research Program of the Chinese Academy of Sciences of China (Grant No. XDA28070301). This work was

Supported by the project of the Key Laboratory of germplasm innovation and physiological ecology of grain crops in cold regions of the Ministry of Education (CXSTOP2021008).

Conflict of Interest

The authors declare no conflict of interest.

References

1. CAO Y.B., WANG X., BAI Z.H., CHADWICK D., MISSELBROOK T., SOMMER S.G., QIN W., MA L. Mitigation of ammonia, nitrous oxide and methane emissions during solid waste composting with different additives: a meta-analysis. *Journal of Cleaner Production*. **235**, 626, **2019**.
2. JIANG H.T., LI X., DAI Y.J. Phosphoric acid activation of cow dung biochar for adsorbing enrofloxacin in water: Icing on the cake. *Environmental Pollution*. **341**, 122887, **2024**.
3. ZHU P.C., LI Y.C., GAO Y.F., YIN M.Q., WU Y.X., LIU L.L., DU N., LIU J., YU X.N., WANG L. S., GUO W.H. Insight into the effect of nitrogen-rich substrates on the community structure and the co-occurrence network of thermophiles during lignocellulose-based composting. *Bioresource Technology*. **319**, 124111, **2021**.
4. ZHU P.C., WU Y.X., RU Y.N., HOU Y.H., SAN K.W., YU X.N., GUO W.H. Industrial-scale aerobic composting of livestock manures with the addition of biochar: Variation of bacterial community and antibiotic resistance genes caused by various composting stages. *Environmental Pollution*. **314**, 120270, **2022**.
5. WAQAS M., HASHIM S., HUMPHRIES U.W., AHMAD S., NOOR R., SHOAIB M., NASEEM A., HLAING P.T., LIN H.A. Composting processes for agricultural waste management: a comprehensive review. *Processes*. **11** (3), 731, **2023**.
6. REN X.N., WANG Q., CHEN X., HE Y.F., LI R.H., LI J., ZHANG Z.Q. Pathways and mechanisms of nitrogen transformation during co-composting of pig manure and diatomite. *Bioresource Technology*. **329**, 124914, **2021**.
7. LOPEA-GONZALEZ J.A., VARGAS-GARCIA M.D.C., LOPEZ M.J., SU'AREZ-ESTRELLA F., JURADO M.D.M., MORENO J. Biodiversity and succession of mycobiota associated to agricultural lignocellulosic waste-based composting. *Bioresource Technology*. **187**, 305, **2015**.
8. QIAO C.C., PENTON C.R., LIU C., TAO C.Y., DENG X.H., OU Y.N., LIU H.J., LI R. Patterns of fungal community succession triggered by C/N ratios during composting. *Journal of Hazardous Materials*. **401**, 123344, **2020**.
9. MA S.S., FANG C., SUN X.X., HAN L.J., HE X.Q., HUANG G.Q. Bacterial community succession during pig manure and wheat straw aerobic composting covered with a semi-permeable membrane under slight positive pressure. *Bioresource Technology*. **259**, 221, **2018**.
10. HERRERO M.H., HENDERSON B., HAVLÍK P., THORNTON P.K., CONANT R.T., SMITH P., WIRSENIUS S., HRISTOV A.N., GERBER P., GILL M., BUTTERBACH-BAHL K., VALIN H., GARNET

- T., STEHFEST E. Greenhouse gas mitigation potentials in the livestock sector. *Nature Climate Change*. **6** (5), 452, **2016**.
11. SPIEHS M.J., JADERBORG J.P., HALES K.E., DICOSTANZO A., CRAWFORD G.I., PARKER D.B. Effect of corn processing and wet distiller's grains with solubles on odorous volatile organic compound emissions from urine and feces of beef cattle. *Applied engineering in agriculture*. **34** (3), 591, **2018**.
 12. WANG L. Y., WANG T.F., XING Z.J., ZHANG Q.F., NIU X.H., YU Y.S., TENG Z.J., CHEN J.X., Enhanced lignocellulose degradation and composts fertility of cattle manure and wheat composting by *Bacillus* inoculation. *Journal of Environmental Chemical Engineering*. **11** (3), 109940, **2023**.
 13. SONG X.Y., YANG J.K., HUSSAIN Q., LIU X.W., ZHANG J.J., CUI D.J. Stable isotopes reveal the formation diversity of humic substances derived from different cotton straw-based materials. *Science of the total environment*. **740**, 140202, **2020**.
 14. SANTOS C., FONSECA J., AIRES A., COUTINHO J., TRINDADE H. Effect of different rates of spent coffee grounds (SCG) on composting process, gaseous emissions and quality of end-product. *Waste Manage*. **59**, 37, **2017**.
 15. DE MENDONÇA COSTA M.S.S., BERNARDI F.H., DE MENDONÇA COSTA L.A., PEREIRA D.C., LORIN H.E.F., ROZATTI M.A.T., CARNEIRO L.J. Composting as a cleaner strategy to broiler agro-industrial wastes: Selecting carbon source to optimize the process and improve the quality of the final compost. *Journal of Cleaner Production*. **142**, 2084, **2017**.
 16. LIANG F., LI B. Z., VOGT R.D., MULDER J., SONG H., CHEN J.S., GUO J.H. Straw return exacerbates soil acidification in major Chinese croplands. *Resources, Conservation and Recycling*. **198**, 107176, **2023**.
 17. PADHAN K., PATRA R.K., SETHI D., PANDA N., SAHOO S.K., PATTANAYAK S.K., SENAPATI A.K. Isolation, characterization and identification of cellulose-degrading bacteria for composting of agro-wastes. *Biomass Conversion and Biorefinery*. **1**, 15, **2023**.
 18. CHEN X.M., CHENG W.T., LI S.Z., TANG X.F., WEI Z.M. The "quality" and "quantity" of microbial species drive the degradation of cellulose during composting. *Bioresource Technology*. **320**, 124425, **2021**.
 19. CARRION C.C., NASROLLAHZADEH M., SAJJADI M., JALEH B., SOUFI G.J., IRAVANI S. Lignin, lipid, protein, hyaluronic acid, starch, cellulose, gum, pectin, alginate and chitosan-based nanomaterials for cancer nanotherapy: Challenges and opportunities. *International Journal of Biological Macromolecules*. **178**, 193, **2021**.
 20. AN X.J., ZONG Z.B., ZHANG Q.H., LI Z.M., ZHONG M., LONG H.Z., CAI C.Z., TAN X.M. Novel thermo-alkali-stable cellulase-producing *Serratia* sp. AXJ-M cooperates with *Arthrobacter* sp. AXJ-M1 to improve degradation of cellulose in papermaking black liquor. *Journal of Hazardous Materials*. **421**, 126811, **2022**.
 21. WU D., QU F.T., LI D., ZHAO Y., LI X., NIU S.J., ZHAO M.Y., QI H.S., WEI Z.M., SONG C.H. Effect of Fenton pretreatment and bacterial inoculation on cellulose-degrading genes and fungal communities during rice straw composting. *Science of The Total Environment*. **806**, 151376, **2022**.
 22. LI Y.X., KURAMAE E.E., NASIR F., WANG E., ZHANG Z.G., LI J., YAO Z.M., TIAN L., SUN Y., LUO S.Y., GUO L.L., REN G.D., TIAN C.J. Addition of cellulose degrading bacterial agents promoting keystone fungal-mediated cellulose degradation during aerobic composting: Construction the complex co-degradation system. *Bioresource Technology*. **381**, 129132, **2023**.
 23. XIE X.Y., ZHAO Y., SUN Q.H., WANG X.Q., CUI H.Y., ZHANG X., LI Y.J., WEI Z.M. A novel method for contributing to composting start-up at low temperature by inoculating cold-adapted microbial consortium. *Bioresource Technology*. **238**, 39, **2017**.
 24. YAO Y., HUANG G., AN C.J., CHEN X.J., ZHANG P., XIN X.Y., SHEN J., AGNEW J. Anaerobic digestion of livestock manure in cold regions: Technological advancements and global impacts. *Renewable and Sustainable Energy Reviews*. **119**, 109494, **2020**.
 25. XU Z.J., WU H.Z., WU M.L. Energy performance and consumption for biogas heat pump air conditioner. *Energy*. **35** (12), 5497, **2010**.
 26. JIANG Y.F., LI X.S., YAO J., WAN X., ZHANG J.X., DAI Y.J. Design and performance simulation of a distributed aerobic composting system assisted by solar PV/T heat pump. *Renewable Energy*. **196**, 547, **2022**.
 27. WANG Y.Z., MA F., ZHU T., LIU Z., MA Y.G., LI T.F., HAO L.Y. Electric heating promotes sludge composting process: Optimization of heating method through machine learning algorithms. *Bioresource Technology*. **382**, 129177, **2023**.
 28. SUN Q.H., WU D., ZHANG Z.C., ZHAO Y., XIE X.Y., WU J.Q., LIU Q., WEI Z.M. Effect of cold-adapted microbial agent inoculation on enzyme activities during composting start-up at low temperature. *Bioresource Technology*. **244**, 635, **2017**.
 29. XIE X.Y., ZHAO Y., SUN Q.H., WANG X.Q., CUI H.Y., ZHANG X., LI Y.J., WEI Z.M. A novel method for contributing to composting start-up at low temperature by inoculating cold-adapted microbial consortium. *Bioresource Technology*. **238**, 39, **2017**.
 30. DE MAAYER P., ANDERSON D., CARY C., COWAN D.A. Some like it cold: understanding the survival strategies of psychrophiles. *EMBO reports*. **15** (5), 508, **2014**.
 31. SMIRNOVA M., LOSADA C.B., AKULAVA V., ZIMMERMANN B., KOHLER A., MIAMIN U., OOSTINDIER M., SHAPAVAL V. New cold-adapted bacteria for efficient hydrolysis of feather waste at low temperature. *Bioresource Technology Reports*. **23**, 101530, **2023**.
 32. LIU X., SHAO Z.Y., WANG Y.X., LIU Y.F., WANG S.Y., GAO F., DAI Y.J. New use for *Lentinus edodes* bran biochar for tetracycline removal. *Environmental Research*. **216**, 114651, **2023**.
 33. MENSRAWY M.N., ABDEL-HAMID A.M., MOHAMED S.K., MO'MEN H. Isolation and molecular identification of cellulose/hemicellulose degrading bacteria from agricultural compost and determination of their hydrolytic potential. *South African Journal of Botany*. **149**, 617, **2022**.
 34. BAO K., WANG X.H., ZHANG W.Q., LUO J.L., NIE M., XIAO M. Screening of cellulolytic microbes and determination of cellulase activity. *Journal of Biology*. **24** (2), 56, **2007** [In Chinese].
 35. WEI Z.W., ZHAO Y.X., ZHANG M.M., ZHENG W.F. Preliminary identification and enzyme activity assay of a cellulose-degrading bacterium. *Jiangsu Agricultural Sciences*. **1**, 329, **2010** [In Chinese].
 36. LIU X.H., LU Y.R. *Environmental Biochemistry Laboratory Tutorial*. Beijing: Science Press. **2006** [In Chinese].

37. LI X., JIANG H.T., ZHU L.Y., TANG J.Q., LIU Z.H., DAI Y.J. Adsorption interactions between typical microplastics and enrofloxacin: relevant contributions to the mechanism. *Chemosphere*. **351**, 141181, **2024**.
38. HUSSAIN A.A., ABDEL-SALAM M.S., ABO-GHALIA H.H., HEGAZY W.K., HAFEZ S.S. Optimization and molecular identification of novel cellulose degrading bacteria isolated from Egyptian environment. *Journal of Genetic Engineering and Biotechnology*. **15** (1), 77, **2017**.
39. WANG W.D., CUI Z.J., WANG X.F., NIU J.L., LIU J.B., IGARASHI Y. Capability and stability of degrading rice straw of composite microbia system MC1. *Environmental Science (in Chinese)*. **26** (5), 156, **2005**.
40. GAO Y., ZHU J.W., WANG K.Y., MA Y., FANG J., LIU G. Discovery of a heterotrophic aerobic denitrification *Pseudomonas* sp. G16 and its unconventional nitrogen metabolic pathway. *Bioresource Technology*. **387**, 129670, **2023**.
41. JIANG H.T., DAI Y.J. Vitamin C modified crayfish shells biochar efficiently remove tetracycline from water: a good medicine for water restoration. *Chemosphere*. **311**, 136884, **2023**.
42. MENG L.X., XU C.X., WU F.L. Microbial co-occurrence networks driven by low-abundance microbial taxa during composting dominate lignocellulose degradation. *Science of the total environment*. **845**, 157197, **2022**.
43. HAN Y., LIU W.Y., CHANG N., SUN L., BELLO A., DENG L.T., ZHAO L.Y., EGBEAGU U.U., WANG B., ZHAO Y., ZHAO M.M., BI R.X., JONG C., XU X.H., SUN Y. Exploration of β -glucosidase-producing microorganisms community structure and key communities driving cellulose degradation during composting of pure corn straw by multi-interaction analysis. *Journal of Environmental Management*. **325**, 116694, **2023**.
44. MALIK W.A., JAVED S. Enhancement of cellulase production by cellulolytic bacteria SB125 in submerged fermentation medium and biochemical characterization of the enzyme. *International Journal of Biological Macromolecules*. **4**, 130415, **2024**.
45. MATKAR K., CHAPLA D., DIVECHA J., NIGHOJKAR A., MADAMWAR D. Production of cellulase by a newly isolated strain of *Aspergillus sydowii* and its optimization under submerged fermentation. *International Biodeterioration & Biodegradation*. **78**, 24, **2013**.
46. HASSAN S.H.A., KIM Y.S., OH S.E. Power generation from cellulose using mixed and pure cultures of cellulose-degrading bacteria in a microbial fuel cell. *Enzyme and Microbial Technology*. **51** (5), 269, **2012**.
47. SAHA S., KHATUN F., NAHIDUZZAMAN M., MAHMUD M.P., RAHMAN M.M., YASMIN S. A greener way of lactic acid production from cellulose-rich waste materials utilising cellulolytic bacteria population from insect gut. *Bioresource Technology Reports*. **18**, 101021, **2022**.
48. LIU S.B., HAN J.H., MA X.W., ZHU X.S., XIN G.R., HUANG X.C. Repeated release of CeO₂ nanoparticles altered algal responses: Growth, photosynthesis, and photosynthetic gene expression. *Eco-Environment & Health*. **2024**.
49. LIU S., HAN J., YAO L., LI H., XIN G., HO S-H, HUANG X. Integrated multilevel investigation of photosynthesis revealed the algal response distinction to differentially charged nanoplastics. *Journal of Hazardous Materials*. **475**, 134815, **2024**.

Supplementary Information

Table S. 1. Bacterial and actinomycete species were identified based on 16S rDNA gene sequences, and fungal species was identified based on 18S rDNA gene sequences.

Candidate strain	Gene bank accession number	Species	Sequence identity
X11	HF562437.1	<i>Pseudomonas</i> sp. sgn 46 partial 16S rRNA gene, isolate sgn 46	100%
F74	EU570472.1	<i>Streptomyces flavovirens</i> strain 173690 16S ribosomal RNA gene, partial sequence	99%
	NR_074561.1	<i>Streptomyces</i> sp. <i>SirexAA-E</i> strain SirexAA-E 16S ribosomal RNA, complete sequence	99%
	JQ924388.1	<i>Streptomyces flavovirens</i> strain CGMCC 4.575 clone 4 16S ribosomal RNA gene, complete sequence	99%
	NR_028988.1	<i>Streptomyces flavogriseus</i> strain CBS 101.34 16S ribosomal RNA, partial sequence	99%
	AB184834.1	<i>Streptomyces flavovirens</i> gene for 16S rRNA, partial sequence, strain: NBRC 3716	99%
	AJ494864.1	<i>Streptomyces flavogriseus</i> partial 16S rRNA gene, type strain CBS 101.34T	99%
	JQ812068.1	<i>Streptomyces</i> sp. QZGY-A11 16S ribosomal RNA gene, partial sequence	99%
	FJ486354.1	<i>Streptomyces anulatus</i> strain HBUM174206 16S ribosomal RNA gene, partial sequence	99%
Z3	EF392680.1	<i>Cladosporium cladosporioides</i> strain TXJ13021 18S ribosomal RNA gene, partial sequence	100%
Z25	KC339216.1	<i>Cladosporium</i> sp. MUT 4306 18S ribosomal RNA gene, partial sequence; internal transcribed spacer 1, 5.8S ribosomal RNA gene, and internal transcribed spacer 2, complete sequence; and 28S ribosomal RNA gene, partial sequence	100%
Candidate strain	Gene bank accession number	Species	Sequence identity
Z25	JQ936096.1	<i>Cladosporium cladosporioides</i> strain M61 18S ribosomal RNA gene, partial sequence; internal transcribed spacer 1, 5.8S ribosomal RNA gene, and internal transcribed spacer 2, complete sequence; and 28S ribosomal RNA gene, partial sequence	100%
	JQ768323.1	<i>Cladosporium cladosporioides</i> strain CFP14 18S ribosomal RNA gene, partial sequence; internal transcribed spacer 1, 5.8S ribosomal RNA gene, and internal transcribed spacer 2, complete sequence; and 28S ribosomal RNA gene, partial sequence	100%
	JF773595.1	<i>Cladosporium</i> sp. CHTAE11 18S ribosomal RNA gene, partial sequence; internal transcribed spacer 1, 5.8S ribosomal RNA gene, and internal transcribed spacer 2, complete sequence; and 28S ribosomal RNA gene, partial sequence	100%




Fig. S1. The growth of mixed bacteria.




Fig. S2. Box plot of optimum temperature for enzyme production activity of microorganisms.