

Original Research

Investigating the Influence of Specific Heavy Metals on the Haemato-Biochemical and Genotoxic Traits of Grass Carp

Sabahat Saeed^{1,2#}, Noman Waheed^{1,2**}, Muhammad Fiaz Khan², Adeeba Naseer³, Shumaila Noreen², Zeeshan Ahmad², Usama Nawaz², Shehzad Ghayyur², Xu Rifu^{1**}

¹College of Animal Sciences, Jilin Agricultural University, Changchun, Jilin, 130118 PR China

²Department of Zoology Hazara University Mansehra (21300), Khyber Pakhtunkhwa, Pakistan

³Department of Zoology, The Islamia University of Bahawalpur, Bahawalpur, Pakistan

Received: 1 May 2024

Accepted: 3 August 2024

Abstract

The current study was conducted to assess hemato-biochemical and genotoxic changes caused by Grass carp's exposure to toxic heavy metals. The fish *C. idella* was exposed to sub-lethal concentrations (1/20th of 96-h LC₅₀) of cadmium (5.7 mg/l), lead (7.6 mg/l), and mercury (0.6 mg/l), for one week. The blood measurements of the treated groups demonstrated a significant decrease ($p < 0.05$) in the levels of RBCs, hemoglobin, MCHC, WBCs, lymphocytes, and platelets, whereas there was a significant increase ($p < 0.05$) in the MCH and MCV values. The analysis of blood biochemistry unveiled a significant increase ($p < 0.05$) in metabolites (such as cholesterol, glucose, creatinine, and urea) and electrolytes (potassium and sodium). Conversely, the levels of triglycerides, serum proteins, calcium, and phosphorus exhibited a significant decline ($p < 0.05$) in comparison to the untreated group. The percentage for DNA damage for erythrocytic cells was noted for the exposed concentrations of Cd, Pb, and Hg with a significant increasing pattern ($p < 0.05$) following the sequence Pb>Hg>Cd. The results demonstrated that the Comet test is precise for determining toxicity and can be applied in environmental monitoring activities. This study concluded that excessive use of these heavy metals can negatively affect the health of fish.

Keywords: *Ctenopharyngodon idella*, heavy metals, blood cell count, biochemical analysis, DNA damage

equal contribution

*e-mail: nomanwaheed9131@gmail.com

**e-mail: poultryxu@jlau.edu.cn

Introduction

Environmental contamination has surfaced as a prominent obstacle for modern society [1], stemming from the swift industrial growth, rising energy needs, and negligent depletion of natural reserves, leading to its escalation over recent decades [2]. Both natural and human-caused sources are constantly releasing different toxic organic and inorganic contaminants into the land and aquatic ecosystem. Heavy metals count among them, and because of their toxic features, perseverance, ability to build up in living organisms, and a tendency to accumulate in the food chain, they have been highlighted as powerful biological poisons that considerably contribute to environmental contamination [3]. The poisoning by heavy metals in aquatic environments is regarded to be harmful to humans and aquatic creatures [4].

Heavy metals are metallic elements that are toxic or dangerous even in small quantities and have a relatively high density ($<5 \text{ g/cm}^3$) [5]. Heavy metals can easily infiltrate in aquatic environments and eventually reach the bodies of aquatic species [6]. In accordance to [7], cadmium (Cd), lead (Pb), and mercury (Hg) are three of the ten heavy metals that are harmful to human health. Lead is used as a production material for batteries, solder cables, anti-rust coatings, and paint-making materials. Cadmium is widely used as a pigment in the paint industry, stabilizer in the manufacture of PVC, in the manufacture of batteries and ceramics, as well as in iron coatings to be stainless steel [8]. Mercury is a non-essential metal that gets released in the surrounding environment due to natural (earth's crust) and anthropogenic (fossil fuels combustion and mining) processes [9, 10].

Fish biomarkers represent a valuable approach for monitoring the stress induced by harmful metals. They gauge cellular, bimolecular, and physiological alterations in response to intoxication resulting from exposure to heavy metals [11]. The assessment of blood and its constituents is a key component of hematological analysis, encompassing different blood indices [12]. Biochemical parameters have been widely utilized to evaluate the adverse impact of contaminants in the environment. These factors encompass the levels of electrolytes and metabolites [13]. Heavy metals are also known to be carcinogenic and potentially genotoxic. Reactive oxygen species (ROS) are produced when heavy metals stimulate oxidative stress, which can lead to DNA damage and cell death [14].

Numerous harmful metals are contaminating water and air, endangering the health of hundreds of millions of organisms around the world. Another risk to human and animal health is heavy metal contamination of food [15]. These heavy metals can bioaccumulate and have a wide range of harmful consequences on different bodily tissues and organs. These metals interfere with a variety of cellular functions, such as apoptosis, growth, proliferation, differentiation, and damage-repairing mechanisms [7]. Similar methods of action, such as hemato-biochemical reactions and genotoxic modification in grass carp, are shown by comparing the mechanisms of action of these

metals. Severe reactions following high-dose exposure result in increased DNA damage and neuropsychiatric disorders [16]. There is an abundance of information on the influence of heavy metals on the physiology of many fish species, and much of the literature focuses on the effects of certain heavy metals on a small number of physiological characteristics specific to a given fish species [17]. A thorough analysis of every heavy metal is necessary to guarantee a healthy environment for food security and safety, given the detrimental effects of heavy metals on fish and eventually humans. Therefore, it was planned to investigate more about the harmful impacts of specific heavy metals, i.e., Cd, Pb, and Hg on Grass carp physiology, particularly on blood chemistry and molecular changes at the DNA level.

The Grass carp, scientifically known as *Ctenopharyngodon idella*, is an herbivorous freshwater fish that falls under the Cyprinidae family. It is simple to upkeep in laboratory settings, holds significant commercial value in managing aquatic vegetation, and can serve as an effective indicator for appraising water quality [18]. Our study was devised to examine the hematological, biochemical, and genotoxic consequences triggered by selected heavy metals in Grass carp. As of now, this is the first research work that determined the impacts of cadmium, lead, and mercury salts on Grass Carp.

Materials and Methods

Ethical Approval

The present research was carried out in the Department of Zoology (Fisheries and Aquatic Toxicology Laboratory), Hazara University Mansehra. The ethics of animal handling were strictly followed, as suggested by the Institutional Bioethics Committee (IBC) of Hazara University Mansehra, Pakistan.

Fish Collection

The *C. idella* specimens, ranging in size from 4 to 5 inches, were collected using a manual net from Hattian fish hatchery Kamra. They were subsequently transported to the zoology laboratory at Hazara University Mansehra in an oxygenated polythene bag for subsequent examination. A 0.2% KMnO_4 solution was administered for two minutes to eliminate any external infections, followed by a rinse with tap water.

Acclimatization

Before commencing the experiment, a two-week acclimatization phase was carried out with all the fish placed in a glass aquarium with a water retaining capacity of 60 liters. All the water quality parameters were agreed to international standards of water quality suitable for freshwater fish species. Standard conditions were maintained throughout the experiment. Commercial fish

feed of 30% protein was given daily at 2% of their body weight. Oxygen was regulated through aerators, and fecal material was removed regularly to avoid ammonia toxicity. Inspection of each aquarium was done on a daily basis to find any mortality and clinical ailments.

Concentration of Heavy Metals

Analytical grade heavy metals in the form of salts were used in the present study, i.e., mercury as (HgCl_2), lead as ($\text{Pb}(\text{NO}_3)_2$), and cadmium as (CdCl_2) were purchased from (Merck, Germany) through a local distributor. For the experimentation, the $\text{LC}_{50-96\text{h}}$ value was initially ascertained for each metal. Over a 96-hour period, toxicity tests for cadmium, lead, and mercury were determined using varied concentrations of Cd (10, 30, 50, 70, 90 mg/l), Pb (20, 40, 60, 80, 100 mg/l), and Hg (02, 04, 06, 08, 10 mg/l).

Experimental Design

After the acclimatization period, fish species were then allocated to four groups (E_0 - E_3) randomly, at which a set of 30 fish were introduced in 3 replicates (10 fish/replicate). Based on 96h LC_{50} calculated value for cadmium, lead, and mercury (23.84 mg/l, 38.01 mg/l, and 3.38 mg/l), a sub-lethal concentration of 5.7 mg/l for cadmium (E_1), 7.6 mg/l for lead (E_2), and 0.6 mg/l for mercury (E_3), corresponding to $1/20^{\text{th}}$ of 96-h LC_{50} used for this study for 1 week. Group E_0 was used as a control group with non-exposed healthy fish for the entire experimentation. A semi-static system was employed, with the test solution being replaced every 24 hours. To evaluate the general health, behavior, and any alterations in response to heavy metal toxicity in fish, daily observations were made.

Hemato-Biochemical Analysis

After one week of experiments, the fish from both the control and exposed groups were carefully taken from the aquariums and then exposed to clove oil anesthesia. After that, blood was collected from the caudal vein of the fish with sterilized 2ml syringes containing heparin as an anticoagulant. The sample was taken and placed into the Tubes containing EDTA. Total erythrocyte and leukocyte count, hemoglobin, hematocrit, and platelet values were examined by using a hematology analyzer (Sysmex, Kx21). Packed cell volume (PCV) was determined through the micro-hematocrit technique. Documentation of erythrocyte indices, including mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), and mean corpuscular hemoglobin concentration (MCHC), was accomplished through a specific process [19]. The serum has been employed to assess a comprehensive range of biochemical indicators, encompassing electrolytes and metabolites. Metabolites include triglycerides, protein, glucose, cholesterol, serum creatinine, blood urea, and uric acid, and serum electrolytes such as serum calcium, phosphorus, iron, sodium, and potassium were

also identified in the bloodstream, utilizing a biochemical analyzer.[20].

Genotoxicity Assessment

The blood was analyzed by Single Cell Gel electrophoresis technique (Comet assay) for DNA damage. Comet assay performed on fish blood was adopted according to the described protocol [21]. Blood samples were tempered by utilizing saline solution. Slides were made using 10 μl of saline solution and 120 ml of low-melting agarose (0.5%) at 37°C . Slides in the lysis solution (1 mL of Triton X-100, 10 mL of DMSO, and 89 mL of lysing solution stock, pH 10.0 – stock solution: 2.5 M of NaCl, 100 mM of EDTA 100, 10 mM to 1 L of Tris) were refrigerated for 60 minutes. After 60 minutes, the slides were retained for 20 minutes at 25 V and 300 mA in a horizontal electrophoresis configuration. Slides were neutralized for approximately 15 minutes with 0.4M Tris (pH 7.5) and then stabilized in ethanol for 10 minutes. Cells without DNA damage exhibit homogeneous movement, but those with DNA damage exhibit fragments of different masses, and smaller cells move more quickly during electrophoresis, like a comet's tail.

Statistical Analysis

We analyzed the data by using statistical tools, such as SPSS (version-24). A t-test was conducted to compare the control and exposed groups, whereas a one-way ANOVA was performed to analyze all exposed groups. For creating graphs, the software GraphPad Prism (version 9) was used.

Results


This study examined the toxic effects of selected heavy metals Cd, Hg, and Pb on the well-being of *C. idella* fish by assessing its genotoxic and hemato-biochemical biomarkers. The findings of this current investigation are outlined in this study.

Hematological Responses

In a recent study, we investigated the exposure to specific heavy metals affecting the hematological parameters of the fish *C. idella*. Alterations in the hematological profiles were observed, indicated by the changes induced by exposure to the toxic substances. Specifically, the blood indices of the experimental groups displayed a significant decline ($p < 0.05$) in the levels of WBCs, RBCs, platelets, Hb, lymphocytes, and MCHC while there was a significant rise ($p < 0.05$) in the levels of MCV and MCH in *C. idella*, in comparison to the control group.

Biochemical Indices

Compared to the untreated or control group, the treated groups exhibited changes in various biological parameters






Fig. 1. Biochemical Parameters in *C. idella* exposed to selected heavy metals in contrast with control. Graphs representing hematological parameters with symbols * ($p < 0.05$) and ** ($p < 0.00000$) denotes significant deviation ($p < 0.05$) between the control and experimental groups of *C. idella*, (A) RBCs count ($10^6/\mu$ l), (B) Hemoglobin (gd/l), (C) PCV (%), (D) MCH (pg), (E) MCV (fl), (F) MCHC (g/dl), (G) Platelets (10^3), (H) WBCs count ($10^6/\mu$ l), (I) Lymphocytes (%), (J) Neutrophils (%), (K) Monocytes (%).

Table 1. Biochemical Parameters in *C. idella* exposed to selected heavy metals in contrast with control.

Biochemical Parameters	Control group	E1 Cadmium	E2 Lead	E3 Mercury
Metabolites				
Glucose (mg/dl)	32.86±2.50	64.20±3.24***a	74.0±4.02**b	39.36±2.20*c
Cholesterol (mg/dl)	92.83±2.57	133.5±3.15***a	116.8±4.60**b	154.8±4.02***c
Triglyceride(mg/dl)	153.7±3.89	136.07±3.27***a	115.9±4.26**b	116.03±3.55***c
Creatinine (mg/dl)	2.86±0.38	8.56±0.58***a	8.15±0.65 **b	8.12±0.43***c
Proteins (g/dl)	8.24±0.37	5.91±0.51***a	8.24±0.59 ^b	6.73±0.60*c
Blood Urea (mg/dl)	41.68±2.52	119.8±3.0***a	114.7±4.11***b	101.1±3.02***c
Uric Acid (mg/dl)	2.56±0.41	3.12±0.23	2.98±0.10	3.16±0.15
Electrolytes				
Sodium (mmol/l)	94.10±2.95	106.8±4.01*a	104.4±3.52*b	100.2±2.94 ^c
Potassium (mmol/l)	2.65±0.05	3.96±0.12***a	3.86±0.55*b	3.84±0.24***c
Phosphorus (mmol/l)	3.85±0.67	2.67±0.59 ^a	2.80±0.68 ^b	1.89±0.20***c
Calcium (mg/dL)	14.50±2.3	9.12±0.33*a	10.91±1.50 ^b	12.04±1.83 ^c
Iron (mg/dL)	83.87±2.16	83.50±2.89 ^a	64.77±4.41***b	79.16±2.67 ^c

Mean±S.D values for some biochemical indices with symbols * and ** denotes significant deviation ($p < 0.05$) between the experimental groups and the control group, and symbols (a,b,c) indicate a significant intergroup difference.

after exposure to specific heavy metals. There was an increase in blood glucose, cholesterol, creatinine, blood urea, and uric acid levels, alongside a decline in proteins across all treated groups in comparison to the untreated group. Additionally, changes in electrolyte values, including phosphorus, iron, calcium, potassium, and sodium were noted.

DNA Damage

The intensifying concerns about the toxic effects of heavy metals compelled scientists to investigate the genotoxic effects of selected metal ions on fish. The genotoxicity of cadmium, lead, and mercury in erythrocytes of *C. idella* was analyzed using single cell gel electrophoresis technique (Comet assay). After the exposure of *C. idella* to different concentrations of selected metal ions, a significant ($p < 0.05$) higher frequency of erythrocytic DNA damage was observed in the following trend $Pb > Hg > Cd$ in all treated groups compared to the control group. Details of DNA damage that occurred in treated groups are presented in Fig. 2 and Table 2.

Discussion

In the present study, various hematological indices demonstrated a significant decrease in the number of RBCs, Hct, and Hb in fish when exposed to different metals

compared to the control group. The distribution and binding of oxygen in fish blood are reflected in the amounts of Hct and Hb. Our results are consistent with the research of [22], who found that silver carp treated with mercuric chloride had lower hemoglobin, red blood cell counts, and HCT. Similarly, [23] noted that exposure to salts of copper and cadmium caused a drop in red blood cell count, hemoglobin, and hematocrit in *C. catla*. Zulfahmil I. et al. [24] examined a significant decline in Hb and RBC in *Chanos chanos* after being subjected to lead nitrate, which concurs with our observations. Hematocrit values typically decline when a fish experiences reduced appetite or stress. The noticeable decrease in hematocrit and hemoglobin values suggests a mechanism of hemodilution resulting from impaired osmoregulation or gill damage. Our results further align with previous research conducted by various scholars, as observed in the works of [25, 26].

The hematological indices, such as MCH, MCV, and MCHC, offer additional insights into the size, relationship, shape, and Hb content of erythrocytes. Moreover, these indices serve as standards for the morphological examination of anemia [27]. The hemoglobin to erythrocyte count ratio is represented by MCH. The MCV value can indicate the efficiency of RBC production throughout the process of erythropoiesis. Our research showed that when grass carp were exposed to particular metals, their total MCH and MCV values rose substantially, while their MCHC values declined. Parallel findings were observed in the milkfish subjected to lead nitrate [24]. Regenerative anemia brought on by




Fig. 2. DNA damage in peripheral erythrocytes of *C. idella* due to metals exposure. Representing peripheral erythrocytic cells (A) control-regular cell (B) Cd (5.77 mg/L) exposed (C) Hg (0.67 mg/l) exposed (D) Pb (7.6 mg/l) exposed.

Table. 2. DNA damage in peripheral erythrocytes of *C. idella* due to metals exposure.

Heavy metals/Concentrations (mg/l)		DNA Damage (%)			
Heavy Metals	Sub-lethal Concentrations	Type I	Type II	Type III	Type IV
Cadmium	5.77	25.6±2.34**	22.39±1.87**	28.18±1.64**	11.32±2.23**
Mercury	0.67	36.17±1.55**	28.45±2.24**	26.62±1.75**	19.86±1.42**
Lead	7.6	48.13±3.43**	41.53±2.72**	29.54±1.32**	15.63±1.73**

Summary of (Mean±S.D) values for genotoxic indices with symbol ** denotes significant deviation ($p < 0.05$). Type I-IV illustrates low damage (type I); moderate damage (type II); high damage (type III); complete damage (type IV).

erythrocyte hemolysis may be the cause of elevated MCH as well as MCV values. The study on common carp conducted by [28] similarly reported a remarkable decrease in MCHC. The same results in *C. batrachus* after exposure to arsenic were also reported by [29].

The white blood cells (WBCs) play a critical role in the cellular and immune systems. Our findings indicated a reduction in WBCs and lymphocyte counts compared to the untreated group. Similarly, [25] observed a significant decrease in WBCs and lymphocyte counts in fish exposed to lead, chromium, and copper, compared to the control. Ergönül M. et al. [30] reported similar results, mirroring our findings. The decline in WBC values may signify an impaired immune system in the fish, as suggested by [31]. In the current investigation, when *C. idella* was subjected to particular heavy metals, the number of platelets was reduced in the treatment groups relative to the untreated group. The decline in the number of platelets suggested a possible hemodilution mechanism that was probably

brought on by gill damage or impaired osmoregulation. Our results confirm those of [32], who subjected *C. punctatus* to heavy metal exposure.

Blood serves as a comprehensive indicator of the overall body's health. Changes in different hematological measures signify the response to pollution and can function as a method for the early identification of particular issues related to aquatic pollution [33]. Research on various fish species indicates that heavy metals can impact the biochemical traits present in tissues and blood. Alterations in tissue or blood composition may induce changes, leading to an elevation or reduction in the observed values. Blood glucose emerges as a reliable and sensitive marker for identifying environmental stress caused by pollutants in fish. In the conducted experiment with *C. idella*, there were significant increases in blood glucose levels. Our study demonstrates similarities to the investigation conducted by [34], in which the Caspian roach was subjected to lead exposure. Islam S.M. et al. [35] documented the occurrence

of hyperglycemia in *Pangasianodon hypophthalmus* exposed to chromium. Cholesterol serves as a crucial structural element in membranes and acts as a precursor to all steroid hormones. The concentration of cholesterol is considered an indicator of environmental stress caused by heavy metals [36]. Our results demonstrated a rise in cholesterol levels across all experimental groups compared to the control. The heightened cholesterol levels induced by heavy metals could potentially be attributed to liver failure, subsequently leading to an elevation in serum concentration. Comparable outcomes were achieved in the case of *C. batrachus* subjected to arsenic exposure, as documented by [29]. Additionally, Tabat J.L. et al. [37] observed increased levels of glucose and cholesterol in catfish fingerlings when exposed to cadmium.

Triglycerides primarily function as a cellular energy source and can be employed to evaluate nutritional status. A recent study identified a decrease in triglyceride levels, indicating liver damage induced by oxidative stress. Heydarnejad M.S. et al. [38] documented comparable findings, indicating a reduction in triglyceride levels in rainbow trout treated with cadmium. Ali Z. et al. [39] demonstrated a decline in triglyceride levels in *C. carpio* following exposure to manganese and chromium, exhibiting similarity with our research findings. The levels of tissue proteins serve as an indicator of stress induced by xenobiotics in aquatic organisms. In all groups treated with metals, there was an observed reduction in the entire protein content of the experimental fish's blood serum. Conditions, including liver cirrhosis, renal nephrosis, or alterations in enzymatic activity necessary for protein synthesis, could be the cause of the drop in plasma protein levels. Similar results in *C. batrachus* exposed to arsenic trioxide were reported by [40]. The results of our research align with those of [37], who observed similar outcomes. In this study, after being exposed to various heavy metals, the experimental fish *C. idella* showed higher blood urea, serum creatinine, and uric acid levels than the untreated group. Fluctuations in levels of creatinine are a sign of renal failure. These results are in alignment with the findings reported by [41], who found that *C. punctatus* exposed to mercuric chloride had higher levels of creatinine. According to research by [42], *C. gariepinus* subjected to cadmium showed elevated uric acid levels. Ali Z. et al. [39] identified a rise in uric acid and urea after manganese and chromium treatment in *C. carpio*. Our results are consistent with those of [41], who examined the effects of mercuric chloride on *C. punctatus*.

The presence of electrolytes in the bloodstream serves as a bioindicator of the health and physiological condition of fish. Exposure of grass carp to cadmium, lead, and mercury resulted in observable changes in electrolyte levels. There was a decrease in calcium and phosphorus levels, accompanied by an increase in sodium, potassium, and iron levels. The exposure of *C. idella* to selected heavy metals induced hypophosphatemia and hypocalcemia. This result is parallel to the study conducted by [43], who also found hypophosphatemia and hypocalcemia in *H. fossilis* subjected to lead. Increased sodium and potassium

activity were shown in grass carp following exposure to zinc and mercury, as reported by [44]. It was observed that the groups that received heavy metal treatment had higher iron concentrations. The study of [45], which found elevated iron levels in common carp subjected to harmful heavy metals, lends credence to our findings.

Comet test was used to identify metal-induced substantial changes in DNA damage levels. Several genetic pollutants can be efficiently screened using this method in the peripheral blood of various fish species [46]. DNA damage significantly increased in *C. idella* peripheral blood erythrocytes during the current experiment as metal concentrations increased. In terms of the percentage of erythrocytic damaged comet cells, the different metals' abilities to damage DNA followed the following sequence Pb>Hg>Cd. Similar results of DNA damage were reported in *C. idella* induced by metals such as Pb, Cr, and Cu [47]. Kousar S. et al. [48] reported the comparable genotoxic effects of different metals, like Cu, As, and Zn in peripheral erythrocytes of four freshwater fish species viz. *L. rohita*, *C. mrigala*, *C. catla*, and *C. idella*. Due to their nucleated form, erythrocytes were chosen as the ideal study material for fish DNA damage, even though the damage may also be found in the tissues of the fish's gills and kidneys [49]. Moreover, metal-induced DNA damage raised severe concerns about the possibility that metals could endanger grass carp's ability to survive in their native aquatic environments.

Conclusions

In conclusion, anthropogenic activities and manufacturing firms are the leading contributors to heavy metal toxicity in aquatic ecosystems. Heavy metals can easily infiltrate aquatic environments and eventually reach the bodies of aquatic species. Fish are a vital source of animal-based protein and come into close contact with heavy metals in the aquatic environment as a result of pollution. The results of our study indicate that grass carp are sensitive to the harmful effects of mercury, lead, and cadmium on their hematological, blood-biochemical, and genotoxic markers. Within the realm of environmental biomonitoring, the factors examined in this research can function as anticipatory markers for evaluating the influence of heavy metals on both fish and other aquatic organisms.

Acknowledgments

We would like to express our deepest gratitude to all those, who contributed to the completion of current research.

Authors contribution

Sabahat Saeed: Conceptualization, Investigation; Formal Analysis, Writing -original draft. Noman Waheed: Data Curation, Investigation, Writing -original draft,

Methodology. Visualization. Muhammad Fiaz Khan: Supervision. Adeeba Naseer: Review and editing. Shumaila Noreen: Review final draft. Usama Nawaz: Review final draft. Zeeshan Ahmad: Review final draft. Shehzad Ghayyur: Co-Supervision. Xu Rifu: Approved final Manuscript.

Funding

This work was supported by grants from the National Natural Science Foundation of China (Grant No. 32172714 and 31902145); the Project of Science and Technology Development Plan of Jilin Province (Grant No. 20220101306JC and 20210202100NC); the Project of Education Development Plan of Jilin Province (Grant No. JJKH20220362KJ); and the China Agriculture Research System of MOF and MARA (Grant No. CARS-41-Z16).

Conflict of Interest

The authors declare no conflict of interest.

References

- ALI H., KHAN E., ILAHI I. Environmental chemistry and ecotoxicology of hazardous heavy metals: environmental persistence, toxicity, and bioaccumulation. *Journal of Chemistry*. **2019** (4), 1, **2019**.
- SAJID M., NAZAL M.K., BAIG N., OSMAN A.M. Removal of heavy metals and organic pollutants from water using dendritic polymers based adsorbents: a critical review. *Separation and Purification Technology*. **191**, 400, **2018**.
- BARLOW N.L., BRADBERRY S.M. Investigation and monitoring of heavy metal poisoning. *Journal of Clinical Pathology*. **76** (2), 82, **2023**.
- SABULLAH M., AHMAD S., SHUKOR M., GANSAU A., SYED M., SULAIMAN M., SHAMAAN N. Heavy metal biomarker: Fish behavior, cellular alteration, enzymatic reaction and proteomics approaches. *International Food Research Journal*. **22** (2), 435, **2015**.
- YADAV M., GUPTA R., SHARMA R.K. Green and sustainable pathways for wastewater purification. In book: *Advances in Water Purification Techniques* (pp.355-383). Ahuja Consulting, Calabash, NC, United States, **2019**.
- AUTHMAN M.M., ZAKI M.S., KHALLAF E.A., ABBAS H.H. Use of fish as bio-indicator of the effects of heavy metals pollution. *Journal of Aquaculture Research and Development*. **6** (4), 328, **2015**.
- BALALI-MOOD M., NASERI K., TAHERGORABI Z., KHAZDAIR M.R., SADEGHI M. Toxic mechanisms of five heavy metals: mercury, lead, chromium, cadmium, and arsenic. *Frontiers in Pharmacology*. **12**, 643972, **2021**.
- MANUHARA Y.S.W., WIBOWO A.T., WINARNI D., ISLAMATASYA U., KHASANAH U.W.N. The comparison toxicity effects of lead and cadmium exposure on hematological parameters and organs of mice. *Ecology, Environment and Conservation*. **26** (4), 1842, **2020**.
- RICE K.M., WALKER JR E.M., WU M., GILLETTE C., BLOUGH E. Environmental mercury and its toxic effects. *Journal of Preventive Medicine and Public Health*. **47** (2), 74, **2014**.
- YANG L., ZHANG Y., WANG F., LUO Z., GUO S., STRÄHLE U. Toxicity of mercury: Molecular evidence. *Chemosphere*. **245**, 125586, **2020**.
- ARDESHIR R.A., MOVAHEDINIA A.-A., RASTGAR S. Fish liver biomarkers for heavy metal pollution: a review article. *American Journal of Toxicology*. **2** (1), 1, **2017**.
- ADESHINA I., EMIKPE B.O., JENYO-ONI A., AJANI E.K., ABUBAKAR M.I. Haematology, plasma biochemistry and serum of table size African catfish, *Clarias gariepinus*, naturally infected with *Listeria* species in Oyo State. *Comparative Clinical Pathology*. **29** (3), 69, **2020**.
- NWANI C.D., EKWUEME H.I., EJERE V.C., ONYEKE C.C., CHUKWUKA C.O., PEACE O.S., NWADINIGWE A.O. Physiological effects of paraquat in juvenile African catfish *Clarias gariepinus* (Burchfiel 1822). *Journal of Coastal Life Medicine*. **3** (1), 35, **2015**.
- MATOS L.A., CUNHA A.C., SOUSA A.A., MARANHÃO J.P., SANTOS N.R., DE MC GONÇALVES M., DANTAS S.M.D.M., E SOUSA J.M.D.C., PERON A.P., DA SILVA F.C.C. The influence of heavy metals on toxicogenetic damage in a Brazilian tropical river. *Chemosphere*. **185**, 852, **2017**.
- LUO L., WANG B., JIANG J., FITZGERALD M., HUANG Q., YU Z., LI H., ZHANG J., WEI J., YANG C. Heavy metal contaminations in herbal medicines: Determination, comprehensive risk assessments, and solutions. *Frontiers in Pharmacology*. **11**, 595335, **2021**.
- GORINI F., MURATORI F., MORALES M.A. The role of heavy metal pollution in neurobehavioral disorders: a focus on autism. *Review Journal of Autism and Developmental Disorders*. **1** (4), 354, **2014**.
- SHAHJAHAN M., TASLIMA K., RAHMAN M.S., AL-EMRAN M., ALAM S.I., FAGGIO C. Effects of heavy metals on fish physiology – a review. *Chemosphere*. **300**, 134519, **2022**.
- VAJARGAH M.F., NAMIN J.I., MOHSENPOUR R., YALSUYI A.M., PROKIĆ M.D., FAGGIO C.. Histological effects of sublethal concentrations of insecticide Lindane on intestinal tissue of grass carp (*Ctenopharyngodon idella*). *Veterinary Research Communications*. **45** (4), 373, **2021**.
- HANEY D., HURSH D., MIX M., WINTON J. Physiological and hematological changes in chum salmon artificially infected with erythrocytic necrosis virus. *Journal of Aquatic Animal Health*. **4** (1), 48, **1992**.
- IBRAHIM A.T.A. Toxicological impact of green synthesized silver nanoparticles and protective role of different selenium type on *Oreochromis niloticus*: hematological and biochemical response. *Journal of Trace Elements in Medicine and Biology*. **61**, 126507, **2020**.
- SINGH N.P., MCCOY M.T., TICE R.R., SCHNEIDER E.L. A simple technique for quantitation of low levels of DNA damage in individual cells. *Experimental Cell Research*. **175** (1), 184, **1988**.
- HEDAYATI A., GHAFARI Z. Effect of Mercuric Chloride on Some Hematological, Biochemical Parameters in Silver Carp (*Hypophthalmichthys Molitrix*). *International Journal of Veterinary Medicine Research & Reports*. **2013**.
- NAZ S., HUSSAIN R., ULLAH Q., CHATHA A.M.M., SHAHEEN A., KHAN R.U. Toxic effect of some heavy metals on hematology and histopathology of major

- carp (*Catla catla*). Environmental Science and Pollution Research. **28** (6), 6533, **2021**.
24. ZULFAHMI I., RAHMI A., MULIARI M., AKMAL Y., PAUJIAH E., SUMON K. A., RAHMAN M. Exposure to Lead Nitrate Alters Growth and Haematological Parameters of Milkfish (*Chanos chanos*). Bulletin of Environmental Contamination and Toxicology. **107** (5), 860, **2021**.
 25. SHAH N., KHISROON M., SHAH S.S.A. Assessment of copper, chromium, and lead toxicity in fish (*Ctenopharyngodon idella Valenciennes, 1844*) through hematological biomarkers. Environmental Science and Pollution Research. **27** (26), 33259, **2020**.
 26. CHAUDHARY A., GUL JAVAID K., BUGHIO E., FAISAL N. Study on histo-biochemical biomarkers of chromium induced toxicity in *Labeo rohita*. Emerging Contaminants. **9** (1), 100204, **2023**.
 27. RAMESH P., RAMACHANDRA MOHAN M. Effects of Cadmium Chloride on Hematological Profiles in Freshwater Fish *Channa punctatus* (Bloch). Journal of Environment and Ecology. **39** (4A), 1096, **2021**.
 28. KHALESİ M., ABEDI Z., BEHROUZI S., ESKANDARI S.K. Haematological, blood biochemical and histopathological effects of sublethal cadmium and lead concentrations in common carp. Bulgarian Journal of Veterinary Medicine, **20** (2), **2017**.
 29. KUMAR R., BANERJEE T. Arsenic induced hematological and biochemical responses in nutritionally important catfish *Clarias batrachus* (L.). Toxicology Reports. **8** (3), 148, **2016**.
 30. ERGÖNÜL M., ALTINDAĞ A., ATASAĞUN S., SHAH S., KARACAKAYA P. Alterations in hematological, immunological and biochemical parameters of tench, *Tinca tinca* after acute zinc exposure. The Journal of Animal and Plant Sciences. **28** (5), 1357, **2018**.
 31. ABDEL-WARITHA.-W.A., YOUNISE.-S.M., AL-ASGAH N.A., RADY A.M., ALLAM H.Y. Bioaccumulation of lead nitrate in tissues and its effects on hematological and biochemical parameters of *Clarias gariepinus*. Saudi Journal of Biological Sciences. **27** (3), 840, **2020**.
 32. RAMESH P., RAMACHANDRA MOHAN M. Effects of Cadmium Chloride on Hematological Profiles in Freshwater Fish *Channa punctatus* (Bloch). Journal of Environment and Ecology. **39** (4A), 1096, **2021**.
 33. MIAH M. Deleterious effects of mercuric chloride on blood biochemistry, liver and kidney histology in female albino mice. Journal of Agriculture, Food and Environment. **2** (2), 18, **2021**.
 34. HEDAYATI A., DARABITABAR F. Lethal and sublethal impacts of lead on some hematological, biochemical and immunological indices in Caspian roach (*Rutilus rutilus*). Pollution. **3** (1), 21, **2017**.
 35. ISLAM S.M., ROHANI M.F., ZABED S.A., ISLAM M.T., JANNAT R., AKTER Y., SHAHJAHAN M. Acute effects of chromium on hemato-biochemical parameters and morphology of erythrocytes in striped catfish *Pangasianodon hypophthalmus*. Toxicology Reports. **7**, 664, **2020**.
 36. ÖZÇELİK S., CANLI M. Combined effects of metals (Cr^{6+} , Hg^{2+} , Ni^{2+} , Zn^{2+}) and calcium on the serum biochemistry and food quality of the Nile fish (*Oreochromis niloticus*). Journal of Food Composition and Analysis. **115** (10), 104968, **2023**.
 37. TABAT J.L., JEHU A., KOGI E., HABILA J.D. Sublethal toxicity effects of cadmium (Cd^{2+}) on serum biochemistry in fingerlings and juveniles of fresh water catfish, *Clarias gariepinus* (Burchell, 1822). International Journal of Fisheries and Aquatic Studies. **9** (4), 1, **2021**.
 38. HEYDARNEJAD M.S., KHOSRAVIAN-HEMAMAI M., NEMATOLLAHI A. Effects of cadmium at sub-lethal concentration on growth and biochemical parameters in rainbow trout (*Oncorhynchus mykiss*). Irish Veterinary Journal. **66** (1), 1, **2013**.
 39. ALI Z., YOUSAFZAI A.M., SHER N., MUHAMMAD I., NAYAB G.E., AQEEL S.A.M., SHAH S.T., ASCHNER M., KHAN I., KHAN H. Toxicity and bioaccumulation of manganese and chromium in different organs of common carp (*Cyprinus carpio*) fish. Toxicology Reports. **8**, 343, **2021**.
 40. PICHHODE M., GAHERWAL S. Effect of heavy metal toxicity, arsenic trioxide on the biochemical parameter of fresh water fish, *Clarias batrachus*. Pollution Research. **39**, 123, **2020**.
 41. GILL H.K., PAUL K., DUA A. Evaluation of Genotoxic And Biochemical Stress Response To Mercuric Chloride In Erythrocytes of Freshwater Fish *Channa Punctatus*. Research Square. **2021**.
 42. ELARABANY N., BAHNSAWY M., ALFAKHERI M. Effects of cadmium on some haematological and biochemical parameters of African catfish, *Clarias graiepinus*. African Journal of Biological Sciences. **15** (1), 43, **2019**.
 43. SRIVASTAV A.K., RAI R., SUZUKI N., MISHRA D., SRIVASTAV S.K. Effects of lead on the plasma electrolytes of a freshwater fish, *Heteropneustes fossilis*. International Aquatic Research. **5** (1), 1, **2013**.
 44. SALAH EL-DEEN M., SALEH R., EL-RAZIK A., ABO-HEGAB E. Recorded Changes in some Biochemical indicators of grasscarp *ctenopharyngodon idella* exposed to mercury and zinc. Egyptian Journal of Aquatic Biology and Fisheries. **5** (2), 57, **2001**.
 45. VINODHINI R., NARAYANAN M. The impact of toxic heavy metals on the hematological parameters in common carp (*Cyprinus carpio* L.). Iranian Journal of Environmental Health Science & Engineering. **6** (1), **2009**.
 46. MATSUMOTO S.T., MANTOVANI M.S., MALAGUTTHI M.I.A., DIAS A.L., FONSECA I.C., MARIN-MORALES M.A. Genotoxicity and mutagenicity of water contaminated with tannery effluents, as evaluated by the micronucleus test and comet assay using the fish *Oreochromis niloticus* and chromosome aberrations in onion root-tips. Genetics and Molecular Biology. **29**, 148, **2006**.
 47. SHAH N., KHAN A., HABIB KHAN N., KHISROON M. Genotoxic consequences in common grass carp (*Ctenopharyngodon idella Valenciennes, 1844*) exposed to selected toxic metals. Biological Trace Element Research. **199**, 305, **2021**.
 48. KOUSAR S., JAVED M. Diagnosis of metals induced DNA damage in fish using comet assay. Pakistan Veterinary Journal. **35** (2), 168, **2015**.
 49. ARSLAN Ö.Ç., BOYACIOĞLU M., PARLAK H., KATALAY S., KARAASLAN M.A. Assessment of micronuclei induction in peripheral blood and gill cells of some fish species from Aliğa Bay Turkey. Marine Pollution Bulletin. **94** (1-2), 48, **2015**.