

Original Research

Leaf Trait-Based Profiling to Select High-Performing Woody Plant Species for Land Restoration

Ridesti Rindyastuti^{1*}, Abban Putri Fiqa², Dewi Ayu Lestari¹, Janis Damaiyani¹, Apriyono Rahadianoro¹, Titut Yulistyarini², Decky Indrawan Junaedi²

¹Research Center for Applied Botany-BRIN, KST Dr. (H.C.) Ir. Soekarno, Jl. Raya Jakarta-Bogor Km. 46, Cibinong, Bogor, West Java 1691, Indonesia

²Research Center for Ecology and Ethnobiology-BRIN, KST Dr. (H.C.) Ir. Soekarno, Jl. Raya Jakarta-Bogor Km. 46, Cibinong, Bogor, West Java 1691, Indonesia

Received: 15 March 2024

Accepted: 5 August 2024

Abstract

Bringing ecosystem services back to degraded areas through land restoration is a complex task across countries. For the establishment of vegetation, selecting high-performing plant species using ecological trait-based profiling was conducted to initiate restoration projects. Six-leaf ecological traits of twelve plant species were investigated, including Specific Leaf Area (SLA), Leaf Dry Matter Content (LDMC), Leaf Nitrogen Concentration (LNC), chlorophyll content, stomatal density, and stomata aperture size. The correlation between SLA and LNC, and also SLA and LDMC, indicated plant trade-offs in growth and life history strategy related to herbivory and flammability, respectively. SLA and LDMC are considered two strong predictors of plant performance which serve as the main requirement for selecting plant species in restoration, especially in growth performance (SLA), flammability, and herbivore resistance (LDMC). Moreover, according to the tropical Malesian region, leaf traits of exotic species *Swietenia macrophylla* and *Ficus religiosa* indicated a risk of invasiveness to some extent. Among ten native species, *Protium javanicum*, *Syzygium polyanthum*, *Canarium vulgare*, and *Artocarpus heterophyllus* are considered the pioneer species in restoration. The pioneers with intensive weed control are *Durio zibethinus* and *Cinnamomum sintoc*. Whilst, the other native species *Cynometra schefferi*, *Dimocarpus longan*, *Garcinia dulcis*, *Diospyros celebica* are considered inserted species. We demonstrated that plant profiling using eco-physiological traits takes an important role in plant species selection, which serves as a critical stage in successful restoration programs.

Keywords: Specific leaf area (SLA), land restoration, functional traits, leaf dry matter content (LDMC), plant life strategy

*e-mail: ride17@gmail.com

Tel.: +62-811-1933-3605

Introduction

Numerous land-use changes and utilization, such as land conversion, agriculture [1] urban settlements, and industries, lead to land degradation and biodiversity loss, either at the species to ecosystem levels [2–4]. Ecological restoration is a project to accelerate ecosystem recovery of degraded areas by restoring community structure, species composition, and ecosystem functions [5, 6]. In general, the establishment of plant cover is the primary objective of the programs [7]. The establishment of plant cover is important, making the selection of plant species a crucial stage in initiating a restoration project.

The selection of plant species for restoration should consider ecosystem sustainability and current environmental changes. Utilizing native plant species, especially endemic and endangered ones, is strongly recommended as they provide historical references for the habitat. A multi-criteria approach can be employed in selecting plant species for restoration, taking into account both ecological and socio-economic aspects [8]. In recent decades, there has been an urgent need for management authorities to conduct plant trait profiling and assessment. This is essential for selecting plant species with high ecological establishment capacity, performance, and competitive ability in ecosystems during restoration or rehabilitation programs [9–12]. The selected species should also demonstrate adaptability to water and temperature stresses [13], resistance to herbivory and flame disturbance, and the ability to outperform other species [14]. Moreover, they should possess the capability to resist invasive species [15, 16].

Plant selection for constructing the recommendation system for effective land restoration of degraded areas could be established by considering plant traits that demonstrate the advantages of plants to ecosystems [14, 17–19]. These traits encompass a range of characteristics, including morphological, eco-physiological, biochemical, socioeconomic, and reproductive factors that may be associated with multiple biological pathways and their services to ecosystems [8, 19]. Several leaf traits were suggested as proxies for the ecological function of carbon assimilation, water relations, and energy balance such as SLA (Specific Leaf Area), LDMC (Leaf Dry Matter Content), leaf thickness, and leaf nutrient [20]. SLA represents area-based photosynthetic activities. Other leaf traits, such as stomatal characteristics, nitrogen levels, and chlorophyll content, are also essential for assessing the eco-physiological profiles of plants [21]. Several leaf traits are also proxies of nutrient uptake and photosynthesis, which could be indicated by leaf nitrogen and chlorophyll content [22]. Plant traits showed a complex interaction effect rather than a single impact; therefore, species may possess a combination of response-and-effect traits that matter to the ecological process [23].

In most developing countries, the plant selection processes for restoration were rarely implemented, because the research which serves as a starting point for the selection

process is still limited [24, 25]. Previously, [26] proposed a list of local species for restoration based on the carbon sequestration capacity and society's preferences for plant species for restoration using questionnaire methods. [27] profiled ten woody species that have ecological adaptation to dry environments based on leaf and photosynthetic characteristics. To support natural biodiversity and ecosystem sustainability, proposing high-performed local plant species is preferable, including the native and introduced plants in natural forest areas in a targeted restored environment [28].

Plant trait-based profiling and the development of a framework for trait comparison are still rare to be conducted either for researchers or the management of restoration programs in Indonesia. Here, we applied a low-cost and affordable method to select and recommend highly performed and adaptive woody plant species. We focused on eco-physiological leaf traits that will be used to predict plant survival rates during restoration stages. The study on the variation and correlation of leaf traits across plant species contributes to ecological perspectives on plant life history strategies in tropical Malesian regions. Plant profiling and role prediction of plant species that have high growth and advantageous eco-physiological strategies will be highly implemented for successful land restoration.

Materials and Methods

We investigated leaf traits of twelve local woody plant species using 2-year-old seedlings under a controlled condition. These traits are SLA, LDMC, total chlorophyll content, LNC, stomatal density, and aperture size. We also examined the correlation between these measured traits.

Plant Materials

A total of twelve species, including ten native and two exotic plant species to the Indonesian region (*Ficus religiosa* and *Swietenia macrophylla*), were used in this research (Table 1). All studied local species (10 natives and 1 exotic species) are broad-leaved-woody evergreen species, while one exotic species (*S. macrophylla*) is a broad-leaved woody semi-deciduous species. These twelve plant species were chosen based on the suggested list by [26, 29], which selected plant species based on society and stakeholders' preferences for restoration and rehabilitation programs. Two exotic species were chosen to compare leaf traits between native and exotic plant species. Plants were propagated and collected in Purwodadi Botanic Garden-Indonesian Research and Innovation Agency (BRIN), East Java, Indonesia, both in adult and seedling stages. Three plants were used for each species for replicability. The information on plant species, habitats [30], and distribution ranges either native or introduction ranges [31] are presented in Table 1.

Table 1. Species names, local names, habitats, and distribution ranges of studied plants.

No	Plant species	Habitat	Distribution
1	<i>Canarium vulgare</i> Leenh.	Primary forests in calcareous areas and lowland rainforest	Jawa, Lesser Sunda Is., Maluku, New Guinea, Solomon Is., Sulawesi
2	<i>Syzygium polyanthum</i> (Wight) Walp.	Lowland primary and secondary forest	Indochina and Malesia
3	<i>Cynometra schefferi</i> (K.Schum.) Rados.	Primary lowland forest	Bismarck Archipelago, Maluku, New Guinea, Sulawesi
4	<i>Artocarpus heterophyllus</i> Lam.	Tropical mix forest	India, introduced to Africa and Southeast Asia, Vietnam, Thailand to Indonesia
5	<i>Ficus religiosa</i> L.	Mountain forest	Africa, Asia, North America, and Oceania, naturalized in Southeast Asia
6	<i>Garcinia dulcis</i> (Roxb.) Kurz	Lowland tropical rainforest 0–500 m asl	Andaman Island, Indonesia, North Queensland
7	<i>Protium javanicum</i> Burm.f.	Tropical rain forest 35–250 m asl	Java, Lesser Sunda Island (Indonesia)
8	<i>Swietenia macrophylla</i> (L.) Jacq.	Tropical forest	Mexico to Bolivia and Brazil, introduced to Seychelles (Africa), Indochina and Solomon Is.
9	<i>Cinnamomum sintoc</i> Blume	Hill forest, lowland, and montane forests up to 2400 m altitude	Thailand, Peninsular Malaysia, Sumatra, Java, the Lesser Sunda Islands, and Borneo
10	<i>Dimocarpus longan</i> Lour.	Tropical rain forest	Southern China, Taiwan, Thailand, Malaysia, Indonesia, Kamboja, Laos, Vietnam, India, Filipina, Australia.
11	<i>Diospyros celebica</i> Bakh.	Tropical lowland forest	Sulawesi Island (Indonesia)
12	<i>Durio zibethinus</i> L.	Mixed Dipterocarp Forest along riverside and hilly areas >800 m asl.	Southeast Asia

Leaf Traits Measurements

Specific Leaf Area (SLA) and Leaf Dry Matter Content (LDMC)

Measurements of Specific Leaf Area (SLA) and Leaf Dry Matter Content (LDMC) were carried out on 2-year-old seedlings using destructive methods, as suggested by [32]. Three plants were used as a replication for each species. Five healthy leaves of each plant were selected and harvested. Leaf area was measured using ImageJ [33]. LDMC was obtained from the dry-oven weights of leaves at 80°C until constant dry weight. Leaf samples were weighed using an analytical balance in each plant species with three individuals as replication. SLA value is the ratio of leaf area value and leaf dry mass (cm^2/g).

Stomatal Density (SD) and Stomata Aperture Size (SA)

Stomata observations were carried out using Scanning Electron Microscope (SEM) procedures [34]. Three plants were used for each species for replicability. In the SEM procedure, the leaves were cut into 5x5 mm sizes and excluded the main and secondary leaf veins. The samples were observed using SEM (FEI type Inspects 25) in the magnification of 500–5000 times. The number of stomata was counted from the magnified

epidermal tissue areas (square millimeters). The stomatal density was calculated as density per square millimeter. The stomata aperture size was determined by measuring the length of the opened area of the stomata.

Total Chlorophyll Content (CCM)

Total chlorophyll content was measured using CCM-200 plus Chlorophyll Content Meter on mature leaves. Three plants for each species were used as replication and 5 leaves of each plant were sampled. Three sites of leaf surface, i.e. apex-near, middle, and base-near leaf sites, were selected as measurement points to obtain representative chlorophyll content data.

Leaf Nitrogen Concentration (LNC)

Leaf nitrogen concentration was measured at the Soil Laboratory, East Java Agricultural Technology Study Center (BPPT), Malang, using the Kjeldahl method [35]. Three plants for each species were used as replications. Two hundred grams of leaf for each plant were sampled and dry-oven baked. Leaf Nitrogen Concentration (LNC) expressed in mg/g was obtained from the calculation of Nitrogen concentration (%) and the leaf dry mass. Whilst, leaf dry mass was calculated from leaf area (mm^2) multiplied by Specific Leaf Weight (SLW) and measured as mg/cm^2 .

Data Analysis

First, we conducted a normality test to assess whether the data were normally distributed or not. Then, a one-way analysis of variance (ANOVA) and Tukey's test were conducted to compare leaf traits among plant species at a confidence level of 95% ($\alpha \leq 0.05$). Pearson's correlation tests were performed to identify the correlation between two leaf traits at a confidence level of 95% ($P \leq 0.05$). Normality test, ANOVA, Tukey's test, and Pearson correlation test were performed in R 4.1.0 [36]. Principal Component Analysis (PCA) was conducted to eliminate weak leaf trait predictors on high plant performances for land restoration. The PC analysis was performed using the statistical software PAST 4.0 [37]. High-performance characteristics, which are shown by high SLA, low LDMC, low LNC, and low SD, are used as criteria for selecting plant species utilized in the restoration programs.

Results and Discussion

Variability of Leaf traits Among Species

Eco-physiological traits showed variability across plant species [38]. There were significant differences in all six-leaf traits across the twelve selected plant species based on the ANOVA test (Fig. 1). The significant differences in all six-leaf traits among twelve plant species in this study indicate how the growth strategies of plants vary across species in tropical habitats. Although there is evidence of infra-specific variation in several plant traits which indicated plasticity and the influence of environmental gradients [39], these plant traits are useful to distinguish plant types based on growth and assimilate allocation strategies at the specific level.

In terrestrial ecosystems, SLA is thought to be a reliable proxy for indicating a plant's photosynthesis, growth rate, and performance in its environment. The higher the SLA value, the higher the photosynthetic activities will be. It ranges from 6.636 to 2040.816 cm²/g of all life forms [32, 40, 41]. *Ficus religiosa* had the highest SLA value in this study relative to other species. SLA of *F. religiosa* widely varies between samples, which indicates infra-specific plasticity [39]. This could be possible because this species has a wide distribution range. On the other hand, an endemic species *Diospyros celebica* had the lowest SLA value, which was not significantly different from the other three native species, i.e., *Durio zibethinus*, *Garcinia dulcis*, and *Arthocarpus heterophyllus*.

The recorded SLA of tropical plants ranges from 90.5 to 298.5 cm²/g [25], and 29.38 to 88.55 mm²/mg for endemic species *Zanthoxylum acanthopodium* [24]. Therefore, the SLA of *F. religiosa* in this study was considered as high compared to those of other species and the global SLA record of plant species. The high SLA indicated a strong plant competition in resource uptake, particularly at the maximum use of sunlight. On the other hand, species with the lowest SLA (*D. zibethinus*, *G. dulcis*,

and *A. heterophyllus*) did not always necessarily indicate poor performance. It might show high survival rates on degraded land with drought and less fertile soil conditions [42]. Moreover, the exotic species *Swietenia macrophylla* showed a relatively high value of SLA. This indicated that the range of SLA value among the native species was relatively narrow, while those of the exotic species were relatively high.

LNC of the studied species ranged from 0.004 to 0.135 mg/g. The LNC of *Protium javanicum* was 0.004±0.00 mg/g, which was significantly lower than that of other species. On the other hand, *A. heterophyllus* had the highest LNC value with 0.135±0.001 mg/g. Different growth strategies between native and exotic species are also indicated by SLA and LNC. Exotic species *S. macrophylla* had relatively high SLA but low LNC value. Whilst, the native species *D. celebica* and *D. zibethinus* had the lowest SLA value, yet relatively high LNC. This SLA comparison between native and exotic species indicates that exotic species commonly outperform natives [43]. However, there are possibilities that native species may perform as high as exotic and even invasive species.

In general, the LNC value is highly associated with the rate of photosynthesis, which indirectly influences plant development [21, 44]. In this research, the species with the highest LNC was *A. heterophyllus*. A species with high LNC and SLA values will have a short life span and will accelerate the nutrition cycle in the ecosystem [45, 46]. However, it should be emphasized that the LNC value strongly depends on environmental conditions, such as soil, humidity, and temperature [21, 47]. Moreover, the use of *A. heterophyllus* will contribute to fast soil nutrient recovery in a restoration.

In contrast to the SLA, the LDMC of *F. religiosa* was the lowest compared to other observed species. Species with high LDMC values will be more resistant to herbivores or wind disturbances [21]. LDMC is a plant trait that describes the nature of a species' resource-conserving strategy [48, 49]. As a result, *Cynometra schefferi*, which had the highest LDMC value of all the species studied, was likely to exhibit wind and herbivore disturbances resistance.

The variability of stomatal traits demonstrates the different growth strategies on leaf gas exchange and water loss regulation between native and exotic species (Fig. 2). The exotic species *S. macrophylla* showed the lowest value of stomata aperture size but with relatively high stomatal density. In contrast, the native species *D. celebica* had the lowest stomatal density and the largest stomata aperture size. The high stomatal density of exotic species *S. macrophylla* represented high photosynthetic activity, fast growth rate, and water loss of plants and soil [27, 50]. Planting this species in restoration may negatively affect the surrounding restored ecosystem and water conservation.

Correlations, Trade-Offs between Leaf Traits and Life History Strategies

We investigated the correlation of measured leaf traits among plant species to identify the bivariate relationship




Fig. 1. Box plot graph of leaf trait values of studied species.

Species 1–12 were listed according to Table 1. Different letters represent significant differences between species according to a post hoc Tukey test ($p < 0.05$).




Fig. 2. Scanning electron micrographs showing the density of stomata on the abaxial surface of 12 species studied, which are listed according to Table 1.

between leaf traits. We extracted essential correlations between leaf traits at $P < 0.05$ and detected eight significant correlations (Fig. 3 and Table 2). A negative correlation between two particular functional leaf traits tends to demonstrate a trade-off in the life history strategy of plant species in adapting to a challenging environment [51]. In this study, negative correlations were detected between LNC with other important traits related to growth and photosynthesis, i.e., SLA and stomatal density. This functional plant group takes advantageous strategies to obtain rapid growth by increasing photosynthetic activity and foliage display, but tends to decrease nitrogen allocation into leaf organs. The high level of N may not be advantageous for plants since it could increase herbivory through the attractiveness of high nutrition [52, 53]. A negative correlation between SLA and LDMC may demonstrate a trade-off between leaf traits that relate to the level of flammability. According to [21, 54], flammability could be predicted through the dry matter content of plant leaves (LDMC). The studied plant group demonstrated the life strategies to maximize productivity and simultaneously implement flame and herbivory risk avoidance mechanisms.

Chlorophyll content had a positive correlation with nitrogen level and stomata aperture size as well as nitrogen level, which had a positive correlation with aperture size. Further, chlorophyll content was supported by stomata aperture size, but not by its density. A positive correlation between chlorophyll content and LNC was revealed by previous studies [55]. The chlorophyll content

is influenced by nitrogen level because nitrogen is one of the most essential nutrients for plant growth and serves as a major component of chlorophyll and protein [56, 57]. Nitrogen levels affect chlorophyll content that may influence the development of stomata to maximize gas exchange of photosynthetic activity. Since SLA was negatively correlated with nitrogen level, the increase of photosynthetic capacity might occur in the leaf area scale (cm^2). Consequently, when leaves had a large foliage display (high SLA value), leaf nitrogen levels tended to be decreased to develop an adaptation mechanism in protecting leaves from mechanical disturbance, which includes predatory herbivores or flame risks.

Strong Trait Predictor and Recommended Species for Restoration

Among the six investigated leaf traits in this study, strong predictors were preferred as the baseline in the determination of high performance-species in growth and life history strategy, which may be ecologically advantageous for restoration programs. Based on the component analyses, for these twelve studied species contexts, two leaf traits i.e. SLA and LDMC were stronger predictors compared to other leaf traits at a total of variance 70.58% which consist of 45.82% and 24.77% from PC1 and PC2 respectively (Fig. 4). We also arranged the information on ecological attributes that relate to light tolerance based on stomatal density and leaf morphology [58] to predict the role of plant species in restoration (Table 3).

Table 2. Bivariate combination of six plant traits.

Bivariate combination	r	P value	Correlation
Chlorophyll vs SLA	-0.418*	0.011	negative
Chlorophyll vs LDMC	-0.178	0.298	negative
Chlorophyll vs SD	-0.348	0.038	negative
Chlorophyll vs LNC	0.331*	0.049	positive
Chlorophyll vs SA	0.428**	0.009	positive
SLA vs LDMC	-0.484**	0.003	negative
SLA vs SD	0.064	0.710	positive
SLA vs LNC	-0.450**	0.006	negative
SLA vs SA	-0.174	0.310	negative
LDMC vs SD	-0.253	0.136	negative
LDMC vs LNC	-0.251	0.141	negative
LDMC vs SA	-0.184	0.284	negative
SD vs LNC	0.346*	0.039	positive
SD vs SA	-0.287	0.090	negative
LNC vs SA	0.469*	0.004	positive

*: Correlation is significant at the 0.05 level (2-tailed); **: Correlation is significant at the 0.001 level (2-tailed). SLA = Specific Leaf Area, LDMC = Leaf Dry Matter Content, C = Chlorophyll, SA = Stomata Aperture size, LNC = Leaf Nitrogen Concentration.




Fig. 3. Detected significant correlation between studied leaf traits according to the Pearson Correlation test ($p < 0.05$).

Specific Leaf Area (SLA) is widely proven as the best predictor of plant photosynthesis, growth rate, and performances in various habitats and ecosystems [59, 60], which play important roles in plant competition, or even in plant invasion context [21]. The large range of SLA among the species in the global plant and woody plant group itself leads to a wider opportunity to explore and identify native species with higher SLA. LDMC indicates a plant adaptation strategy that relates to the risk of flammability. The lower the LDMC value, the higher the plant species' ability to avoid destruction caused by a flame in their natural habitats [21, 61]. Wüest R.O. et al. [62] found that LDMC is a more reliable predictor relative to other plant traits, especially those related to environmental services. Therefore, the result of the study also supports the previous result about the importance of SLA and LDMC in plant functional ecology.

High growth rate and high resistance to flame and herbivory were important criteria in the restoration, indicated by high SLA and low LDMC values, respectively. The resistance to herbivores was also supported by low LNC. The determination of these criteria is consistent with most study results on plant traits carried out on seedlings under controlled conditions [53, 63]. The component of plant growth serves as species attributes rather than measured plant characteristics, which are more influenced by environmental conditions.

We recommend the use of native plant species that had high SLA but low LDMC, which demonstrated high photosynthetic activities and advantageous strategy in decreasing the risk of herbivory at restored habitats (Fig. 4; Table 3). Exotic species *Ficus religiosa* and *S. macrophylla* showed high SLA. Numerous study results showed that SLA strongly impacts plant growth rate associated with plant invasive occurrences. Species with

a high value of SLA are considered to show a high risk of invasiveness in recipient-restored tropical habitats [64, 65]. Therefore, these two species were less recommended to be planted in restored areas in tropical regions of Southeast Asia. Effective controls of invasive species are rarely in restoration projects because it is long-term and high-cost efforts. Providing basic information through research on intrinsic aspects of plant species could guide successful restoration projects [16].

On the other hand, a native species *P. javanicum* had a high SLA, higher than those of the exotic species *S. macrophylla*. Moreover, four species were relatively sun tolerant and had high SLA, therefore were highly recommended as strong-performing pioneers i.e. *P. javanicum*, *S. polyanthum*, *C. vulgare*, and *A. heterophyllum*. Regarding to the light tolerance and SLA value, *C. schefferi*, *G. dulcis*, *D. longan*, and *D. celebica* were recommended as inserted species in land restoration. *Cinnamomum sintoc* and *D. zibethinus* were two sun-tolerant species with relatively low SLA, therefore these two species were recommended as pioneer species on restored land with the intensive control of weeds.

Durio zibethinus with the local name “durian” is a native-tolerant species with a low value of SLA and high stomatal density. This species will be preferable for restoration because of its economic value by producing favored and delicious fruit. Because of the low SLA, as a pioneer plant, this species had low photosynthetic activities that led the plants to sufficiently compete with the weeds and invasive plants on the restored land. The high stomatal density of *D. zibethinus* indicated that this species might lose water rapidly due to high transpiration rates. Therefore, intensive weed control and regular watering are necessary treatments for this species in restored areas.




Fig. 4. Principal-component analysis of 6 leaf traits of 12 plant species. Different dots color represent different species.

Table 3. Ecological attributes and predicted role of the studied species on land restoration.

No	Species	Light tolerance	SD	SLA	LDMC	LNC	Role prediction on restoration
1	<i>C. vulgare</i>	sun	th	th	th	th	pioneer
2	<i>S. polyanthum</i>	sun	th	th	tl	tl	pioneer
3	<i>C. schefferi</i>	semi shade	tl	tl	h	tl	inserted
4	<i>A. heterophyllus</i>	sun	h	tl	l	h	pioneer
5	<i>F. religiosa</i>	semi shade	l	h	l	tl	risk as invasive
6	<i>G. dulcis</i>	shade	tl	l	th	tl	inserted
7	<i>P. javanicum</i>	semi sun	th	h	h	l	pioneer
8	<i>S. macrophylla</i>	sun	h	h	tl	tl	high risk as invasive
9	<i>C. sintoc</i>	sun	h	tl	tl	th	pioneer
10	<i>D. longan</i>	shade	l	tl	h	th	inserted
11	<i>D. celebica</i>	shade	l	l	h	th	inserted
12	<i>D. zibethinus</i>	sun	h	l	th	th	pioneer

Remarks (h = high, l = low, th = tend to high, tl = tend to low)

In field conditions, plant growth will be influenced by the nature of plants (shade or sunlight tolerance) and the characteristics of the habitat (shaded or opened environment). Shade-tolerant species could form small leaves to reduce the heat load, but in tropical forests, light-demanding species had large leaves compared to shade-tolerant plants to compete with their co-existing plants. Sun-tolerant species have a high SLA and Leaf Area Ratio (LAR) to increase photosynthetic rates by absorbing more sunlight in a light-limited environment, whereas shade-tolerant species have a low SLA compared to light-demanding species [66]. Plants that grow in shady environments could show a high survival rate by growing slowly since plants develop conservative strategies by sustaining in low-energy environments rather than maximizing net carbon gain in the shade [67].

In some restoration cases, native plant species showed high and moderate growth rates when seedlings were planted in degraded areas with open habitats, plenty of light, and adequate ground cover. Kestari D.A. et al. [68] revealed that native Bornean species such as *Shorea balangeran* and *S. lamellata* were growing well on open degraded land in coal-ex-mined areas. *Shorea balangeran* was also reported to have high survival and growth rates in the restoration of peat swamps in Borneo [69]. In the field situations, SLA, which represents growth rates, could also be positively correlated with plant survival rate likewise revealed by [13].

Conclusions

This study adjusts the manifest to ecological knowledge that eco-physiological traits of plants vary across species. The correlation between two leaf traits, especially SLA,

LNC, and LDMC, showed plant trade-offs in plant growth and life history strategy related to herbivory and the risk of flammability. The results strengthen the global perspective on SLA and LDMC as strong predictors in plant performance, which serve as the baseline of plant species selection for restoration. The two exotic species, i.e., *S. macrophylla* and *F. religiosa* showed a risk of invasiveness in tropical habitats of Southeast Asia based on sun adaptation and high SLA. Native plant species group in this study, i.e., *P. javanicum*, *S. polyanthum*, *C. vulgare*, and *A. heterophyllus* are recommended as pioneer species in restoration based on sun adaptation and high SLA, with two species, i.e., *C. sintoc* and *D. zibethinus* should be accompanied with intensive weeds control, while other native species, i.e., *C. schefferi*, *G. dulcis*, *D. longan*, and *D. celebica* are recommended as inserted species. Besides, we recommend using of high-performing plant species in restored habitats. We also demonstrate that the study on eco-physiological traits takes an important role in plant species selection, which serves as a critical stage in successful restoration programs.

Acknowledgments

The authors would like to thank the In-house Research scheme conducted by the Research Centre of Plant Conservation, Botanic Garden, and Forestry–National Research and Innovation Agency (BRIN), Indonesia, and several technicians in Purwodadi Botanic Garden.

Conflict of Interest

The authors declare no conflict of interest.

Author contribution

All authors equally contributed as main contributors to this study. All authors developed the conceptual ideas and the outline, conducted the literature reviews, were involved in data collection, drafted the manuscript, and revised, offered critical feedback on each section and finalized the manuscript. The authors specifically focused on particular work: R.R. worked more for the research concept; A.P.F. provided data collection and performed the analysis; D.A.L. carried out the literature reviews, was involved in data collection, and interpreted the data; J.D. carried out the literature reviews, provided data collection, and interpreted the data; A.R. conducted the literature reviews and collected the data; T.Y. conducted the literature reviews, collected the data, and offered critical feedback on each section; D.I.J. offered critical feedback on each section, revised and finalized the manuscript. All authors have read and agreed to the published version of the manuscript.

References

- MONTFORT F., BÉGUÉ A., LEROUX L., BLANC L., GOND V., CAMBULE A.H., REMANE I.A., GRINAND C. From land productivity trends to land degradation assessment in Mozambique: Effects of climate, human activities and stakeholder definitions. *Land Degradation & Development*. **32** (1), 49, **2021**.
- ASSENATO F., DI LEGINIO M., D'ANTONA M., MARINOSCI I., CONGEDO L., RIITANO N., LUISE A., MUNAFÒ M. Land degradation assessment for sustainable soil management. *Italian journal of agronomy*. **15** (4), 299, **2020**.
- SCHULZE K., MALEK Ž., VERBURG P.H. How will land degradation neutrality change future land system patterns? A scenario simulation study. *Environmental Science & Policy*. **124**, 254, **2021**.
- VON RINTELEN K., ARIDA E., HÄUSER C. A review of biodiversity-related issues and challenges in megadiverse Indonesia and other Southeast Asian countries. *Research Ideas and Outcomes*. **3**, e20860, **2017**.
- HUANG C., ZHOU Z., PENG C., TENG M., WANG P. How is biodiversity changing in response to ecological restoration in terrestrial ecosystems? A meta-analysis in China. *Science of the Total Environment*. **650**, 1, **2019**.
- ROHR J.R., BERNHARDT E.S., CADOTTE M.W., CLEMENTS W.H. The ecology and economics of restoration. *Ecology and Society*. **23** (2), **2018**.
- DUCHICELA S.A., CUESTA F., PINTO E., GOSLING W.D., YOUNG K.R. Indicators for assessing tropical alpine rehabilitation practices. *Ecosphere*. **10** (2), e20595, **2019**.
- GIANNINI T.C., GIULIETTI A.M., HARLEY R.M., VIANA P.L., JAFFE R., ALVES R., PINTO C.E., MOTA N.F., CALDEIRA JR C.F., IMPERATRIZ-FONSECA V.L. Selecting plant species for practical restoration of degraded lands using a multiple-trait approach. *Austral Ecology*. **42** (5), 510, **2017**.
- BALAZS K.R., KRAMER A.T., MUNSON S.M., TALKINGTON N., STILL S., BUTTERFIELD B.J. The right trait in the right place at the right time: Matching traits to environment improves restoration outcomes. *Ecological Applications*. **30** (4), e02110, **2020**.
- MULER A.L., CANHAM C.A., VAN ETEN E.J., STOCK W.D., FROEND R.H. Using a functional ecology approach to assist plant selection for restoration of Mediterranean woodlands. *Forest Ecology and Management*. **424**, 1, **2018**.
- NAVARRO-CANO J.A., GOBERNA M., VERDÚ M. Using plant functional distances to select species for restoration of mining sites. *Journal of Applied Ecology*. **56** (10), 2353, **2019**.
- WANG C., LIU H., ZHU L., REN H., YAN J., LI Z., ZHANG H. Which traits are necessary to quickly select suitable plant species for ecological restoration? *Ecological Solutions and Evidence*. **2** (4), e12102, **2021**.
- ZEBERIO J.M., PÉREZ C.A. Rehabilitation of degraded areas in northeastern Patagonia, Argentina: Effects of environmental conditions and plant functional traits on performance of native woody species. *Journal of Arid Land*. **12** (4), 653, **2020**.
- LEGER E.A., BARGA S., AGNERAY A.C., BAUGHMAN O., BURTON R., WILLIAMS M. Selecting native plants for restoration using rapid screening for adaptive traits: methods and outcomes in a Great Basin case study. *Restoration Ecology*. **29** (4), e13260, **2021**.
- HOLMES P.M., ESLER K.J., GAERTNER M., GEERTS S., HALL S.A., NSIKANI M.M., RICHARDSON D.M., RUWANZA S. Biological invasions and ecological restoration in South Africa. B.W. van Wilgen et al. Eds., *Biological Invasions in South Africa, Invading Nature-Springer Series in Invasion Ecology* 14, pp 665–700, **2020**.
- WEIDLICH E.W., FLÓRIDO F.G., SORRINI T.B., BRANCALION P.H. Controlling invasive plant species in ecological restoration: A global review. *Journal of Applied Ecology*. **57** (9), 1806, **2020**.
- BALACHOWSKI J.A., VOLAIRE F.A. Implications of plant functional traits and drought survival strategies for ecological restoration. *Journal of Applied Ecology*. **55** (2), 631, **2018**.
- CARLUCCI M.B., BRANCALION P.H., RODRIGUES R.R., LOYOLA R., CIANCIARUSO M.V. Functional traits and ecosystem services in ecological restoration. *Restoration Ecology*. **28** (6), 1372, **2020**.
- NAM K.J., LEE E.J. Variation in leaf functional traits of the Korean maple (*Acer pseudosieboldianum*) along an elevational gradient in a montane forest in Southern Korea. *Journal of Ecology and Environment*. **42** (1), 33, **2018**.
- MIGLIAVACCA M., MUSAVI T., MAHECHA M.D., NELSON J.A., KNAUER J., BALDOCCHI D.D., PEREZ-PRIEGO O., CHRISTIANSEN R., PETERS J., ANDERSON K. The three major axes of terrestrial ecosystem function. *Nature*. **598** (7881), 468, **2021**.
- CORNELISSEN J.H., LAVOREL S., GARNIER E., DÍAZ S., BUCHMANN N., GURVICH D., REICH P.B., TER STEEGE H., MORGAN H., VAN DER HEIJDEN M. A handbook of protocols for standardised and easy measurement of plant functional traits worldwide. *Australian journal of Botany*. **51** (4), 335, **2003**.
- YE M., PENG S., LI Y. Intraspecific variation in photosynthetic nitrogen-use efficiency is positively related to photosynthetic rate in rice (*Oryza sativa* L.) plants. *Photosynthetica*. **57** (1), **2019**.
- TAN X., GUO X., GUO W., LIU S., DU N. Invasive *Rhus typhina* invests more in height growth and traits associated with light acquisition than do native and non-invasive alien shrub species. *Trees*. **32**, 1103, **2018**.

24. JUNAEDI D.I. Ecology of *Zanthoxylum acanthopodium*: Specific leaf area and habitat characteristics. *Biodiversitas Journal of Biological Diversity*. **20** (3), 732, **2019**.
25. RINDYASTUTI R., SANCAYANINGSIH R.P. The growth strategies analysis of ten woody plant species for effective revegetation. *Biotropia*. **25** (1), 43, **2018**.
26. DANARTO S.A., BUDIHARTA S., FAUZIAH F. Tree species preference and rehabilitation perspective by local community: Case study in Bondowoso, East Java, Indonesia. *Asian Journal of Forestry*. **3** (2), **2019**.
27. RINDYASTUTI R., HAPSARI L. Adaptasi ekofisiologi terhadap iklim tropis kering: studi anatomi daun sepuluh jenis tumbuhan berkayu. *Jurnal Biologi Indonesia*. **13** (1), **2017**.
28. FIQA A.P., ARISOESILANINGSIH E. SOEJONO Konservasi Mata Air DAS Brantas Memanfaatkan Diversitas Flora Indonesia. Fakultas MIPA-Universitas Brawijaya, Malang, **2005**.
29. DANARTO S.A., YULISTYARINI T. Seleksi tumbuhan dataran rendah kering yang berpotensi tinggi dalam sekuestrasi karbon untuk rehabilitasi kawasan terdegradasi. *Pros Sem Nas Masy*. **5** (1), 33, **2019**.
30. YUDHOYONO A., SUKARYA D. 3500 plant species of the Botanic Gardens of Indonesia. PT. Sukarya dan Sukarya Pendetama. Jakarta. **2013**.
31. POWO Plants of The World Online. Royal Botanic Garden Kew. **2024**.
32. PEREZ-HARGUINDEGUY N., DIAZ S., GARNIER E., LAVOREL S., POORTER H., JAUREGUIBERRY P., BRET-HARTE M., CORNWELL W., CRAINE J., GURVICH D. New handbook for standardised measurement of plant functional traits worldwide. *Australian Journal of Botany*. **61**, 167, **2013**.
33. AGEHARA S., PRIDE L., GALLARDO M., HERNANDEZ-MONTERROZA J. A Simple, Inexpensive, and Portable Image-Based Technique for Nondestructive Leaf Area Measurements: HS1395, 11/2020. EDIS. **2020** (6), **2020**.
34. KHAN R., AHMAD M., ZAFAR M., ULLAH A. Scanning electron and light microscopy of foliar epidermal characters: A tool for plant taxonomists in the identification of grasses. *Microscopy Research and Technique*. **80** (10), 1123, **2017**.
35. OKTAVIA D., JIN G. Variations in leaf morphological and chemical traits in response to life stages, plant functional types, and habitat types in an old-growth temperate forest. *Basic and Applied Ecology*. **49**, 22, **2020**.
36. R CORE TEAM (2020) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. **2021**.
37. HAMMER Ø., HARPER D.A. Paleontological data analysis. John Wiley & Sons. **2024**.
38. POORTER L., BONGERS F. Leaf traits are good predictors of plant performance across 53 rain forest species. *Ecology*. **87** (7), 1733, **2006**.
39. PAŻ-DYDERSKA S., DYDERSKI M.K., NOWAK K., JAGODZIŃSKI A.M. On the sunny side of the crown—quantification of intra-canopy SLA variation among 179 taxa. *Forest Ecology and Management*. **472**, 118254, **2020**.
40. DÍAZ S., KATTGE J., CORNELISSEN J.H., WRIGHT I.J., LAVOREL S., DRAY S., REU B., KLEYER M., WIRTH C., COLIN PRENTICE I. The global spectrum of plant form and function. *Nature*. **529** (7585), 167, **2016**.
41. KUNSTLER G., FALSTER D., COOMES D.A., HUI F., KOOYMAN R.M., LAUGHLIN D.C., POORTER L., VANDERWEL M., VIEILLEDENT G., WRIGHT S.J. Plant functional traits have globally consistent effects on competition. *Nature*. **529** (7585), 204, **2016**.
42. HE D., CHEN Y., ZHAO K., CORNELISSEN J., CHU C. Intra- and interspecific trait variations reveal functional relationships between specific leaf area and soil niche within a subtropical forest. *Annals of Botany*. **121** (6), 1173, **2018**.
43. FANAL A., MAHY G., MONTY A. Can we foresee future maple invasions? A comparative study of performance-related traits and invasiveness of eight *Acer* species. *Plant Ecology*. **223** (10), 1181, **2022**.
44. EVANS J.R., CLARKE V.C. The nitrogen cost of photosynthesis. *Journal of Experimental Botany*. **70** (1), 7, **2019**.
45. YAO G. Q., NIE Z.F., ZENG Y.Y., WASEEM M., HASAN M.M., TIAN X.Q., LIAO Z.Q., SIDDIQUE K.H., FANG X.W. A clear trade-off between leaf hydraulic efficiency and safety in an aridland shrub during regrowth. *Plant, Cell & Environment*. **44** (10), 3347, **2021**.
46. ZHOU J., CIERAAD E., VAN BODEGOM P.M. Global analysis of trait–trait relationships within and between species. *New Phytologist*. **233** (4), 1643, **2022**.
47. BOONMAN C.C., BENÍTEZ-LÓPEZ A., SCHIPPER A.M., THUILLER W., ANAND M., CERABOLINI B.E., CORNELISSEN J.H., GONZALEZ-MELO A., HATTINGH W.N., HIGUCHI P. Assessing the reliability of predicted plant trait distributions at the global scale. *Global Ecology and Biogeography*. **29** (6), 1034, **2020**.
48. ARCHIBALD S., ALLINNE C., CERDÁN C.R., ISAAC M.E. From the ground up: Patterns and perceptions of herbaceous diversity in organic coffee agroecosystems. *Ecological Solutions and Evidence*. **3** (3), e12166, **2022**.
49. SIMPSON K.J., ATKINSON R.R., MOCKFORD E.J., BENNETT C., OSBORNE C.P., REES M. Large seeds provide an intrinsic growth advantage that depends on leaf traits and root allocation. *Functional Ecology*. **35** (10), 2168, **2021**.
50. LAWSON T., VIALET-CHABRAND S. Speedy stomata, photosynthesis and plant water use efficiency. *New Phytologist*. **221** (1), 93, **2019**.
51. LI S., WANG H., GOU W., WHITE J.F., KINGSLEY K.L., WU G., SU P. Leaf functional traits of dominant desert plants in the Hexi Corridor, Northwestern China: Trade-off relationships and adversity strategies. *Global Ecology and Conservation*. **28**, e01666, **2021**.
52. BALA K., SOOD A., PATHANIA V.S., THAKUR S. Effect of plant nutrition in insect pest management: A review. *Journal of Pharmacognosy and Phytochemistry*. **7** (4), 2737, **2018**.
53. BLUMENTHAL D.M., MUELLER K.E., KRAY J.A., OCHELTREE T.W., AUGUSTINE D.J., WILCOX K.R. Traits link drought resistance with herbivore defence and plant economics in semi-arid grasslands: The central roles of phenology and leaf dry matter content. *Journal of Ecology*. **108** (6), 2336, **2020**.
54. POPOVIĆ Z., BOJOVIĆ S., MARKOVIĆ M., CERDÀ A. Tree species flammability based on plant traits: A synthesis. *Science of the Total Environment*. **800**, 149625, **2021**.
55. FIORENTINI M., ZENOBI S., GIORGINI E., BASILI D., CONTI C., PRO C., MONACI E., ORSINI R. Nitrogen and chlorophyll status determination in durum wheat as influenced by fertilization and soil management: Preliminary results. *PloS One*. **14** (11), e0225126, **2019**.
56. LEGHARI S.J., WAHOCHO N.A., LAGHARI G.M., HAFEEZLAGHARI A., MUSTAFABHABHAN G.,

- HUSSAINTALPUR K., BHUTTO T.A., WAHOCHO S.A., LASHARI A.A. Role of nitrogen for plant growth and development: A review. *Advances in Environmental Biology*. **10** (9), 209, **2016**.
57. RAZAQ M., ZHANG P., SHEN H.-L., SALAHUDDIN. Influence of nitrogen and phosphorous on the growth and root morphology of *Acer mono*. *PloS One*. **12**, (2), e0171321, **2017**.
 58. MATHUR S., JAIN L., JAJOO A. Photosynthetic efficiency in sun and shade plants. *Photosynthetica*. **56**, 354, **2018**.
 59. GOUD E.M., AGRAWAL A.A., SPARKS J.P. A direct comparison of ecological theories for predicting the relationship between plant traits and growth. *Ecology*. **104** (4), e3986, **2023**.
 60. WILLIAMS G., NELSON A. Spatial variation in specific leaf area and horizontal distribution of leaf area in juvenile western larch (*Larix occidentalis* Nutt.). *Trees*. **32**, 1621, **2018**.
 61. ALAM M.A., WYSE S.V., BUCKLEY H.L., PERRY G.L., SULLIVAN J.J., MASON N.W., BUXTON R., RICHARDSON S.J., CURRAN T.J. Shoot flammability is decoupled from leaf flammability, but controlled by leaf functional traits. *Journal of Ecology*. **108** (2), 641, **2020**.
 62. WÜEST R.O., MÜNKEMÜLLER T., LAVERGNE S., POLLOCK L.J., THUILLER W. Integrating correlation between traits improves spatial predictions of plant functional composition. *Oikos*. **127** (3), 472, **2018**.
 63. WIGLEY B.J., FRITZ H., COETSEE C. Defence strategies in African savanna trees. *Oecologia*. **187**, 797, **2018**.
 64. HELSEN K., MATSUSHIMA H., SOMERS B., HONNAY O. A trait-based approach across the native and invaded range to understand plant invasiveness and community impact. *Oikos*. **130** (6), 1001, **2021**.
 65. MILANOVIĆ M., KNAPP S., PYŠEK P., KÜHN I. Linking traits of invasive plants with ecosystem services and disservices. *Ecosystem Services*. **42**, 101072, **2020**.
 66. POORTER L., ROZENDAAL D.M. Leaf size and leaf display of thirty-eight tropical tree species. *Oecologia*. **158**, 35, **2008**.
 67. FALSTER D.S., DUURSMA R.A., FITZJOHN R.G. How functional traits influence plant growth and shade tolerance across the life cycle. *Proceedings of the National Academy of Sciences*. **115** (29), E6789, **2018**.
 68. LESTARI D.A., FIQA A.P., FAUZIAH F., BUDIHARTA S. Growth evaluation of native tree species planted on post coal mining reclamation site in East Kalimantan, Indonesia. *Biodiversitas Journal of Biological Diversity*. **20** (1), 134, **2019**.
 69. LAMPELA M., JAUHIAINEN J., SARKKOLA S., VASANDER H. To treat or not to treat? The seedling performance of native tree species for reforestation on degraded tropical peatlands of SE Asia. *Forest Ecology and Management*. **429**, 217, **2018**.