

Original Research

Heavy Metals Pollution Evaluation, Health Risk Assessment, and Source Identification in the Reservoirs, Northern Anhui Province, China

Yu Liu^{1, 2, 3}, Jie Ma^{4, 5*}, Kai Chen^{1, 5}, Qimeng Liu^{1, 5}, Huichan Chai⁵, Haitao Zhang⁴

¹The Key Laboratory of Universities in Anhui Province for Prevention of Mine Geological Disasters, Anhui University of Science and Technology, Huainan 232001, China

²Joint National-Local Engineering Research Centre for Safe and Precise Coal Mining, Anhui University of Science and Technology, Huainan 232001, China

³State Key Laboratory of Mining Response and Disaster Prevention and Control in Deep Coal Mines, Anhui University of Science and Technology, Huainan 232001, China

⁴School of Resources and Civil Engineering, Suzhou University, Suzhou 234000, China

⁵School of Earth and Environment, Anhui University of Science and Technology, Huainan 232001, China

Received: 11 May 2024

Accepted: 15 August 2024

Abstract

Reservoirs, distinctive water bodies, serve as vital water sources for irrigation, electricity generation, and aquaculture. In this study, 31 water samples from 16 reservoirs in Northern Anhui Province's Jiagou region were collected to reveal their pollution status, health risks, and sources. The decreasing order of the heavy metal concentrations is as follows: Mn > Cr > Ni > As > Zn > Cu > Mo > Co. The heavy metal pollution index values of all the samples are below 100, indicating the reservoirs' water quality is within acceptable standards. The Nemerow comprehensive pollution indicates that M4 and M17 are classified as slightly polluted categories, while the remaining points belong to the non-polluted categories. The USEPA health risk assessment indicates that Cr is higher than the maximum acceptable risk levels recommended by certain agencies; the reservoirs' samples are not suitable as drinking water resources. Correlation analysis and principal component analysis both suggest that the sources of Co, Ni, Cu, and Cr may be attributed to human activities involving pesticides and fertilizers. Mo, As, and Zn may originate from coal combustion, while Mn may be derived from pesticide use. These findings offer valuable insights for environmental policymakers and inform decision-making for ecological protection.

Keywords: heavy metals, pollution assessment, health risk assessment, source identification, reservoir

*e-mail: ahszumajie@163.com

Introduction

Water constitutes the fundamental basis for the existence of the Earth's organisms. Water resources play a crucial role in maintaining the planet's ecological environment and are essential for the sustainable development of the global environment. The water resources encompass surface water and groundwater, and these water resources can be utilized for irrigation, power generation, water supply, shipping, aquaculture, and other purposes. The surface water bodies include rivers, lakes, wells, springs, tides, and reservoirs. Wherein reservoirs are a unique artificial construction that functions as water storage, electric power generation, tourism, and aquaculture. With rapid urbanization, industrialization, and agricultural development, the demands for clean and safe water resources have escalated, leading to a global water shortage. Many countries prioritize the development of water conservation projects, investing heavily in the construction of reservoirs, embankments, and various water storage facilities to balance supply and demand. For example, China has constructed 97,264 reservoirs with a storage capacity of approximately 810.41 billion m³, and 756 new reservoirs are being built with a capacity of 12.190 billion m³ [1, 2].

Nowadays, reservoirs have increasingly served as drinking water sources for major cities, and they play a significant role in agricultural water use, hydropower generation, flood control and disaster prevention, and ecological preservation. However, reservoir construction also poses potential adverse effects, including geological disasters and ecological damage. For example, reservoirs may induce and increase the frequency of earthquakes in the reservoir's area and surrounding regions [3]. Reservoir construction can lead to the gradual reduction of biological species, hindering biodiversity. Industrial and agricultural activities can release heavy metals into water bodies, resulting in their accumulation in reservoirs. Heavy metals are persistent pollutants characterized by bioaccumulation, environmental persistence, and toxicity. Even more, some certain heavy metals (e.g., Cd, Cr, As, and Pb) exhibit toxic effects even at low concentrations [4, 5].

Numerous studies have investigated conventional ions, isotopes, and heavy metals in various water bodies [6–8]. With rapid industrialization, heavy metal accumulation in these water bodies has escalated, posing significant threats to human health and the ecological environment [9]. Consequently, researchers have emphasized heavy metal pollution in different water bodies, including reservoirs [10]. Various heavy metal pollution assessment methods have been proposed, such as the single-factor pollution index method [11], the Nemerow pollution index method [12], the heavy metal pollution index method (HPI) [13], the heavy metal evaluation index method, and the potential ecological risk index method [14]. Ecological risk assessment of heavy metals can be conducted using the United States Environmental Protection Agency's (USEPA) health risk assessment model (AHP model) [15]. Principal component analysis (PCA) [16], correlation analysis, positive matrix

factorization (PMF) [17], cluster analysis, unmix analysis, and weighted alternating least squares (MCR-WALS) are commonly used to identify heavy metal sources in water bodies.

Extensive research has been conducted on rivers, lakes, subsidence pools, shallow groundwater, and deep groundwater in Northern Anhui Province, China, and this research has focused on investigating their hydrochemical characteristics, heavy metal pollution, and isotope features [18–21]. However, studies on reservoirs remain relatively limited. Therefore, the objectives of this study aim to 1) characterize the concentrations and their variation of heavy metals (Mn, Zn, Co, As, Mo, Cu, Cr, and Ni), 2) evaluate heavy metal pollution using the Nemerow composite pollution index and the HPI, 3) assess health risks to adults and children using the AHP model, and 4) identify the heavy metal sources through correlation and principal component analyses.

Materials and Methods

Research Area

The study area is located in the Jiagou Region, Suzhou City, Northern Anhui Province, China (117°3'34"E, 33°53'27"N), covering an area of 178 square kilometers. Suzhou City comprises Yongqiao District, Dangshan County, Xiaoxian County, Lingbi County, and Sixian County, and the study area is situated within Yongqiao District, which can be seen in Fig. 1. The region experiences a warm temperate monsoon climate, with an average annual precipitation of 1200 mm, primarily concentrated between June and August. Annual sunshine hours range from 250 to 2500. The study area lies in the center of the Huaibei Plain, characterized by diverse geomorphological features, including hills, terraces, and plains; the Jiagou district is located in the plain area. Agriculture, small-scale industry, and tourism constitute the primary industries in the region. Commonly, the loose strata develop three aquifers and two aquicludes; the upper confined aquifer is the primary water supply source among these three aquifers. Plenty of reservoirs have been constructed for irrigation, domestic living, shipping, and local aquaculture.

Sample Collection and Testing

A total of 32 water samples were collected from both sides of 16 reservoirs in the Jiagou region. During sampling, GPS positioning was used to determine the coordinates of each sampling point. Additional information was recorded, including in-situ conditions (e.g., temperature, pH, traffic conditions, topography, crops, and industrial facilities). The sampling locations are depicted in Fig. 1. At each sampling point, the sampling bottle was triple-rinsed with the original water. The water sample was then collected in a 500-ml polyethylene bottle, sealed, and then transported to the Key Laboratory of Mine Water Resource Utilization of Anhui Higher Education Institutes, Suzhou




Fig. 1. Sampling points in the study area.

City, Anhui Province. In the laboratory, the water sample was filtered through a 0.22- μ m microporous water filtration membrane and acidified with 5% concentrated nitric acid, guaranteeing a pH below 2. The acidified sample with 97 g, the concentrated nitric acid with 2 g, and the 500 ppb Rh internal standard with 1 g were added together to a 100-ml volumetric flask. Finally, inductively coupled plasma mass spectrometry (ICP/MS) was employed to measure the concentrations of Co, Cu, Zn, Mo, As, Cr, Mn, and Ni. Finally, only thirty-one samples with testing precision of less than 10% were considered for further analysis, which can offer the original data for studying the concentrations and variation characteristics, pollution assessment, and the source apportionment of the heavy metals.

Analysis Methods

Descriptive Statistical Method

Descriptive statistics provide a statistical summary of relevant data collected during a survey. Complex datasets can be described by using a few representative indices, allowing for an intuitive understanding of data variations. In this study, descriptive statistics were used to analyze the minimum value, maximum value, mean value, coefficient of variation (C.V.), and standard deviation of the data. These indices can also be employed to assess the extent to which human activities influence heavy metal enrichment in different water bodies [22].

The Heavy Metal Pollution Index Method (HPI)

The heavy metal pollution index method (HPI) is assessed based on weighted arithmetic mean. It evaluates the pollution degree of various heavy metals, and the calculation involves the following steps [23].

$$W_i = k/S_i \quad (1)$$

W_i is the i^{th} heavy metal indicator weight and k is the constant of proportionality; usually, it is 1. S_i is the recommended standard for the i^{th} parameter.

$$Q_i = 100 \times C_i/S_i \quad (2)$$

Q_i is the quality grade index of the i^{th} heavy metal index. C_i is the tested concentration of the i^{th} water sample's heavy metal element.

$$HPI = \frac{\sum_{i=1}^n (Q_i W_i)}{\sum_{i=1}^n W_i} \quad (3)$$

The acceptable contamination value is 100, and more than 100 is considered serious pollution.

The Nemerow Composite Pollution Index

The Nemerow composite pollution index, which highlights the effect of heavy metal pollutants on water quality, is calculated by using the average and maximum values of the single-factor pollution index, and the calculation formula is as follows:

$$P_F = \sqrt{\frac{P_{\text{imax}}^2 + P_a^2}{2}} \quad (4)$$

In the formula, P_I is synthesized as a comprehensive pollution index, P_a is the average of individual pollution indexes of the single pollution index, and P_{imax} is the largest value of the single pollution index. The evaluation indicators are defined as follows: $P_I < 1.0$ indicates no pollution; $1.0 \leq P_I < 2.0$ suggests slight pollution; $2.0 \leq P_I < 3.0$ implies moderate pollution; $P_I \geq 3.0$ represents heavy pollution [24].

The Health Risk Assessment (AHP Model)

Numerous studies have demonstrated that the primary routes of heavy metal exposure in humans are through drinking water ingestion and skin contact. Excessive exposure to heavy metals can increase the risk of cancer [25]. In this study, the health risk assessment model recommended by the USEPA is employed to evaluate health risks for adults (male and female) and children. Heavy metals in the reservoir can be classified into two categories: chemical non-carcinogenic elements (Mn, Co, Ni, Cu, Zn, and Mo) and chemical carcinogenic elements (Cr and As).

The calculation methods for each category vary based on the exposure route:

For drinking water pathway:

R_i^c is the annual carcinogenic risk caused by drinking water. The calculation formula is as follows:

$$R_i^c = \frac{1 - \exp(-D_i \times SF)}{L} \quad (5)$$

D_i represents the average daily exposure dose per unit weight of drinking water route. The calculation formula is as follows:

$$D_i = \frac{\theta C}{W} \quad (6)$$

R_i^n is the annual non-carcinogenic risk caused by drinking water. The calculation formula is as follows:

$$R_i^n = \frac{D_i}{Rf D_i \times L} \times 10^{-6} \quad (7)$$

For skin infiltration pathway:

R_i^p represents the annual risk of carcinogens caused by the skin pathway, which can be determined according to formula (8).

$$R_i^p = \frac{1 - \exp(-CDI \times SF)}{L} \quad (8)$$

R_i^f represents the annual risk of non-carcinogenics caused by the skin pathway. It can be determined according to formula (9):

$$R_i^f = \frac{CDI}{Rf D_i \times L} \times 10^{-6} \quad (9)$$

CDI is the daily dose of heavy metal element d per unit of body weight, and its unit is $\text{mg}/(\text{kg} \cdot \text{d})$, which can be formulated as formula (10).

$$CDI = \frac{I \times ASD \times EF \times FE \times ED}{W \times AT \times f} \quad (10)$$

I is the adsorption amount of heavy metal element per unit area of each bath, and its unit is $\text{mg}/(\text{cm}^2 \cdot \text{times})$. It can be determined according to formula (11).

$$I = 2 \times 10^{-6} \times k \times C_i \times \sqrt{\frac{6 \times \tau \times TE}{\pi}} \quad (11)$$

The relevant parameter values of the above formulas are detailed in Table 1 [26], and the non-carcinogenic reference dose and carcinogenic intensity coefficient are shown in Table 2 [27–29]. The maximum acceptable risk level and negligible risk level recommended by relevant research institutions can be obtained in Table 3 [30]. In accordance with Table 3, it can be concluded that the maximum acceptable risk level of some institutions is between 1×10^{-6} and 1×10^{-4} .

Table 1. The practical significance, reference values, and units of each symbol in the formula.

Sign	Practical significance	Reference values	Units
RfD	Reference dose of drinking water exposure	Calculated according to the formula	mg/(kg*d)
SF	The carcinogenic slope coefficient of the substance	Consult data	(kg*d)/mg
C _i	The exposure concentration of pollutants in the study area	Measured value	ug/L
L	Mean age	80	a
θ	Daily water intake	Adult 2.2; children 1.0	L/d
EF	Exposure frequency	365	d/a
ED	Exposure duration	Non-carcinogenic: 35a Carcinogens: 70a	a
W	Average weight	Adult: Male 63.3; Female 55.1 Children: 25	kg
AT	Average exposure time	Non-carcinogenic: 35a Carcinogens: 70a	d
ASD	The surface area of the human body	Adult 18000; children 6660	cm ²
FE	Bath frequency	0.3	times/d
TE	Bath time	0.4	h
f	Adsorption frequency in the intestine	1	zero dimension
k	Adsorption parameters of skin	0.001	cm/h
τ	Pollutant retention time	1	h

Table 2. Contaminant carcinogenic slope factor SF and non-carcinogen reference RfD.

Component	SF/(kg*d)/mg	RfD/mg/(kg*d)
As	15	-
Cr	41	-
Ni	-	0.02
Mn	-	1.4
Co	-	0.0003
Cu	-	0.005
Zn	-	0.3
Mo	-	0.005

Table 3. The maximum acceptable risk level and negligible risk level are recommended by some relevant research institutions.

Different agency	Maximum acceptable risk level/a ⁻¹	Negligible risk level/a ⁻¹
Swedish Environmental Protection Agency	1×10 ⁻⁶	-
Ministry of Construction and Environment of the Netherlands	1×10 ⁻⁶	1×10 ⁻⁸
Royal Society of England	1×10 ⁻⁶	1×10 ⁻⁷
USEPA	1×10 ⁻⁴	-

Correlation Analysis

Correlation analysis is a statistical technique that can be used to assess the relationship between two or more correlated variables. By eliminating the influence of extraneous factors, this method retains objective and reliable information. It is commonly employed in source apportionment studies of water samples. The strength of the correlation is indicated by the correlation coefficient, which ranges from -1 to +1. A correlation coefficient between 0.6 and 0.8 indicates a positive correlation, while a correlation coefficient greater than 0.8 suggests a strong positive correlation, indicating these two elements may have originated from the same parent material. Conversely, a negative correlation coefficient indicates an inverse relationship between the variables, suggesting that these two elements are derived from different parent materials [30].

Principal Component Analysis

PCA is a multivariate statistical technique that reduces a large number of correlated variables into a smaller set of uncorrelated variables, known as principal components. These components capture most of the variance in the original data. PCA is commonly used with SPSS software to transform multiple indicators into a few comprehensive indicators or to identify major pollution sources from multiple pollution sources. Based on the contribution rate of each principal component, the main factors can be identified, providing insights into the primary sources of the original elements [26].

In this study, the descriptive statistics using SPSS 17.0 are employed in delineating the concentration and variation characteristics; the HPI and the Nemerow composite pollution index are used for evaluating the pollution status; the health risk assessment (AHP model) are conducted for assessing the health risks for adults (male and female) and children through drinking water ingestion and skin contact; and finally, the correlation analysis and the principal component analysis by using SPSS 17.0 are utilized for the source apportionment of the heavy metals.

Results and Discussion

Descriptive Statistics

The concentrations in 8 heavy metals (Co, Cu, Zn, Mo, As, Cr, Mn, and Ni) of 31 samples from 16 reservoirs are listed in Table 4. The descending order of average concentrations are: Mn ($20.08 \mu\text{g}\cdot\text{L}^{-1}$) > Cr ($7.75 \mu\text{g}\cdot\text{L}^{-1}$) > Ni ($3.57 \mu\text{g}\cdot\text{L}^{-1}$) > As ($2.34 \mu\text{g}\cdot\text{L}^{-1}$) > Zn ($1.82 \mu\text{g}\cdot\text{L}^{-1}$) > Cu ($0.94 \mu\text{g}\cdot\text{L}^{-1}$) > Mo ($0.87 \mu\text{g}\cdot\text{L}^{-1}$) > Co ($0.29 \mu\text{g}\cdot\text{L}^{-1}$). Among these heavy metals, the element of Mn is within the widest concentration range. The average values of Mn, Mo, Co, and Ni are below the standard limits; the maximum values of Mn and Ni exceed the limits for drinking water. The average values of Cu, Zn, and As do not exceed Class I of the environmental quality standard for surface water [31, 32], but the average Cr concentration reaches Class V. Fig. 2 illustrates the percentage contribution of each heavy metal at different sampling points. As can be seen from Fig. 2, Cr and Ni have significantly higher contributions in sites 1–12 compared with other sites.

The coefficient of variation (C.V.) reflects the dispersion of heavy metal concentrations; higher C.V. values indicate greater dispersion [33, 34]. Based on previous studies [34], variability can be classified as follows: Low variability: $\text{C.V.} \leq 20\%$, moderate variability: $20\% < \text{C.V.} \leq 50\%$, and high variability: $\text{C.V.} > 50\%$. As and Zn exhibit moderate variability, while Mn, Co, Ni, Cu, Mo, and Cr show high variability. This suggests a significant spatial variation in heavy metal concentrations, which is potentially influenced by pollution sources.

Pollution Assessment

Fig. 3 presents the HPI values for each sampling point. As shown in Fig. 3, all values were below 100, indicating that these samples are within acceptable levels of heavy metal pollution in the 16 reservoirs [35]. All of the samples are below the national standard limit, and these samples can be classified as acceptable levels. However, among these samples, M4 and M19 have a trend of relatively elevated HPI values, which may be related to pesticide use in the study region.

Table 4. Analysis and statistics of heavy metal concentration in Jiagou Reservoir.

Item	Mn	Co	Ni	Cu	Zn	Mo	Cr	As
Min/ $\mu\text{g}\cdot\text{L}^{-1}$	0.73	0.09	0.23	0.11	0.28	0.22	0	0.79
Max/ $\mu\text{g}\cdot\text{L}^{-1}$	237.79	0.81	21.93	2.48	3.76	2.48	80.88	4.37
Mean/ $\mu\text{g}\cdot\text{L}^{-1}$	20.08	0.29	3.57	0.94	1.82	0.87	7.75	2.34
SD/ $\mu\text{g}\cdot\text{L}^{-1}$	46.44	0.17	5.25	0.6	0.8	0.62	17.27	0.98
C.V.	231.27	58.62	147.06	63.83	43.96	71.26	222.84	41.88




Fig. 2. Heavy metal content histogram of each sampling points.




Fig. 3. Heavy metal pollution index histogram of reservoirs' heavy metals.

Simultaneously, the results of the Nemerow comprehensive pollution assessment are shown in Fig. 4. The descending order of the Nemerow comprehensive pollution index is: $M17 > M4 > M18 > M9 > M2 > M6 > M21 > M3 > M11 > M10 > M12 > M13 > M7 > M5 > M15 > M28 > M32 > M31 > M27 > M22 > M19 > M29 > M20 > M30 > M16 > M14 > M23 > M24 > M25 > M26$. M4 and M17 are classified as slightly polluted categories, while the remaining points belong to the non-polluted categories. Compared with the results of other related research on groundwater [24], the reservoirs' samples and the groundwater samples are both within a good water quality environment. Meanwhile, these results are consistent with the results of the HPI, suggesting the results

of these two pollution evaluation methods can be mutually corroborated and credible.

Health Risk Assessment

According to different water acquisition routes, the USEPA health risk assessment model commonly considers selecting different parameters. In this study, this model is employed to calculate the range and the average of annual health risks posed by heavy metals in water bodies through skin contact and drinking water, which are presented in Tables 5 and 6, respectively. Table 5 and Fig. 5(a) summarize the calculated health risks for males, females, and children via skin contact. Except for




Fig. 4. Nemerow pollution index histogram of reservoir s' heavy metals.

Table 5. The average annual health risk value of skin contact pathway.

Skin Contact	Males			Females			Children		
	Min	Max	Mean	Min	Max	Mean	Min	Max	Mean
Mn	3.57×10^{-13}	1.16×10^{-10}	9.72×10^{-12}	4.10×10^{-13}	1.33×10^{-10}	1.12×10^{-11}	3.34×10^{-13}	1.08×10^{-10}	9.11×10^{-12}
Co	1.98×10^{-10}	4.13×10^{-9}	7.67×10^{-10}	2.28×10^{-10}	4.75×10^{-9}	8.81×10^{-10}	1.86×10^{-10}	3.87×10^{-9}	7.19×10^{-10}
Ni	7.82×10^{-12}	2.41×10^{-9}	1.93×10^{-10}	8.99×10^{-12}	2.77×10^{-9}	2.22×10^{-10}	7.33×10^{-12}	2.26×10^{-9}	1.81×10^{-10}
Cu	1.48×10^{-11}	9.88×10^{-10}	1.54×10^{-10}	1.70×10^{-11}	1.14×10^{-9}	1.77×10^{-10}	1.39×10^{-11}	9.26×10^{-10}	1.44×10^{-10}
Zn	6.44×10^{-13}	9.49×10^{-12}	4.30×10^{-12}	7.40×10^{-13}	1.09×10^{-11}	4.94×10^{-12}	6.03×10^{-13}	8.89×10^{-12}	4.03×10^{-12}
Mo	2.96×10^{-11}	3.38×10^{-10}	1.25×10^{-10}	3.40×10^{-11}	3.88×10^{-10}	1.43×10^{-10}	2.77×10^{-11}	3.17×10^{-10}	1.17×10^{-10}
Cr	0	4.44×10^{-3}	3.38×10^{-4}	0	4.94×10^{-3}	3.82×10^{-4}	0	4.21×10^{-3}	3.19×10^{-4}
As	8.10×10^{-6}	4.46×10^{-5}	2.37×10^{-5}	9.31×10^{-6}	5.12×10^{-5}	2.72×10^{-5}	7.59×10^{-6}	4.18×10^{-5}	2.22×10^{-5}




Fig. 5. The box plot of carcinogenic risk caused by the skin contact route (a) and the drinking water route (b).

Table 6. The average annual health risk value of drinking water pathway.

Drinking Water	Males			Females			Children		
	Min	Max	Mean	Min	Max	Mean	Min	Max	Mean
Mn	2.28×10^{-13}	7.38×10^{-11}	6.23×10^{-12}	2.62×10^{-13}	8.48×10^{-11}	7.16×10^{-12}	2.62×10^{-13}	8.50×10^{-11}	7.17×10^{-12}
Co	1.27×10^{-10}	1.18×10^{-9}	4.20×10^{-10}	1.46×10^{-10}	1.35×10^{-9}	4.83×10^{-10}	1.46×10^{-10}	1.36×10^{-9}	4.84×10^{-10}
Ni	4.99×10^{-12}	4.76×10^{-10}	7.76×10^{-11}	5.737×10^{-12}	5.47×10^{-10}	8.91×10^{-11}	5.748×10^{-12}	5.48×10^{-10}	8.93×10^{-11}
Cu	9.47×10^{-12}	2.15×10^{-10}	8.12×10^{-11}	1.09×10^{-11}	2.47×10^{-10}	9.33×10^{-11}	1.090×10^{-11}	2.48×10^{-10}	9.35×10^{-11}
Zn	4.11×10^{-13}	5.45×10^{-12}	2.64×10^{-12}	4.72×10^{-13}	6.26×10^{-12}	3.03×10^{-12}	4.731×10^{-13}	6.27×10^{-12}	3.04×10^{-12}
Mo	1.89×10^{-11}	2.16×10^{-10}	7.56×10^{-11}	2.17×10^{-11}	2.48×10^{-10}	8.69×10^{-11}	2.172×10^{-11}	2.48×10^{-10}	8.71×10^{-11}
Cr	0	1.36×10^{-3}	1.34×10^{-4}	0	1.550×10^{-3}	1.529×10^{-4}	0	1.55×10^{-3}	1.53×10^{-4}
As	5.17×10^{-6}	2.85×10^{-5}	1.52×10^{-5}	5.94×10^{-6}	3.27×10^{-5}	1.75×10^{-5}	5.95×10^{-6}	3.28×10^{-5}	1.75×10^{-5}

Mn, children exhibit slightly lower carcinogenic and non-carcinogenic risks compared with females. The average annual non-carcinogenic health risks for adults associated with the six non-carcinogenic heavy metals (Mo, Ni, Mn, Co, Cu, and Zn) are all below the negligible risk level of 1×10^{-7} recommended by the Royal Society. Therefore, the health risks posed by these six heavy metals for adults can be considered negligible.

Regarding carcinogenic risks, the average annual health risk and range for As are lower than the acceptable risk level of 1×10^{-4} recommended by the USEPA. However, the health risks associated with heavy metal Cr are higher than those for As. Notably, the risk value for Cr exceeds 1×10^{-4} at some sampling sites, indicating that the carcinogenic risk posed by Cr requires further attention [36].

Table 6 and Fig. 5(b) present the calculated health risks for males, females, and children via drinking water. Children exhibit slightly higher carcinogenic and non-carcinogenic risks than adults through this route [37]. The average annual health risks associated with Cr exceed the maximum acceptable risk levels set by all relevant agencies, indicating its potential carcinogenic risk. The health risk value for As is lower than the maximum acceptable risk level recommended by the USEPA but higher than the limits established by the Swedish Environmental Protection Agency, the Dutch Ministry of Construction and Environment, and the Royal Society. By contrast, the health risk values for Ni, Mn, Co, Cu, Zn, and Mo are significantly below the maximum acceptable risk levels set by relevant agencies, which means that their health risks are negligible.

Overall, the descending order of health risks to humans posed by the eight heavy metals was $\text{Cr} > \text{As} > \text{Co} > \text{Cu} > \text{Ni} > \text{Mo} > \text{Mn} > \text{Zn}$. Owing to Cr being higher than the maximum acceptable risk levels recommended by certain agencies, the reservoirs in the study area are not suitable as drinking water resources. The health risks associated with heavy metal exposure via skin contact

were higher than those via drinking water [38]. Notably, the health risk values for Cr and As exceeded the maximum acceptable risk level at certain sampling points; these points may be related to human activities, such as the application of chemical fertilizers and pesticides in surrounding agricultural areas.

Correlation Analysis

The correlation analysis of heavy metal concentrations can be implied to reveal the relationship between two elements. When they have a strong positive correlation coefficient, this indicates that these two elements have homology; on the contrary, a strong negative correlation coefficient between two elements shows a non-homology. As shown in Table 7, there are strong positive correlations between Ni and Cr (0.943), Co and Ni (0.887), and Cr and Co (0.858). Mo and As show a moderately positive correlation coefficient of 0.721, suggesting that these elements may have originated from similar and/or the same sources. Conversely, negative correlation coefficients are observed between Mn and Ni (-0.079), Mn and Mo (-0.169), Mn and Cr (-0.03), and Mn and As (-0.066). These negative correlation coefficients indicate that these elements may originate from different sources.

Principal Component Analysis

In this study, the principal component analysis (PCA) is applied to reveal the source of these heavy metals. As can be seen from Table 8, with rotation and an eigenvalue greater than 1, three principal components can be summarized. Principal component 1 (PC1) accounts for 43.05% of the variance, and the high loadings on Co, Ni, Cu, and Cr indicate a strong positive correlation among these heavy metals. PC2 accounting for 23.15% of the variance exhibits a high loading on Mo and As, with a moderate correlation with Zn, which suggests a relatively

Table 7. Correlation analysis of heavy metals.

Heavy metals	Mn	Co	Ni	Cu	Zn	Mo	Cr	As
Mn	1							
Co	0.046	1						
Ni	-0.079	0.887**	1					
Cu	0.098	0.555**	0.628**	1				
Zn	0.051	0.297	0.178	0.202	1			
Mo	-0.169	0.378*	0.127	-0.02	0.236	1		
Cr	-0.03	0.858**	0.943**	0.547**	0.198	0.148	1	
As	-0.066	0.359*	0.056	-0.102	0.323	0.721**	0.043	1

Note: * indicates a significant correlation at the 0.05 level and ** indicates a significant correlation at the 0.01 level.

Table 8. Principal component analysis of heavy metals.

Component	Before Rotation			After Rotation		
	FC1	FC2	FC3	VF1	VF2	VF3
Mn	-0.02	-0.22	0.88	-0.02	-0.12	0.90
Co	0.96	0.04	0.02	0.88	0.37	0.06
Ni	0.93	-0.27	-0.16	0.97	0.06	-0.09
Cu	0.67	-0.40	0.14	0.76	-0.12	0.22
Zn	0.39	0.34	0.47	0.21	0.50	0.44
Mo	0.38	0.81	-0.09	0.08	0.88	-0.17
Cr	0.91	-0.25	-0.13	0.94	0.07	-0.05
As	0.31	0.86	0.10	-0.01	0.92	0.00
Eigen Values	3.44	1.85	1.07	3.24	2.04	1.09
Var/%	43.05	23.15	13.39	40.46	25.52	13.61
Cum/%	43.05	66.20	79.59	40.46	65.98	79.59

strong positive correlation. PC3 accounting for 13.39% shows a strong correlation with Mn. Combining related research results from the surrounding research regions and the recording information during the sampling [39, 40], PC1 may represent the influence of human activities such as pesticide and fertilizer use, PC2 may be associated with coal combustion, and PC3 may be related to local pesticide applications.

Conclusions

In this study, 31 surface water samples were collected from 16 reservoirs for studying the pollution evaluation, health risk assessment, and source apportionment

of the reservoirs' samples. The key findings are summarized as follows:

(1) The descending order of average concentrations are: Mn > Cr > Ni > As > Zn > Cu > Mo > Co. Mn, Co, Ni, Cu, Mo, and Cr show a high variability; this suggests that there is a significant spatial variation in heavy metal concentrations, potentially influenced by pollution sources.

(2) The heavy metal pollution index (HPI) values for all samples were below 100, indicating acceptable heavy metal pollution levels. The Nemerow comprehensive pollution assessment indicates that M4 and M17 are classified as slightly polluted categories, while the remaining points belong to the non-polluted categories.

(3) Health risk assessment reveals that Cr is higher than the maximum acceptable risk levels recommended by

certain agencies; the reservoirs' samples are not suitable as drinking water resources.

(4) Correlation analysis and the principal component analysis both identify three principal components: PC1 may represent the influence of human activities such as pesticide and fertilizer use, PC2 may be associated with coal combustion, and PC3 may be related to local pesticide applications.

(5) Regular water quality monitoring, attention to primary pollution sources, and identification of the source apportionment need to be engaged to guarantee the water quality of reservoirs.

Acknowledgments

This research was supported by Natural Science Research Project of Anhui Educational Committee (No. 2022AH050797), Open Research Grant of Joint National-Local Engineering Research Centre for Safe and Precise Coal Mining (No. EC2023012), Open Foundation of the Key Laboratory of Universities in Anhui Province for Prevention of Mine Geological Disasters (No. 2022-MGDP-01), Independent Research Fund of the State Key Laboratory of Mining Response and Disaster Prevention and Control in Deep Coal Mines (No. SKLMRDPC19ZZ06), the Coal Research Centre for Comprehensive Prevention and Control of Mine Water Hazard (134092220042204), the Key Scientific Research Project of Suzhou University (2022yzd01), and the Outstanding Academic and Technical Backbone of Suzhou University (2020XJGG11).

Conflict of Interest

The authors declare that they have no conflict of interest.

References

- ZANG N., ZHU J., WANG X., LIAO Y., CAO G., LI C., LIU Q., YANG Z. Eutrophication risk assessment considering joint effects of water quality and water quantity for a receiving reservoir in the South-to-North Water Transfer Project, China. *Journal of Cleaner Production*. **331**, 129966, **2022**.
- SUN Z., ZHANG L., DUAN Z. The number and distribution of reservoir projects in China. *China Water Conservancy*. **7**, 10, **2013** [In Chinese].
- NA J., CEN C., SHI Z., FENG B. Distinct response of deep and shallow geothermal reservoir temperature following earthquakes in the Kangding Fault zone hydrothermal system. *Journal of Hydrology: Regional Studies*. **52**, 101743, **2024**.
- ZHANG K., LI K., XIN R., XU Z., ZHANG Z., HE S., ZHAO Z., TONG M., CUI Y. Distribution pattern and ecological risk assessment of heavy metals in Henan section of the Yellow River. *Water Supply*. **22** (7), 6195, **2022**.
- RAHMAN Z., SINGH V.P. The relative impact of toxic heavy metals (THMs) (arsenic (As), cadmium (Cd), cadmium (Cr) (VI), mercury (Hg), and lead (Pb)) on the total environment: an overview. *Environmental Monitoring and Assessment*. **191** (7), 419, **2019**.
- CHEN J., ZHU M., LIU G., GAO M., LIU J., ZHANG W. Determining the factors controlling the chemical composition of groundwater using geochemical methods and groundwater health assessment in Qixing Town, southwest China. *Water Supply*. **23** (9), 3715, **2023**.
- KAUR S., MEHRA R., KUMAR M.R. Quantification of health risks and spatial distribution of heavy metals in groundwater of Lower Himalayas, India. *International Journal of Environmental Science and Technology*. **19**, 3201, **2022**.
- WARDHANI E., NOTODAMOJO S., ROOSMINI D. Assessment of Heavy Metal Contamination in Saguling Reservoir Water West Java Province Indonesia. *Web of Conferences*. **73**, 06009, **2018**.
- WANG Y., HAO C., HE K., PENG Y., SUN X., LIU M. Spatial distribution, formation mechanism, and health risk assessment of high-fluoride groundwater at the largest antimony mine in Hunan Province, China. *Water Supply*. **23** (8), 3402, **2023**.
- HAO R., YIN W., JIA H.Y., XU J.F., LI N.X., CHEN Q.Z., ZHONG Z.M., WANG J., SHI Z.H. Dynamics of dissolved heavy metals in reservoir bays under different hydrological regulation. *Journal of Hydrology*. **595**, 126042, **2021**.
- BI Y., WANG H., XIA B., JIANG C., WU W., LI Z., LI S., SU H., BAI Z., XU S., ZHUANG X. Pollution Characterization and Comprehensive Water Quality Assessment of Rainsource River: A Case Study of the Longgang River in Shenzhen. *Environmental Science*. **43** (2), 782, **2022** [In Chinese].
- ZHANG Q., FENG M., HAO X. Application of Nemerow Index Method and Integrated Water Quality Index Method in Water Quality Assessment of Zhangze Reservoir. *Earth and Environmental Science*. **128**, 012160, **2018**.
- SAHOO M.M., SWAIN J.B. Modified heavy metal Pollution index (m-HPI) for surface water Quality in river basins, India. *Environmental Science and Pollution Research*. **27**, 15350, **2020**.
- SHENG Y., WANG Z., FENG X. Potential ecological risk and zoning control strategies for heavy metals in soils surrounding core water sources: A case study from Danjiangkou Reservoir, China. *Ecotoxicology and Environmental Safety*. **252**, 114610, **2023**.
- MISHRA L.R., SINGH P.K., KOTHARI M., BHAKAR S.R., YADAV K.K., MEENA S.S. Appraisal of irrigation land suitability of semi-arid climatic zone of Rajasthan, India. *Water Supply*. **23** (11), 4800, **2023**.
- VERMA S.K., CHAURASIA S. A hydrogeochemical study of groundwater of district Ballia, U.P., India. *Water Supply*. **23** (8), 3221, **2023**.
- LI X., BING J., ZHANG J., GUO L., DENG Z., WANG D., LIU L. Ecological risk assessment and sources identification of heavy metals in surface sediments of a river-reservoir system. *Science of the Total Environment*. **842**, 156683, **2022**.
- YU H., GUI H., JIANG Y., LI Z., WANG M., FANG H., WANG C., CHEN C., QIU H., ZHANG Y. Characteristics of Dissolved Organic Matter Content in Urban Rivers under Different Environmental Impact Zones: A case study of China's Tuo River. *Polish Journal of Environmental Studies*. **29** (5), 3891, **2020**.
- XU J., GUI H., CHEN J., LI C., LI Y., ZHAO C., GUO Y. Hydrogeochemical Characteristics and Formation Mechanisms of the Geothermal Water in the Qingdong

- Coal Mine, Northern Anhui Province, China. *Mine Water and the Environment*. **4**, 1015, **2022**.
20. QIU H., GUI H., XU H., CUI L., YU H. Occurrence, controlling factors and noncarcinogenic risk assessment based on Monte Carlo simulation of fluoride in mid-layer groundwater of Huaibei mining area, North China. *Science of the Total Environment*. **856**, 159112, **2023**.
 21. MA J., CHEN S., FENG B., DING D. Hydrochemical characteristics and groundwater quality appraisal for irrigation uses in Lan-gan region, Northern Anhui Province, East China. *Water Supply*. **22** (3), 2644, **2022**.
 22. CHHEANG L., THONGKON N., SRIWIRIYARAT T., THANASUPSIN S.P. Heavy Metal Contamination and Human Health Implications in the Chan Thnal Reservoir, Cambodia. *Sustainability*. **13** (24), 13538, **2021**.
 23. JALEES M.I., IRFAN I., ALI A., BATOOL M. Spatial distribution and water quality index of Lahore Canal, Pakistan. *Water Practice and Technology*. **18** (2), 1791, **2023**.
 24. CHENG M., PENG W., ZHANG X., LIN M., YU H. Heavy Metal Content and Water Quality Assessment of Township Water Plants in Yongqiao District, Suzhou City. *Environmental Monitoring and Forewarning*. **16** (2), 94, **2024** [In Chinese].
 25. GAO Q., LI Y., CHENG Q., YU M., HU B., WANG Z., YU Z. Analysis and assessment of the nutrients, biochemical indexes and heavy metals in the Three Gorges Reservoir, China, from 2008 to 2013. *Water Research*. **92**, 262, **2016**.
 26. WANG W. Characteristics of Trace Elements and Health Risk Assessment in Surface Water on the Loess Plateau. Northwest A&F University, China, Master's thesis. **2022** [In Chinese].
 27. LUO M., ZHANG Y., LI H., HU W., XIAO K., YU S., ZHENG C., WANG X. Pollution assessment and sources of dissolved heavy metals in coastal water of a highly urbanized coastal area: The role of groundwater discharge. *Science of the Total Environment*. **807**, 151070, **2021**.
 28. ZHANG Y., TANG Y., CHEN J., SUN X., WU J., LIU B. Pollution Characteristics and Health Risk of Heavy Metals in Surface Water of Pingshui River. *Nonferrous Metals*. **7**, 116, **2021** [In Chinese].
 29. FOLARIN G.M., BADMUS B.S., AKINYEMI O.D., IDOWU O.A., OKE A.O., OMODELE T., BADMUS G.O. Assessment, classification and mapping of groundwater quality for agriculture in selected farm settlements of southwestern Nigeria. *Water Supply*. **23** (9), 3959, **2023**.
 30. ZHAO W. Study on pollution and risk assessment of heavy metals surrounding electroplating plants in Dongguan. China University of Geosciences, China, Doctor's thesis. **2016** [In Chinese].
 31. LIN S.Y. Bringing resource management back into the environmental governance agenda: eco-state restructuring in China. *Environment Development and Sustainability*. **23** (8), 12272, **2021**.
 32. ICS. Environmental quality standards for surface water, State Environmental Protection Administration of China, Beijing, China. **2002** [In Chinese].
 33. WANG Z., SU Q., WANG S., GAO Z., LIU J. Spatial distribution and health risk assessment of dissolved heavy metals in groundwater of eastern China coastal zone. *Environmental Pollution*. **290**, 118016, **2021**.
 34. LI Z., YAN X., WANG G., GUI H., YU H., XU X. Geochemical Baseline Establishment, Pollution Assessment, and Source Apportionment of Heavy Metals in Estuary Sediments of Northwestern Taihu Lake, China. *Polish Journal of Environmental Studies*. **32** (5), 4653, **2023**.
 35. REDDY Y.S., SUNITHA V. Assessment of Heavy metal pollution and its health implications in groundwater for drinking purpose around inactive mines, SW region of Cuddapah Basin, South India. *Total Environment Research Themes*. **8**, 100069, **2023**.
 36. ZOU X., LI Y., WANG L., AHMED M.K., CHEN K., WU J., XU Y., LIN Y., XIAO X., CHEN B., WU K., LIU J., LIN F. Distribution and assessment of heavy metals in suspended particles in the Sundarban mangrove river, Bangladesh. *Marine Pollution Bulletin*. **181**, 113856, **2022**.
 37. LI L., WU J., LU J., LI K., ZHANG X., MIN X., GAO C., XU J. Water quality evaluation and ecological-health risk assessment on trace elements in surface water of the northeastern Qinghai-Tibet Plateau. *Ecotoxicology and Environmental Safety*. **241**, 113775, **2022**.
 38. ZHAO L., GONG D., ZHAO W., LIN L., YANG W., GUO W., TANG X., LI Q. Spatial-temporal distribution characteristics and health risk assessment of heavy metals in surface water of the Three Gorges Reservoir, China. *Science of the Total Environment*. **704**, 134883, **2020**.
 39. YANG X. The Pollution Characteristic and Risk Assessment of Heavy Metals in the WeiHe River of Shaanxi Region. Xi'an University of Architecture and Technology, China, Doctor's thesis. **2015** [In Chinese].
 40. WANG Y., LI B., LI J., WANG Z., XIAN F., FAN Y. Spatial distribution characteristics and pollution levels of heavy metals in surface water and sediments of the Heihe cascade reservoir system, China. *Water Supply*. **22** (11), 8353, **2022**.