

Original Research

Isolation and Identification of Oil Degrading Bacteria from Oil Tanker Accident Locations in Jordan

Mohammad M. Aladwan*, Basem F. Dababneh, Husni S. Farah, Mai A.H. Abusalah

Al-Ahliyya Amman University, Amman 19111, Jordan

Received: 24 April 2024

Accepted: 9 December 2024

Abstract

Oil-degrading bacteria were found in soil samples polluted with oil in the oil tanker accident location and identified using biochemical tests and morphological analysis; fourteen bacterial isolates were identified. Bacterial isolates were identified at the molecular level using a universal primer, 16S rDNA gene. For soil samples, gas chromatography was used to evaluate total petroleum hydrocarbons. Heavy metal contamination (Cu, Cd, Mn, Zn, and Pb) was examined in all soil samples. CFU/g of the bacterial growth count ranged from 2.49×10^5 to 1.87×10^7 . The following bacterial genera were identified: *Staphylococcus*, *Pseudomonas*, *Rhodococcus*, *Serratia*, *Bacillus*, *Acetobacter*, *Micrococcus*, and *Lactobacillus*. *Lactobacillus casei*, *Micrococcus luteus*, and *Pseudomonas fluorescens* showed high growth rates on different types of hydrocarbons, such as toluene, naphthalene, and hexane. The highest Fe, Cd, Pb, Mn, and Cu concentrations were found in soil sample M2B. However, M1A has the highest Zn, Cu, and Fe concentrations. However, M1C has the lowest heavy metal concentrations. M2B, however, has the lowest Zn content. *Pseudomonas* species, which exhibit the greatest oil-degrading ability among all the isolates, may be utilized for bioremediation, as confirmed by this study. At one accident site, a high concentration of Cu showed high toxicity and low ability for hydrocarbon degradation in all bacterial isolates. Oil-contaminated locations might be cleaned up both in-situ and ex-situ using the species found in this study.

Keywords: bioremediation, *Pseudomonas*, oil degrading bacteria, 16S rDNA gene

Introduction

Crude oil is the most prevalent organic pollutant in all ecosystems due to spills during production, storage, transportation, and tank failures. It is classified as hazardous waste because of its cytotoxic, mutagenic,

and carcinogenic effects on living organisms. These pollutants, present in contaminated desert soils, can impact both human health and the ecosystem by seeping into nearby subsoil and groundwater [1]. Since many hydrocarbons are water-insoluble, they remain separated in the non-aqueous phase of the liquid [2]. The contamination of hydrocarbons has major negative impacts on ecosystems, such as decreased plant growth and seed germination. Hydrocarbons can coat plant

*e-mail: m.aladwan@ammanu.edu.jo

roots, reducing the absorption of water and nutrients. Moreover, these molecules can infiltrate plant tissues, causing damage to cell membranes. This damage can lead to the leakage of cell contents and blockage of intercellular gaps, which lowers the rates of respiration and metabolite transport [3].

According to the analysis of desert soil samples, the amount of lead (total or bioavailable) in soils contaminated with crude oil was three times higher than in uncontaminated soils [4, 5]. A high concentration of copper can decrease bacterial growth and viability, thereby inhibiting the ability of bacterial isolates to degrade hydrocarbons. Overall, high concentrations of heavy metals can increase the toxicity of the culture and decrease bacterial growth [6].

Desert soils with low organic carbon content, sandy texture, and reduced degradation rates generally exhibit poor microbial diversity and proliferation. Higher C ratios and lower internal surface areas are associated with lower abundances of degraders in the sandy component of polluted soils. Traditionally, the fertility of desert soils has been assessed using various physical and chemical characteristics, including aggregate stability, water-holding capacity, soil erosion, carbon fractions, and nutrient content. However, biological parameters such as microbial biomass and basal respiration, along with biochemical characteristics like hydrolases involved in the C, N, and P cycles, are more sensitive indicators closely linked to the microorganisms present in desert soil, offering a more accurate representation of the true degradation state [7].

Currently, chemical and mechanical treatments are employed in decontamination processes; however, these methods are expensive and often inefficient. In response to these challenges, bioremediation has emerged as a promising approach for cleaning oil-contaminated environments [8]. Bioremediation employs living organisms, most often microorganisms, plants, or byproducts produced by living organisms, to degrade, detoxify, or sequester toxic chemicals present in natural waters and soils. Bioremediation can be adapted for use in treating soil, sediment, sludge, water, or even air. Treatments can be either *ex situ* or *in situ*. *Ex-situ* procedures involve removing contaminated materials from a polluted site for treatment in another location, while *in situ* approaches treat contaminants in place without transferring samples to a treatment facility [9].

According to previous studies, there are several methods for utilizing microorganisms, especially bacteria, to break down environmentally hazardous waste materials such as crude oil. The existence and activity of these microorganisms, which are greatly influenced by factors such as pH, temperature, moisture content, and the composition of the microbial community (including fungi and bacteria), determine the effectiveness of bioremediation [10].

Recently, molecular and biochemical methods have been employed to identify and characterize bacteria and their specific properties at contaminated sites.

These techniques serve as complementary tools to classical chemical and physiological analytical methods (such as measuring pollutant concentrations and speciation, redox potential, etc.) to monitor spatial and temporal changes in bacterial community composition and functioning during bioremediation processes. The 16S rRNA gene has become a crucial DNA/gene fragment for identifying bacterial species, supporting comparative and phylogenetic research and classification efforts [11].

Moreover, bacterial isolates can effectively degrade a significant quantity of organic pollutants associated with oil, such as toluene, n-alkanes, aliphatic compounds, monoaromatic compounds, and polycyclic aromatic hydrocarbons (PAHs). However, high-molecular-weight PAHs are challenging to degrade for most genera [12]. A promising new technique for detecting petroleum hydrocarbons in environmental samples is comprehensive gas chromatography (GC). In gas chromatography, compound peaks are arranged along the x-axis and y-axis according to their chemical class and carbon number, respectively. This results in distinct chemical classes for petroleum, including cycloalkanes, alkanes, and one-, two-, and multi-ring aromatics. Additional groupings within each class display homologous series. Gas chromatography has proven to be an effective method for identifying and quantifying specific compounds and classes of compounds for crude oil and refined petroleum products [13].

Bacterial isolates utilize organic contaminants as nutrient sources, energy, and carbon. *Acinetobacter*, *Bacillus*, *Comamonas*, *Alcaligenes*, *Pseudomonas*, and *Micrococcus* are some of the more effective bacterial genera for the initiation of bioremediation [14-19]. However, previous studies indicate a noticeably greater decrease in bacterial numbers and diversity in newly polluted soils compared to older ones, with *Pseudomonas* being more common than other detected genera [20]. Furthermore, a similar study was conducted to isolate, purify, and identify bacterial strains from the Yanbu coastal region in Saudi Arabia. These strains can consume and break down hazardous substances from unintentional spills, oil leaks, and contamination from petrochemical industrial waste. According to this study, *Pseudomonas* species, which demonstrated the highest oil-degrading ability among all isolates, could be effectively utilized for bioremediation [21].

The highway between Jordan and Iraq, used for transporting crude oil, has two locations contaminated by oil spills resulting from tanker accidents, potentially impacting local flora. Desert soil samples collected from these sites were dry. This investigation aimed to identify and characterize bacterial isolates from oil-polluted arid soil at these accident locations and evaluate their ability to degrade various hydrocarbons, including crude oil, toluene, naphthalene, and hexane. In a subsequent phase known as '*In situ* bioremediation', these bacterial isolates could be utilized to remediate crude oil spills and facilitate plant regrowth.

Materials and Methods

Soil Sampling

Five soil samples, weighing 250 g were taken from two oil-contaminated sites in the northeastern region of Jordan at depths of 5, 10, and 15 cm. The soil samples were ground, put in polyethylene bags, and kept at 4°C until needed. The samples were then sieved through a 2 mm pore size sieve.

Culture Methods

150 g of soil samples were suspended in 300 mm of minimal-salt medium (Stanier's media) in 500 mm Erlenmeyer flasks supplemented with 400 ppm of crude petroleum oil. The flasks were then shaken at 160 rpm at 37°C. As controls, samples devoid of crude oil contamination [22].

Enumeration of Bacteria

Using Stanier's medium as the diluent, a range of serial dilutions (10^{-5} to 10^{-18}) were utilized to create isolated colonies on tryptic soy agar (TSA) and nutrient agar (N.A). Different media plates were used to isolate and select each colony type [22].

Oil-Degrading Bacteria Isolation Using Different Hydrocarbon Sources

After being cultivated on Stanier's agar, a minimal salt medium, 400 ppm of hydrocarbon compounds (crude oil, toluene, naphthalene, and hexane) were added, and the bacterial colonies were incubated for ten days at 37°C.

Morphological Features and Biochemical Characterizations

Each bacterial isolate was characterized morphologically based on gram staining, size, color, and colony characteristics (form and elevation). Biochemical and physiological tests were used to further identify the bacterial isolates [23]. The isolated bacterial colonies were a variety of colors, including yellow, brown, creamy white, and even green. Their shapes were described as circular, filamentous, punctiform, and rhizoid.

Molecular Characterization

Extraction of Genomic DNA

The protocol outlined in the previous study was followed for bacterial DNA extraction. Following an 18-h incubation period at 37°C, 10 mm of nutrient broth were inoculated with each bacterial isolate. 15 min at 14,000 rpm was used for centrifuging 2 mm

of the broth culture. 30 µl 25% w/v SDS, 3 µl Proteinase K (20 mg/ml), and 567 µl TE buffer were used to suspend the pellet. It was then incubated at 65°C for 30 min. The mixture was incubated with 100 µl of 5 M NaCl and 80 µl of CTAB/NaCl for 10 min at 65°C. After adding equal amounts of phenol, chloroform, and isoamyl alcohol (25:24:1), the mixture was mixed and centrifuged for 5 min at 14,000 rpm. After a second centrifugation of the mixture for 5 min at 14,000 rpm, the solution's supernatant was moved to a fresh tube. After adding 1.4 mg/ml of RNase enzyme to the mixture and letting it sit at 37°C for 35 min, the mixture was cooled with ice, and an equal amount of ice-cold isopropanol was added to precipitate the DNA. After rehydration in 100 µl of TE buffer (pH 7.5), the DNA pellets underwent three rounds of washing with 70% ethanol [24].

Polymerase Chain Reaction (PCR)

Identification of Bacterial Isolates through Sequencing of 16S rDNA

The 16S rDNA target sequence was amplified using the method outlined by a previous study [25]. 25 µl of the master mix (10 mM Tris-HCl pH 8.3, 50 mM KCl, 1.5 mM $MgCl_2$, 200 µM deoxynucleoside triphosphates, and 2.5 U of Taq DNA polymerase) is included in the 50 µl PCR mixture. Additionally, there are 2 µl of the reverse and 2 µl of the forward 16S rDNA universal primer, 5 µl of DNA template, and 16 µl of dH_2O . Reaction mixtures were subjected to 35 cycles of PCR reaction (denaturation at 94°C for 45 sec, annealing at 55°C for 1 min, and extension at 72°C for 1 min) in a DNA thermal cycler (Xp cycler, USA, 2008) at 96°C heat shock for 1 min. Every reaction mixture was kept cold at 4°C. 5-AGAGTTTGATCCTGGCTCAG-3 using the forward primer sequence (FD1) and 5-GGTTACCTTGTTACGACTT-3 as the reverse primer sequence (RD1) [26]. Finally, with the previous study description, 16S rDNA samples were sent to Macrogen, Inc., Korea, for sequencing [26].

Gel Electrophoresis and Photography

Each amplification reaction's aliquot (5 µl) was examined on 1.5% w/v agarose gels. The gels were analyzed using BioDocAnalyze (Biometra, Germany) after being stained with ethidium bromide (0.5 µg/ml). Each gel contained a marker consisting of 1000 bp.

Petroleum Hydrocarbon Determination

The EPA method 3510 was followed to extract soil samples for GC analysis. A 40 mL VOA vial was filled with 10 g of soil. Subsequently, 10 mL of methylene chloride, 15 mL of surrogate working solution (phenylacetylene), and 5 g of anhydrous sodium sulfate were added to the VOA vial. The sealed vial was submerged in a Grant, Inc., Germany, sonic bath

for 5 min. After shaking the vial, it was submerged in the sonic bath for five more min. Gas chromatography (Calruse 500 autosampler, Perkin Elmer, USA) was used to quantify the hydrocarbon compounds in the extract. The apparatus included a flame ionization detector (FID) and a capillary column, Rtx-1 (30 m x 0.53 mm; 0.1 µm film thickness; Silica fused, Philadelphia, Pa., USA). At 290 and 320°C, respectively, the temperatures of the injection port and detector were maintained constant. After being maintained at 50°C for 5 min, the oven's temperature was raised to 320°C at a rate of 10°C per min. The air and hydrogen gas flow rates were set at 2 mm per min for the flame ionization detector. The carrier was helium gas. Based on the retention time of the sample, the starting temperature and progress rate were chosen [27].

Heavy Metal Determination

Using a diluted HCl acid extraction method, heavy metals, including lead, zinc, cadmium, copper, ferrous, manganese, and zinc, were extracted from soils [28]. A 100 mL plastic bottle was filled with 200 mg (dry weight) of soil samples, followed by adding 2 mL of HF, 4 mL of 25% HNO₃, and 4 mL of 25% HCl. The samples were incubated in a water bath at 70°C for 2 h after being shaken for 2 min. After adding 50 mL of boric acid, the sample was once again submerged in the water bath for 15 min. Using dH₂O, the volume of each sample filtrate was increased to 100 mL, and an atomic absorption spectrophotometer (Analytic Jena, Inc.).

Results and Discussion

Enumeration of Bacterial Colonies

Results presented in Table 1 show the number of bacterial colonies (CFU) on nutrient agar in each soil sample. A range of crude oil concentrations (400, 600, 800, 1000, 1500, and 2000 ppm) were used to enhance the biodegradation ability of the tested bacterial isolates over a period that ranged from 50 days to 300 days.

Results indicated that M2A generated the highest CFU at either the least (400 ppm) or the highest (2000 ppm) concentrations of crude oil, where the obtained CFU values were 1.66×10^7 and 1.87×10^{17} , respectively. In contrast, the M1A sample showed the least (2.49×10^5 and 1.25×10^{14}) CFU at a concentration of 400 ppm and 2000 ppm, respectively. This study compared the number of viable bacterial colonies in samples isolated from oil-contaminated (test samples ranging from 2.49×10^5 to 1.87×10^{17} CFU·g⁻¹ soil) and uncontaminated (negative control; 9.8×10^4 CFU·g⁻¹ soil) soil. The variations in colony counts observed in the test samples could be attributed to their origins in both ancient and recently polluted locations. The highest toxicity levels of crude oil pollutants and the most detrimental effects on microbial diversity were observed in the old, contaminated areas. [29]. Additionally, Table 1 illustrates how adding various crude oil concentrations (400-2000 ppm) to the bacterial cultures at each site increased bacterial growth in tandem with rising crude oil concentrations.

Growth of Oil-Degrading Bacteria on Minimally Salted Media with a Different Source of Hydrocarbons

Among the 14 bacterial isolates, 7 were capable of degrading specific n-alkanes, such as hexane; 8 could degrade compounds with a single benzene ring, like toluene; and 5 isolates were capable of degrading compounds with two benzene rings, such as naphthalene. *Pseudomonas fluorescens* exhibited the highest growth rate on all hydrocarbons, while *Micrococcus luteus* and *Lactobacillus casei* appeared with the highest growth rate on hexane (Fig. 1). Conversely, low levels of growth on naphthalene have been observed in other isolates, including *Rhodococcus rhodochrous*, *Bacillus subtilis*, *Pseudomonas putida*, and *Serratia liquefaciens*. Furthermore, only toluene could be used as the exclusive energy and carbon source by *Staphylococcus simulans*, *Rhodococcus rhodochrous*, *Bacillus subtilis*, *Staphylococcus epidermidis*, *Staphylococcus saprophyticus*, and *Serratia liquefaciens* (Fig. 1).

Table 1. Growth of bacterial isolates from soil samples at different concentrations of crude oil.

Crude oil (ppm) \ Soil samples	M1A	M1B	M1C	M2A	M2B
400	2.49×10^5	1.24×10^7	1.32×10^6	1.66×10^7	1.54×10^7
600	2.07×10^7	2.31×10^8	2.17×10^8	1.19×10^9	1.17×10^8
800	1.67×10^9	1.45×10^{11}	2.49×10^9	0.89×10^{11}	1.59×10^9
1000	2.08×10^{10}	2.41×10^{13}	1.79×10^{11}	2.32×10^{12}	2.10×10^{10}
1500	1.89×10^{12}	1.31×10^{14}	1.27×10^{13}	1.53×10^{15}	1.07×10^{13}
2000	1.25×10^{14}	1.99×10^{15}	1.45×10^{15}	1.87×10^{17}	2.00×10^{14}

M: name of the site, A: 0-5 cm depth of soil samples, B: 5-10 cm depth of soil samples, C: 10-15 cm depth of soil samples, D: 15-20 cm depth of soil samples. ppm: part(s) per million from crude oil.




Fig. 1. Growth rate of oil-degrading bacteria on minimal salt media supplemented with four kinds of hydrocarbons, A) Hexane, B) Crude oil, C) Naphthalene, D) Toluene. II- +1: very low number of scattered colonies, +2: large number of medium-sized colonies, +3: bacterial growth all over the plate. III- These results appeared after triplicate experiments with these isolates.

According to our findings, the identified bacterial isolates utilize the supplemented hydrocarbons solely as a source of carbon and energy [30].

Biochemical Characterization and Morphological Features of Oil-Degrading Bacteria

The isolated bacterial colonies exhibited a variety of colors, including yellow, brown, creamy white, and even green. Their shapes varied, described as circular, filamentous, punctiform, and rhizoid. In terms of Gram staining, the bacterial isolates included gram-negative bacilli, gram-positive bacilli, and gram-positive cocci (Table 2). Table 2 summarizes biochemical and

morphological test results identifying bacterial genera such as *Staphylococcus*, *Pseudomonas*, *Rhodococcus*, *Serratia*, *Bacillus*, *Acetobacter*, *Micrococcus*, and *Lactobacillus*. The medium's temperature and pH significantly influenced the bacterial isolates' oil degradation capacity. An optimal temperature of 37°C was identified, facilitating growth across all bacterial species, initiating enzymatic processes, and enhancing hydrocarbon degradation.

Based on their morphological and biochemical traits, the 14 bacterial isolates obtained from culture on a minimal medium enhanced with crude oil were identified (Table 2). The isolates represented a variety of bacterial species. These bacterial isolates

exhibited diverse colors, such as yellow, brown, and white, indicating the production of different pigments (Table 2). Recent studies have highlighted that the genus *Pseudomonas* predominates and demonstrates the highest oil-degrading ability among all isolates in crude oil-contaminated soils. [31, 32]. This study also confirmed that *Pseudomonas* was the predominant genus isolated from the two contaminated sites. Fig. 1 illustrates that *Pseudomonas fluorescens* exhibited the highest growth ability across various hydrocarbons, as noted in recent references [33]. The findings of this study suggest potential applications for bioremediation.

On the other hand, according to other research, certain bacterial species are associated with the following: *Lactobacillus* [34], *Rhodococcus* [35], *Serratia* [36], *Acetobacter* [37], *Micrococcus* [38], *Staphylococcus* [39], and *Bacillus* [40].

Molecular Identification of the Bacterial Isolates

All 14 bacterial isolates had their genomic DNA extracted, and the target sequence of the 16S rDNA was amplified using a DNA thermal cycler (Xp cycler, USA, 2008). The 16S rDNA sequences of all isolates were identified through sequencing (Macrogen, Inc., Korea), and bacterial species were classified (see Table 3). As previously described, the isolated bacterial strains were identified using 16S rDNA sequencing [41]. Table 3 presents the 16S rDNA sequences of the newly isolated species and strains deposited in the NCBI database. Fig. 2 illustrates the phylogenetic tree of oil-degrading bacteria based on their 16S rDNA sequences. PCR-amplified 16S rRNA sequences were used for molecular identification, a sensitive and specific method for detecting microorganisms [41]. Each of the 14 bacterial isolates was molecularly identified, with all isolates producing a PCR product of approximately 1500 bp using universal primer pairs specific to the 16S rDNA gene (Fig. 2). Refer to Table 4 for details on the molecular identification of each bacterial isolate using universal 16S rDNA primers, confirming their species identity.

Petroleum Hydrocarbon Determination

Total petroleum hydrocarbons (TPH) were determined by GC/FID analyses of n-alkanes (C4-C30), monoaromatic hydrocarbons, and polyaromatic hydrocarbons (PAHs). The five oil-contaminated soils showed TPH contents ranging from 6,825 to 107,590 ppm (Fig. 3). Specifically, the M1A soil sample exhibited the highest levels of TPH (107,590 ppm), along with o-xylene and naphthalene. In contrast, the M2B soil sample had the lowest TPH concentration (6,825 ppm). Gas chromatography was utilized to quantify TPH.

The type and concentrations of total petroleum hydrocarbons (ppm) in the examined soil samples were determined using gas chromatography. As depicted in Fig. 3, soil samples M1A and M1C exhibited the highest

Table 2. a) Morphological Features, Biochemical and Physiological characterization of oil degrading bacteria.

	No	Bacterial isolates		1	2	3	4	5	6	7
		Tests								
Morphological characterization	1	Gram Stain	+C	+C	+B	+C	-C	+B	+C	
	2	Spore stain	NA	NA	+	NA	NA	-	NA	
	3	Acid Fast stain	NA	NA	NA	NA	NA	-	NA	
	4	growth Media	TSA	N.A	N.A	N.A	TSA	N.A	TSA	
	5	Colony Color	yellow	yellow	White	Brown	C. White	Brown	yellow	
	6	Colony Size	Small	Large	Large	Medium	Punctiform	Small	Small	
	7	Colony Form	Circular	Irregular	Circular	Circular	Circular	Circular	Circular	
	8	Colony elevation	Convex	Raised	Flat	Raised	Convex	Flat	Flat	
	9	Colony margin	Entire	Entire	Undulated	Entire	Undulated	Undulated	Entire	

Biochemical and physiological characterization	1	Catalase test	+	+	—	+	+	—	+	+	+	+
	2	Oxidase test	+	+	—	+	+	—	—	—	—	+
	3	Nitrate reduction test	+	+	+	+	+	+	+	+	+	+
	4	Methyl red test	—	—	+	—	—	+	+	+	—	—
	5	Degradation of starch	—	+	—	+	+	—	—	—	—	—
	6	Degradation of urea	+	—	—	—	—	—	—	—	—	—
	7	Degradation of casein	+	+	+	+	+	+	—	—	+	+
	8	Degradation of Tween-20	—	+	—	—	—	—	—	—	+	—
	9	Degradation of Tween-80	—	—	—	—	—	—	—	—	+	+
	10	Degradation of gelatin	+	—	+	—	—	+	—	—	+	+
	11	Gas and acid production from D-lactose	+G	—	—	—	—	—	+	—	—	+
	12	Gas and acid production from D-galactose	+G	—	—	—	—	—	+	—	—	+
	13	Gas and acid production from D-sucrose	—	+	+	+	+	+	+	+	+	+G
	14	Gas and acid production from D-inositol	+	—	+	—	—	+	+	—	—	+
	15	Gas and acid production from D-maltose	+	+	+	+	+	+	+	—	—	+
	16	Gas and acid production from D-fructose	+	—	+	—	—	+	+	—	—	+
	17	Utilization of citrate and propionate	+	+	—	+	+	—	+	—	+	+
	18	Blood hemolytic	γ	γ	α	γ	γ	α	γ	γ	α	α
	19	Triple sugar iron test (TSI)	K/A H	NC/NC	K/A	A/A	A/A	K/A	A/A	A/A	A/A	K/A
	20	Growth temperature 4°C	—	—	—	—	—	—	—	—	—	—
	21	Growth temperature 37°C	+	+	+	+	+	+	+	+	+	+
	22	Growth temperature 50°C	—	—	—	—	—	—	—	—	—	—
	23	Growth in the presence of NaCl 3%	+	—	+	—	—	+	+	+	+	+
	24	Growth in the presence of NaCl 10%	+	—	+	—	—	+	+	—	—	—
	25	Growth in the presence of NaCl 15%	—	—	—	—	—	—	—	—	—	—
	26	Eosin methylene blue (EMB)	+	—	+	—	—	+	+	+	—	+
	27	Macconkey agar	+	—	+	—	—	+	+	+	+	+
	28	Mannitol salt agar	—	—	ND	—	+	ND	+	ND	—	—
		Bacterial Species	<i>Staphylococcus simulans</i>	<i>Rhodococcus rhodochrous</i>	<i>Bacillus subtilis</i>	<i>Staphylococcus saprophyticus</i>	<i>Staphylococcus epidermidis</i>	<i>Micrococcus roseus</i>	<i>Lactobacillus casei</i>			

C: Cocci, B: Bacilli, F: Fluorescent, NA: Not Available, N.A.: Nutrient Agar, TSA: Tryptic Soya Agar, G: Gas.

Table 2. b) Morphological Features, Biochemical and Physiological characterization of oil degrading bacteria.

No	Tests	Bacterial isolates																	
		1	2	3	4	5	6	7	8	9	10	11	12	13	14				
Morphological characterization	Gram Stain								+C	-B	-B	-B	+C	-B	-B				
	Spore stain								NA	NA	NA	NA	NA	NA	NA				
	Acid Fast stain								NA	NA	NA	NA	NA	NA	NA				
	growth Media								N.A	N.A	N.A	N.A	N.A	N.A	TSA				
	Colony Color								yellow	yellow	Green	yellow	Clearness	White	White				
	Colony Size								Large	Large	Large	Medium	Medium	Small	Medium				
	Colony Form								Irregular	Irregular	Irregular	Circular	Circular	Irregular	Circular				
	Colony elevation								Raised	Convex	Flat	Raised	Flat	Flat	Flat				
	Colony margin								Filamentous	Entire	Undulated	Irregular	Undulated	Undulated	Entire				
Biochemical and physiological characterization	Catalase test								+	+	+	+	+	-	+				
	Oxidase test								+	+	-	+	-	+	-				
	Nitrate reduction test								+	+	+	+	+	+	+				
	Methyl red test								-	+	-	-	-	+	-				
	Degradation of starch								-	-	-	+	-	+	-				
	Degradation of urea								-	+	-	-	-	-	-				
	Degradation of casein								-	+	-	+	-	+	+				
	Degradation of Tween-20								-	-	+	-	-	-	-				
	Degradation of Tween-80								-	-	+	-	-	-	-				
	Degradation of gelatin								-	+	-	-	-	+	+				
	Gas and acid production from D-lactose								+	+	+	-	+	-	-				
	Gas and acid production from D-galactose								-	+G	-	-	+G	-	-				
	Gas and acid production from D-sucrose								-	+	-	-	+G	+	+				
	Gas and acid production from D-inositol								-	+	-	-	+	+	+				
	Gas and acid production from D-maltose								+	+	-	-	+G	+	+G				
	Gas and acid production from D-fructose								+	+	-	-	+G	+	+				
	Utilization of citrate and propionate								+	+	+	+	+	+	+				
	Blood hemolytic								γ	α	α	γ	γ	β	β				
	Triple sugar iron test (TSI)								K/A	K/A H	K/K	K/NC	K/A G	K/A	A/A				
	Growth temperature 4°C								-	-	-	-	-	-	-				

Table 3. Bacterial species identification based on 16s rDNA sequencing data.

Isolates No.	Total Length	Gene bank accession No.	Similarity %	Identification result
1	1200 bp	MF678966.1	100	<i>Staphylococcus simulans</i>
2	1059 bp	LT906450.1	100	<i>Rhodococcus rhodochrous</i>
3	1190 bp	MT554031.1	100	<i>Bacillus subtilis</i>
4	770 bp	CP054440.1	98	<i>Staphylococcus saprophyticus</i>
5	1128 bp	MT605363.1	99	<i>Staphylococcus epidermidis</i>
6	1080 bp	MH793526.1	97	<i>Micrococcus roseus</i>
7	1190 bp	MK774613.1	98	<i>Lactobacillus casei</i>
8	1115 bp	MN905159.1	98	<i>Micrococcus luteus</i>
9	1165 bp	MT300520.1	97	<i>Pseudomonas fluorescens</i>
10	1317 bp	KC662508.1	97	<i>Acetobacter aceti</i>
11	1048 bp	KP072763.1	95	<i>Pseudomonas putida</i>
12	1050 bp	CP054434.1	98	<i>Staphylococcus saprophyticus</i>
13	910 bp	NR_042062.1	98	<i>Serratia liquefaciens</i>
14	1337 bp	CP014017.2	98	<i>Serratia liquefaciens</i>




Fig. 3. I-Total petroleum Hydrocarbons in 5 soil samples and the control sample. II- M: soil contaminated by oil, C: soil uncontaminated by crude oil, A: 0-5 cm of depth, B: 5-10 cm of depth, C: 10-15cm of depth and D: 15-20 cm of depth. III- www.Restek.com (Retention time of hydrocarbons in the same condition from this website).

Table 4. List of the oil degrading bacterial isolates extracted from different soil samples.

Number of isolates	Bacterial isolates codes	Species
1	M1A(1)	<i>Staphylococcus simulans</i>
2	M1A(2)	<i>Rhodococcus rhodochrous</i>
3	M1A(3)	<i>Bacillus subtilis</i>
4	M1A(4)	<i>Staphylococcus saprophyticus</i>
5	M1A(5)	<i>Staphylococcus epidermidis</i>
6	M1A(6)	<i>Micrococcus roseus</i>

7	M1A(7)	<i>Lactobacillus casei</i>
8	M1A(8)	<i>Micrococcus luteus</i>
9	M1B(9)	<i>Pseudomonas fluorescens</i>
10	M1B(10)	<i>Acetobacter aceti</i>
11	M2B(1)	<i>Pseudomonas putida</i>
12	M2B(2)	<i>Staphylococcus saprophyticus</i>
13	M2B(3)	<i>Serratia liquefaciens</i>
14	M2B(4)	<i>Serratia liquefaciens</i>

M: name of the site, A: 0-5 cm depth of soil samples, B: 5-10 cm depth of soil samples, C: 10-15 cm depth of soil samples, D: 15-20 cm depth of soil samples.




Fig. 4. I- Showed the concentration of many heavy metals (Zn, Cu, Pb, Mn, Cd, Fe) for different sites (M1A, M1B, M1C, M2A, M2B, C1, C2). II- M: soil contaminated by oil, C: soil uncontaminated by crude oil, A: 0-5 cm of depth, B: 5-10 cm of depth, C: 10-15 cm of depth and D: 15-20 cm of depth. III- Reference soils, that were taken from soil uncontaminated by crude oil.

of Cu was detected in the M1A sample, leading to high toxicity and reduced hydrocarbon degradation ability in all bacterial isolates. Consequently, the total petroleum hydrocarbon (TPH) concentration remains high with long-term contamination.

Heavy metal contamination has adversely affected the viability of crude oil biodegradation in oil-contaminated soil. Certain heavy metals can be detrimental to microorganisms, inhibiting their growth. Copper toxicity, for instance, primarily affects microorganisms through interactions with nucleic acids, altering enzyme active sites, oxidizing membrane constituents, and potentially generating harmful hydroxyl free radicals [43].

Flame atomic absorption spectroscopy was employed to analyze heavy metals, which influenced bacterial growth and subsequently affected crude oil degradation. The highest Cu, Fe, Mn, and Pb concentrations were observed in M2 soil samples. Consequently, these samples exhibited the lowest isolate counts, likely due to growth inhibition. Cadmium, lead, cobalt, copper, and zinc were among the metals detected most frequently. These metals exhibited increased toxicity in the following order: lead<zinc<copper<cadmium [44].

Conclusions

This study aimed to identify oil-degrading bacteria from contaminated soil near the Jordanian–Iraqi border. All isolates have been designated in the NCBI database and were identified through molecular detection and biochemical analysis of bacterial species. The optimum temperature for the isolates' oil degradation activity was found to be 37°C. *Pseudomonas fluorescens* exhibited the highest growth across all selective hydrocarbons used in this study. High copper concentrations were found at the accident site, resulting in high toxicity and reduced hydrocarbon degradation ability among all bacterial isolates. As a result, total petroleum hydrocarbon (TPH) concentrations remain high despite the presence of aged crude oil spills at the site. The species identified in this study could be utilized for both in situ and ex-situ cleanup of contaminated sites, such as oil-contaminated desert soil. Isolated bacteria have demonstrated the ability to degrade oil, making them suitable for remediation efforts to clean up oil-contaminated soil.

Acknowledgments

We would like to express our deepest gratitude and appreciation to our family and to those who have supported us in finishing this work.

Conflict of Interest

The authors declare no conflict of interest.

References

- MÜNZEL T., HAHAD O., DAIBER A., LANDRIGAN P.J. Soil and Water Pollution and Human Health: What Should Cardiologists Worry about? *Cardiovascular Research*. **119** (2), 440, **2023**.
- CUBITTO M.A., MORAN A.C., COMMENDATORE M., CHIARELLO M.N., BALDINI M.D., SENERIZ F. Effects of *Bacillus subtilis* O9 biosurfactant on the bioremediation of crude oil-polluted soils. *Biodegradation*. **15** (5), 281, **2004**.
- HAIDER F.U., EJAZ M., CHEEMA S.A., KHAN M.I., ZHAO B., LIQUN C., SALIM M.A., NAVEED M., KHAN N., NÚÑEZ-DELGADO A., MUSTAFA A. Phytotoxicity of petroleum hydrocarbons: Sources, impacts and remediation strategies. *Environmental Research*. **197**, 111031, **2021**.
- KHUDUR S., SHAHSAVARI E., WEBSTER G.T., NUGEGODA D., BALL A.S. The impact of lead co-contamination on ecotoxicity and the bacterial community during the bioremediation of total petroleum hydrocarbon-contaminated soils. *Environmental Pollution*. **253**, 939, **2019**.
- CETIN M., ALJAMA A.M.O., ALRABITI O.B.M., ADIGUZEL F., SEVIK H., CETIN I.Z. Determination and Mapping of Regional Change of Pb and Cr Pollution in Ankara City Center. *Water, Air & Soil Pollution*. **233**, 163, **2022**.
- MUHAMAD F.H., MANOGARAN M., YASID N.A. Inhibitory Effect of Copper on the Growth Rate of *Serratia marcescens* strain DRY6 on SDS. *Bioremediation Science and Technology Research*. **10** (2), 23, **2022**.
- HU Y., WANG Z., ZHANG Z., SONG N., ZHOU H., LI Y., WANG Y., LI C., HALE L. Alteration of desert soil microbial community structure in response to agricultural reclamation and abandonment. *Catena*. **207**, 105678, **2021**.
- RAJASULOCHANA P., PREETHY V. Comparison on Efficiency of Various Techniques in Treatment of Waste and Sewage Water. *Resource-Efficient Technologies*. **2** (4), 175, **2016**.
- BALA S., GARG D., THIRUMALESH B.V., SHARMA M., SRIDHAR K., INBARAJ B.S., TRIPATHI M. Recent Strategies for Bioremediation of Emerging Pollutants: A Review for a Green and Sustainable Environment. *Toxics*. **10**, 484, **2022**.
- EMMANUEL E., USMAN A.A., DARA OBONG U., QUEEN O. Bioremediation Potentials of Some Indigenous Microorganisms Isolated from Auto Mechanic Workshops on Irrigation Water used in Lokoja Kogi State of Nigeria. *Journal of Bacteriology & Mycology*. **4** (4), 121, **2017**.
- ALADWAN M.M., DABABNEH B.F., FARAH H.S., ABUSALAH M.A.H. Identification of Oil Degrading Bacteria from Oil-Contaminated Soil in the Northeastern Part of Jordan. *Journal of Ecological Engineering*. **25** (5), 306, **2024**.
- KHANAFER M., AL-AWADHI H., RADWAN S. Coliform Bacteria for Bioremediation of Waste Hydrocarbons. *BioMed Research International*. **2017**, 1, **2017**.
- NELSON R.K., KILE B.M., PLATA D.L., SYLVA S.P., XU L., REDDY C.M., GAINES R.B., FRYSSINGER G.S., REICHENBACH S.E. Tracking the weathering of an oil Spill with comprehensive two-dimensional gas chromatography. *Environmental Forensics*. **7** (1), 33, **2006**.

14. RAJU M.N., LEO R., HERMINIA S.S., MORÁN R.E., VENKATESWARLU K., LAURA S. Biodegradation of Diesel, Crude Oil and Spent Lubricating Oil by Soil Isolates of *Bacillus* spp. *The Bulletin of Environmental Contamination and Toxicology*. **98** (5), 698, **2017**.
15. PHULPOTO A.H., QAZI M.A., MANGI S., AHMED S., KANHAR N.A. Biodegradation of oil-based paint by *Bacillus* species monocultures isolated from the paint warehouses. *International Journal of Environmental Science and Technology*. **13** (1), 125, **2016**.
16. WU Y., MOHANTY A., CHIA W.S., CAO A. Influence of 3-Chloroaniline on the Biofilm Lifestyle of *Comamonas testosteroni* and Its Implications on Bioaugmentation. *Applied and Environmental Microbiology*. **82** (14), 4401, **2016**.
17. FADHIL G.F., AL-HADITHI H.T., AL-RAZZAQ E.A. Bioremediation of Polycyclic Aromatic Hydrocarbon by *Acinetobacter* Species Isolated from Ecological Source. *Journal of Environmental Biology*. **38** (5), 785, **2017**.
18. OJEWUMI E.E., OKENIYI J.O., IKOTUN J.O., OKENIYI E.T., EJEMEN V.A., POPOOLA A. Bioremediation: Data on *Pseudomonas aeruginosa* effects on the bioremediation of crude oil polluted soil. Data in brief. **19**, 101, **2018**.
19. KAWO A.H., BACHA H.Y. Crude Oil Degradation by *Bacillus* and *Micrococcus* Species Isolated from Soil Compost in Kano, Nigeria. *Bayero. Journal of Pure and Applied Sciences*. **9** (1), 108, **2016**.
20. SAADOUN I., MOHAMMAD M.J., HAMEED K.M., SHAWAQFAH M. Microbial population of crude oil spill polluted soils at the Jordan- Iraq desert (the Badia region). *The Brazilian Journal of Microbiology*. **39** (3), 453, **2008**.
21. EL HANAFY A., ANWAR Y., MOHAMED S.A., AL-GARNI M.S., SABIR J.S.M., ABUZINADAH O.A., AL-MEHDAR H., ALFAIDI A.W., AHMAD M.M. Isolation and identification of bacterial consortia responsible for degrading oil spills from the coastal area of Yanbu, Saudi Arabia. *Biotechnology & Biotechnological Equipment*. **30** (1), 69, **2016**.
22. AL-DEEB T.M., MALKAWI H.I. Isolation, molecular and biochemical characterization of oil degrading bacteria from contaminated soil at an oil refinery. *The Journal of Applied Research and Technology*. **14**, 1, **2009**.
23. ROHOMANIA T., SAHA M.L., HOSSAIN A., RAHMAN M.S. Morphological and Biochemical Characterization of Bacteria Isolated from Fresh and Salted Hilsa, *Tenualosa ilisha* (Hamilton, 1822). *Bangladesh Journal of Microbiology*. **32**, 7, **2016**.
24. CHACHATY E., SAULNIER P. Isolation chromosomal DNA from bacteria. In: Rapley, R. (eds) *The Nucleic Acid Protocols Handbook*. Springer Protocols Handbooks. Humana Press: Totowa, NJ, USA. pp. 29 -32, **2000**.
25. CHEN Y.L., LEE C.C., LIN Y.L., YIN K.M., HO C.L., LIU T. Obtaining long 16S rDNA sequences using multiple primers and its application on dioxin-containing samples. *BMC Bioinformatics*. **16**, 13, **2015**.
26. ADAMS D.D., BAWA M.D., ABDULLAHI U.Y., WUYEP P.A., ABBA D. Molecular Characterization and Determination of Bioremediation Potentials of Some Bacteria Isolated from Spent Oil Contaminated Soil Mechanic Workshops in Kaduna Metropolis. *World Applied Sciences Journal*. **34** (6), 750, **2016**.
27. BRZESZCZ J., STELIGA T., RYSZKA P., KASZYCKI P., KAPUSTA P. Bacteria degrading both n-alkanes and aromatic hydrocarbons are prevalent in soils. *Environmental science and pollution research international*. **31** (4), 5668, **2024**.
28. CHOI K.Y., KIM S.H., CHON H.T. Relationship between total concentration and dilute HCl extraction of heavy metals in sediments of harbors and coastal areas in Korea. *Environmental Geochemistry and Health*. **34**, 243, **2012**.
29. RAMDASS A.C., RAMPERSAD S.N. Diversity and Oil Degradation Potential of Culturable Microbes Isolated from Chronically Contaminated Soils in Trinidad. *Microorganisms*. **9** (6), 1167, **2021**.
30. KALANDER E., ABDULLAH M.M., AL-BAKRI J. The Impact of Different Types of Hydrocarbon Disturbance on the Resiliency of Native Desert Vegetation in a War-Affected Area: A Case Study from the State of Kuwait. *Plants (Basel)*. **10** (9), 1945, **2021**.
31. YOU X.H., ZHANG S., KIM H., CHIANG P.C., YUN W., ZHANG L., HE M. Comparison of Petroleum Hydrocarbons Degradation by *Klebsiella pneumoniae* and *Pseudomonas aeruginosa*. *Applied Science*. **8** (12), 2551, **2018**.
32. RAEID M.M.A., AL-SABAHI J., AL-MAQRASHI F., AL-HABSI A., AL-HINAI M. Characterization of hydrocarbon-degrading bacteria isolated from oil-contaminated sediments in the Sultanate of Oman and evaluation of bioaugmentation and biostimulation approaches in microcosm experiments. *International Biodeterioration & Biodegradation*. **89**, 58, **2014**.
33. PANDEY A.K. Isolation of Oil Degrading Bacteria from oil Contaminated Soil and Expression of oil degrading genes in non-oil degrading bacteria. *Journal of Drug Discovery and Therapeutics*. **1**, **2013**.
34. MUKJANG N., CHITOV T., MHUANTONG W., CHAMPREDA V., PATHOM-AREE W., SATTAYAWAT P., BOVONSOMBUT S. Bacterial Communities Associated with Crude Oil Bioremediation through Composting Approaches with Indigenous Bacterial Isolate. *Life (Basel)*. **12** (11), 1712, **2022**.
35. YU T., LIU X., AI J., WANG J., GUO Y., LIU X., HE X., DENG Z., JIANG Y. Microbial community succession during crude oil-degrading bacterial enrichment cultivation and construction of a degrading consortium. *Frontiers in Microbiology*. **13**, 1044448, **2022**.
36. BIDJA A.M.T., CHEN G., CHEN Z., ZHENG X., LI S., LI T., ZHONG W. Microbial diversity changes and enrichment of potential petroleum hydrocarbon degraders in crude oil-, diesel-, and gasoline-contaminated soil. *3 Biotech*. **10** (2), 42, **2020**.
37. MUKJANG N., CHITOV T., MHUANTONG W., CHAMPREDA V., PATHOM-AREE W., SATTAYAWAT P., BOVONSOMBUT S. Bacterial Communities Associated with Crude Oil Bioremediation through Composting Approaches with Indigenous Bacterial Isolate. *Life (Basel)*. **12** (11), 1712, **2022**.
38. BRZESZCZ J., STELIGA T., RYSZKA P., KASZYCKI P., KAPUSTA P. Bacteria degrading both n-alkanes and aromatic hydrocarbons are prevalent in soils. *Environmental science and pollution research international*. **31** (4), 5668, **2024**.
39. EDDOUAOUDA K., MNIF S., BADIS A., YOUNES S.B., CHERIF S., FERHAT S., MHIRI N., CHAMKHA M., SAYADI S. Characterization of a novel biosurfactant produced by *Staphylococcus* sp. strain 1E with potential application on hydrocarbon bioremediation. *Journal of Basic Microbiology*. **52** (4), 408, **2012**.
40. OBI L.U., ATAGANA H.J., ADELEKE R.A. Isolation and characterization of crude oil sludge degrading bacteria. *Springer Plus*. **5** (1), 1946, **2016**.

41. THIJS S., BEECK O.D., BECKERS B., TRUYENS S., STEVENS V., VAN HAMM J.D., WEYENS N., VANGRONSVELD J. Comparative Evaluation of Four Bacteria-Specific Primer Pairs for 16S rRNA Gene Surveys. *Frontiers in microbiology*. **8**, 1, **2017**.
42. POPOOLA B. M., OLANBIWONNINU A.A. Biotreatment of Crude Oil Contaminated Soil. *Microbiology Research Journal International*. **27** (4), 1, **2019**
43. MUHAMAD F.H., MANOGARAN M., YASID N.A. Inhibitory Effect of Copper on the Growth Rate of *Serratia marcescens* strain DRY6 on SDS. *Bioremediation Science and Technology Research*. **10** (2), 23, **2022**.
44. PIRI M., SEPEHR E., SAMADI A., FARHADI K.H., ALIZADEH M. Contaminated soil amendment by diatomite: chemical fractions of zinc, lead, copper and cadmium. *International Journal of Environmental Science and Technology*. **18**, 1191, **2021**.