

Review

Bioremediation towards Environmental Sustainability: Molecular Approaches to Tackle Inorganic and Organic Pollution

Magdalena Nikolić^{o*}, Vladimir Tomašević

School of Engineering Management, University Union Nikola Tesla,
Bulevar Vojvode Misica 43, 11000 Belgrade, Serbia

Received: 24 September 2024

Accepted: 21 February 2025

Abstract

Development in every aspect interconnects society, economy, and environment. Significant advances in technology and industry have led to various environmental problems affecting ecosystems, climate, and biodiversity. We must face the challenge of the adjustment of industry and the progress of society to a clean environment. The framework of this article is built around imperatives given by the European Council through important documents related to environmental protection, climate change, and clean energy. In the European Green Deal, Horizon Europe, and EU partnership for R&I, researchers are given the benefit of the doubt to halt and remove pollution. In line with the EU Circular Talks platform that includes bioremediation as a new–old approach to dealing with pollution sustainably, the study aims to present substantive examples of the capacity of some microorganisms to remove toxic substances from the environment. Therefore, the study focuses on the determinants in native and engineered organisms. The article aims to provide biochemical data valuable in cleaning particular pollutants in vivo/in vitro, showing scientific data available to researchers and stakeholders. This paper aims to clarify the potential of in situ biological techniques and their application to remove different inorganic or organic contaminants in the environment.

Keywords: pollution removal, bioremediation, environmental sustainability, genetic determinants

Introduction

Faced with climate change, energy, agriculture, tourism, and pollution issues, the EU is determined to address them in such a way as to provide its citizens with the highest environmental standards. Objectives and goals to be achieved within 2020-2050 are set

and prioritized by research programs, legislation, and funding [1]. Pollution is a global problem that affects human health, biodiversity, and the economy. It is a problem that is undoubtedly growing, creating a cascade of consequences that require a completely different approach with the same outcome: a clean, healthy, and sustainable environment. In recent years, many types of pollutants have been uncontrolled and released into the environment. Hydrocarbons, heavy metals, and pesticides are organic and inorganic contaminants with the highest toxic effects.

*e-mail: magdalena.nikolic@fim.rs

^o ORCID iD 0000-0002-2021-153X

Ecology can be seen as the study of nature's household viewed through abiotic and biotic perspectives, researched via the metabolism of living organisms, and understood as a weapon against domicile [2]. The reasoning behind the classical definition of ecology is an accurate understanding of humanity's demands on the biosphere. Thus, frequently exposed to a mixture of xenobiotics such as heavy metals, pesticides, and other toxic substances, bioaccumulation and biosorption can be only evolutionary steps [3]. In line with what was previously said, natural techniques consider the use of biological remediation methods for the contaminated environment as essential. The biological methods available for soil remediation are phyto- and bioremediation, considered the safest and, thus, ecologically acceptable [4, 5].

Phytoremediation is a natural process in which plants uptake and transform a variety of pollutants. This method also involves rhizosphere stabilization, boosting microbial growth in plants' root zone [6-8]. Although applicable in water, phycoremediation is categorized as phytoremediation and bioremediation [9], in which algae and microalgae naturally remove nitrogen, phosphorus, sulfur, and minerals from water. A simple explanation would be that algae feed on these pollutants in water [10]. Also, using algae and microalgae to remove phenolic compounds from household wastewater was useful in treating communal wastewater [11]. Sing et al. (2021) showed that filamentous algae, a genus *Spirogyra*, can be used for Cr(VI) removal and decolorization of methylene blue dye in wastewater [12]. Another method called mycoremediation, or fungal bioremediation, includes fungal microbes' capability to effectively adapt and clean the environment [13]. The bioremediation method includes physical, chemical, and biological techniques. These include mechanisms categorized into in-situ and ex-situ methods in the polluted place or outside it. Certain organisms have the genetic potential to remove, degrade, metabolize, or immobilize a wide range of contaminants from soil and water. The biochemical ability to tolerate high concentrations of toxic organic and inorganic pollutants is related to specific genes involved in detoxification by enzymes, transport protein families, and efflux pump proteins due to adaptation to specific environmental conditions [7, 14]. Bioremediation as a technique presents a series of reactions in which the main role is played by enzymes, synthesized by the actors of bioremediation themselves, microorganisms. This process involves the participation of archaea, bacteria, algae, and fungi whose application showed successful, inexpensive, and safe methods [15].

Sustainable Development, Ecology/Environment, and Bioremediation Are Closely Intertwined Fields

Sustainable development aims to balance human needs with using natural resources (soil, air, water). Considering economic growth and social rights,

17 sustainability development goals are brought by the UN Agenda 2030 [16]. All sustainable goals are viewed through different sets of indicators; they include life quality and ecological/environmental sustainability. Sustainable development is supported by three pillars (environment, economics, and society), and goals are set for each of them [17]. Particularly, environmental sustainability is related to the capacity of the planet to give and stay preserved. Therefore, development must include remediation of polluted soil and water and/or proportionately using the resources of the planet Earth [18]. Therefore, the European Union set goals and objectives within the European Green Deal, focusing on zero pollution and research and innovation (R&I) action [18, 19]. One segment of this document, imperative for achieving environmental and ecological goals, presents the missions on soil, ocean, sea, and water remediation [20]. The EU supports science as a foundation for research and innovation, and this topic is detailed and elaborated under Cluster 6, which tackles the loss of biodiversity, healthy soil and water shrinkage, pollution problems, and environmental deterioration [21]. Specifically, the Horizon Europe program through Cluster 6 boosts sound solutions by implementing a circular economy and bioremediation to preserve, prevent, and remediate the environment. R&I in this cluster aims to prioritize the selected headline indicators built on the European Green Deal and the 8th Environment Action Program (2022-2030) [21].

Within Cluster 6, some projects prioritize molecular aspects of bioremediation. The Biosysmo project (BIOremediation systems exploiting SYnergieS for improved removal of mixed pollutants) brings novelty to bioremediation applications. Bioinformatics methods and genetic material manipulation techniques were applied to create a synergetic biosystem. There was also a special focus on secondary toxic substances and their neutralization. There are several publications funded by this project [22]. Dvořák et al. (2023) analyzed the potential of *Pseudomonas putida* mt-2 to biodegrade m-xylene (aromatics) under different water conditions, such as water potential. The research focused on the expression of the biodegradative xyl genes that encode for the XylR A domain. When modifying water potential, matrix stress tolerance was shaped by the expression of xyl genes (the TOL plasmid pWW0). It modified the A domain of XylR (the structure of the binding pocket and entry tunnel). Water-dependent transcription changes led to low fitness of genotypes but adapted strains to new conditions [23]. Avendaño et al. (2023) revealed the *Pseudomonas putida* KT2440 catabolic pathway in the selenite to selenium nanoparticles transformation. The transformation involves *sucA*, *D2HGDH*, and *PP_3148* genes that encode enzymes that bond the 2-ketoglutarate and enzymes of oxidative stress GSH by glutathione. An important role is the sulfite reductase encoded by *cysG*, *sqr*, *pdo2*, *sqrR*, *ssuEF*, and *sfnCE* sulfur metabolism genes. Certain genes revealed in *P. putida* (WT strain), such as *sqrR*, *pdo2*, and *sqr*, haven't

been found in other bacterial strains that metabolize selenium. Furthermore, these genes are of great interest because switching them off increases selenium nanoparticle production [24]. The GREENER project aims to provide natural, low-cost, and safe solutions by introducing bioremediation and bio-electrochemical technologies. To successfully remove metals and metalloids, PAHs, TPHs, antibiotics, pesticides, etc., a combination of bioremediation methods and risk assessment will be applied. [25]. The results of this project are presented in several studies. Martínez-Cuesta et al. (2023) explored aliphatics and aromatics breakdown rates in polluted soils from Ireland using more than one bioremediation technique (Ecopiling). During the process, DSeq2 software was used for microorganism identification and appearance in the given remediation period. It was notable that at the beginning of the process, there were more TPH degraders (*Alcaligenaceae* or *Pseudomonadaceae*, *Gammaproteobacteria*); afterward, bacterial populations inhabit less polluted soils such as *Actinobacteria*, *Saccharimonadia*, *Acidobacteriota*, and some *Actinobacteriota* or *Firmicutes* were noticed. At the end of bioremediation, about 90% of TPHs in polluted soil were degraded [26]. Similarly, Verdel et al. (2021) explored the degradation rate of organic additives used in paper mills. Consortia of native and engineered microorganisms were used. The four bacteria in the process of bioaugmentation, *Xanthomonadales bacterium* and *Sphingomonas* sp., and *Cellulosimicrobium* and *Aeromonas* sp., showed the biggest potential [27]. Garrido-Sanz et al. (2020) showed that within the *Rhodococcus* genus, the majority can break down aromatic and aliphatic compounds; however, only some strains, like *Rhodococcus* sp. WAY2 can degrade short-chain n-alkanes [28]. Further investigation of *Rhodococcus* sp. WAY2 showed independent clusters of bph, etb, and nah genes involved in the biodegradation pathway of aromatic compounds. The hydrogenase enzyme system coded by these clusters enables *Rhodococcus* sp. WAY2 to metabolize biphenyl, naphthalene, and xylene. The ability to use short and long chains of n-alkanes is due to the presence of chromosomal alkB and ladA genes [29]. Another project funded by the EU, MIBIREM, brings solid solutions regarding bioremediation. This project aims to create a toolbox to gather molecular methods and bioinformatics tools. Microorganisms will then be applied in cyanides, hexachlorocyclohexane (HCH), and petroleum hydrocarbons (PHC) polluted cities within Europe. The expected outcome is the creation and application of the MIBIREM TOOLBOX, which includes identified bioremediation and their genetic information, as well as IT modeling and forecasting tools, resources, and protocols [30]. The objective mentioned in previous documents is achieving the SDGs regarding the environmental pillar "planet" by using biocenosis (decomposers) in pollution removal.

The introduction of organic and inorganic pollutants into an ecosystem causes adverse impacts on health,

biota, and environmental change. The remediation process includes pollution elimination to preserve the ecosystem's health [31]. An ecological approach includes decomposers, as they can break down pollutants into harmless byproducts, therefore restoring ecosystems. Biological processes include bioremediation and biotechnology. Apart from biosensors and their wide use in bioremediation, genetic engineering/genetic modification or manipulation still keeps the top position by using living organisms as a tool in technology [32].

Bioremediation is considered economical, as a natural way of eradicating toxins and restoring polluted sites by choosing the right techniques. Following the principles of bioremediation, there are two main techniques, biostimulation and bioaugmentation, used to degrade harmful and toxic substances.

Bioaugmentation introduces specialized microorganisms to enhance pollutant degradation, and they act as biocontrol agents in contaminated soil, whereas biostimulation involves the transfer of certain microorganisms that act as biostimulants to indigenous bacterial populations [32-34]. The relationship between ecology and bioremediation lies in their shared goal of understanding and restoring environmental health. Ecological principles inform bioremediation strategies by providing insights into the complex interactions within ecosystems and the potential impacts of contaminants on organisms and their habitats. Biotic and abiotic factors that have anthropogenic factors influence every ecosystem and, thus, all processes within. Additionally, the speed and degree of breaking down pollutants depend on the physicochemical characteristics such as solubility, volatility, and bioavailability of pollutants, as well as abiotic factors such as pH, temperature, and moisture content, and biotic factors such as the microbial community and interactions with living and nonliving nature. Overall, integrating ecology and bioremediation is essential for developing effective, sustainable solutions to environmental pollution and ensuring the long-term health and resilience of ecosystems [35, 36]. Therefore, successful bioremediation depends on microbial ecology and metabolism. Thus, developments in molecular microbiology can optimize bioremediation processes in a targeted manner. Novel advances in biotechnology introduced their genetic background, namely enzymes and genes that enable the degradation or chemical change of harmful organic and inorganic pollutants. The promising technology of genetic engineering provides a myriad of possibilities when it comes to removing As, Cd, Cu, Fe, Hg, and Ni, as well as POPs such as PAHs and PCBs [37], by using genetically modified organisms (GMOs) [38, 39] / genetically engineered microorganisms (GEMs) in bioremediation [40].

Materials and Methods

The framework of this article is built around imperatives given by the European Council through

important documents related to environmental protection, climate change, and clean energy. Also, under this topic and with funds provided through the European Green Deal, Horizon Europe, and EU partnership for R&I, researchers are given the benefit of the doubt to halt and remove pollution [19-21, 39]. In this document, the application of living organisms native to polluted areas brings green solutions and deals with issues related to agricultural soil, farming, and unhealthy soils. Also, the freshwater issue and polluted oceans and seas can be addressed by bioremediation. Under Horizon Europe, the EU research and innovation program (2021-27) aims to bring solutions and results by 2030 with 5 missions. This research focuses on living labs and lighthouses within “A Soil Deal for Europe: 100 Living Labs and Lighthouses to Lead the Transition towards Healthy Soils by 2030 and Restore our Oceans and Waters by 2030” [20]. The term living labs refers to on-site dealing with pollution, while lighthouses will promote and share good practices. Mission hotspots include soil pollution, restoration, clean oceans, and freshwater. With 60-70% of European soils polluted and 100 living labs and lighthouses across the EU, the estimations are 30-40% remediated soil by 2030 [41]. The mission to restore oceans and waters includes lighthouses set to be enforced across the EU seas and rivers [42]. One important document, EU Circular Talks, brings bioremediation as techniques that use living organisms such as bacteria, fungi, and algae to reuse and recycle valuable materials, such as heavy metals and organic residues, from hazardous waste fractions. Also, in this context, bioremediation reduces soil and water pollution, thus removing it [43]. This study aims to present substantive examples of the capacity of some microorganisms to remove toxic substances from the environment. Therefore, the study focuses on the determinants in native and engineered organisms. The article aims to provide biochemical data valuable in cleaning particular pollutants *in vivo*/*in vitro*, showing scientific data available to researchers and stakeholders. The paper aims to clarify the potential of *in situ* biological techniques for their application to different inorganic or organic soil contaminants in the environment.

What Makes Microorganisms Valuable in Bioremediation

What is behind the ability to clean the environment from different types of pollutants? How do they effortlessly and successfully do it? All living organisms are determined by their genetic material (genotype) and their environmental interaction (phenotype/phenotypic plasticity).

The adaptive capacity of microorganisms, based on their phenotypic plasticity, confers a fitness advantage in different environments. The genomic plasticity of biodegradation strains enables unhindered growth and development in the presence of very toxic pollutants [44].

The great success lies in the fact that microorganisms can transform pollutants into useful micronutrients. Therefore, the catabolic mechanism of these organisms is subject to many research articles, with genes and proteins/enzymes being target participants. Although proven to break down very toxic organic pollutants, some artificial organic chemicals, such as xenobiotics, are difficult to remove. Also, partial degradation can create more harmful compounds than the targeted ones. Engineered microorganisms have the genetic capacity to overcome these obstacles with efficient catabolic pathways [15, 45-47].

Genetic engineering has been widely used when it comes to removing mixtures of pollutants. Many bacteria can now grow and naturally remove both organic and inorganic pollutants. Targeted enzyme genes are cloned and expressed for successful bioremediation applications. For instance, the genetically modified strain Pb2-1 and the *Rhizobium strain* 10320D, *Pseudomonas putida* 06909, can degrade trichloroethylene (PCBs) but can remove/degrade HMs (Cd^{2+}) by EC20 metal binding protein placed on the cell surface [46].

Finding appropriate bacterial strains that are able to remove mixtures of organic and inorganic substances in one place/polluted soil presents a quite difficult task [49]. Predominantly, inorganic pollutants such as NH_3 , NO_3 , and NH_4ClO_4 can be successfully transformed by microbes, while removing heavy metals is more demanding. This process involves a change in the valence states of metals. The catabolic pathway includes a genomic background. Genes that encode specific enzymes are located on chromosomes or belong to plasmids, transposons, or integrative-conjugative elements (extrachromosomal DNA). Molecular aspects of bioremediation via the catabolic pathway describe the strain's physiology and metabolism, the species' biodegradation potential, and homologous genes, directly indicating similarities and differences among different biodegradation strains [50]. Nowadays, omics technologies (genomics, transcriptomics, proteomics, and metabolomics) are the basis of successful remediation.

To be in a position to decide which microorganisms can apply or whether some should or need to be modified, microbe-based whole-cell systems and their enzymes are investigated by different biotechnological methods (direct DNA isolation, sequence-based techniques (Sanger sequencing (SS), capillary electrophoresis (CE), and next-generation sequencing (NGS), PCR methods) molecular fingerprint techniques (denaturing gradient gel electrophoresis (DGGE) and temperature gradient gel electrophoresis (TGGE)), fluorescent *in situ* hybridization (FISH), terminal-restriction fragment length polymorphism (TRFLP), single-strand conformation polymorphism (SSCP), etc. [51-53]. Several transport protein groups enable microorganisms to have the influx and efflux of different metal ions. This mechanism protects them from overtoxicity. Microorganism HM resistance is linked

to encoded metal-binding peptides metallothioneins (MTs) and enzymatically produced phytochelators (PCs). Also, there are various transporters belonging to Natural Resistance-Associated Macrophage Proteins (NRAMP), Zrt-Irt-like proteins (ZIP), Fe-Transporter (FTR), and Cu-transporter (CTR). The other group is responsible for HM exports from cells by the cation diffusion facilitator (CDF), P1B-type ATPases, FerroPortin (FPN), and the families Ca (II)-sensitive cross-complementin1/Vacuolar iron transporter1 (Ccc1/VIT1). Microalgae are well known for their successful removal of Zn. The most researched proteins are metallothioneins, which play a role in metal chelation. MT_s has several classes with different roles; MT_s class III is linked with successful Zn chelation. Although not that effective, MTs-class II is found in *Chlorella*, *Aureococcus*, *Symbiodinium*, *Nannochloropsis*, *Thalassiosira*, and *Ostreococcus* genera, also proven to be heavy metal-tolerant microorganisms [54]. Although MT-III is significantly more reactive than MTs-I/II, the class II MTs are polypeptides obtained from direct gene products and bind with Cu, Cd, Hg, and Zn [54]. Additionally, heavy metals can boost PAH biodegradation by enzyme-metal-substrate complexes. Since bioremediation of heavy metals is pollution concentration dependent. Being coenzymes, HMs stimulate a biological response in microorganisms towards PAH removal, even in a concentrated environment. Also, PAHs influence the microorganism's membrane, changing the cell membrane permeability and the electrical potential of the cell membrane, etc., and enhancing HM intake [55]. When tackling PAH removal, enzyme-based metabolic pathways include genes encoding oxygenase, peroxidases, reductases, hydroxylases, monooxygenases, and dehydrogenases [56]. In PAHs, aromatic ring cleavage, side chain, and central aromatic processes participate in two types of oxygenases: cytochrome P450 monooxygenases (CYPs) and ring-hydroxylating dioxygenases (RHDs). Microorganism PAH degradation results in CO₂ and H₂O. Furthermore, redesigning microorganisms for specific purposes can be achieved using synthetic biology tools. A broad spectrum of sequence-specific nucleases, known as gene editing tools, have been applied so microorganisms with new abilities can accomplish a given task. To manipulate any genomic sequence, target DNA double-strand breaks (DSBs) at specific sites and, if needed, repair them by non-homologous end joining (NHEJ) or homologous recombination (HR) to give a specific sequence. Systems biology and gene editing tools used to enhance bioremediation processes are the CRISPR-Cas system, ZFN, and TALEN. While ZFN and TALEN use artificial endonucleases, CRISPR-Cas uses naturally occurring RNA-directed endonucleases. CRISPR stands for clustered regularly interspaced short palindromic repeat DNA sequences with Cas-added protein. There are 3 CRISPR-Cas types and several subtypes defined by the Cas protein. The CRISPR-Cas 9 system consists of Cas9, a DNA endonuclease, and two noncoding

RNAs, crRNA and tracrRNA (guide RNA (gRNA)) [57]. CRISPR presents 30-40 bp of repetitive DNA sequences with "spacer" DNA sequences in between, with the repeats being translated into crRNA. CrRNA and Cas protein form crRNP (ribonucleoprotein), which causes a break in the intruder's DNA/RNA. In the Cas9-gRNA complex, gRNA directs Cas9 to cut both strands of a targeted region of DNA, forming the DSB of a target site [58, 59]. Recent research on prokaryotic CRISPR-Cas systems is leading to models in which inactive and RNA-targeting Cas proteins are coupled to a variety of effector proteins to regulate gene expression and epigenetic modifications, additionally preventing mutations [60]. More advanced tools, TALENs, or transcription activator-like (TAL) effector nucleases, are originally secreted from the pathogenic bacterial genus *Xanthomonas*. The DNA-binding domains of TALENs form repetitive sequences isolated from *Xanthomonas*. TALEs combine the carboxy-FokI cleavage domain (C-terminal) and amino-DNA-binding domain (N-terminal). The advantage of TALENs lies in using a 'protein-DNA code' that joins DNA-binding TALE repeat domains to target one of the DNA bases at the binding site. Further, Zinc Finger Nucleases (ZFNs) are artificial restriction enzymes that consist of 3-4 domains, each approximately 30 amino acids long, organized in a ββα motif. Like TALEs, they have the carboxy-FokI cleavage domain (C-terminal) isolated from *Flavobacterium okeanokoites*. These ZFPs have between 9 and 18 bp that can target a wide range of possible DNA sequences and present a zinc-finger DNA-binding domain surrounding the cleavage domain, depending on the target site [60]. All three above-mentioned gene editing tools improved the process of bioremediation.

Pesticides are known for their high toxicity, biomagnification, and forming secondary pollutants if bioremediation isn't completed correctly. Applying CRISPR-Cas, ZFN, and TALEN gene editing tools changes the outcome of bioremediation by improving microorganisms with functional pesticide degradation genes, improving pesticide degradation to less toxic compounds, thus preventing biomagnification [60]. To remediate the environment burdened with harmful compounds, the best thing to do is to heal it with organisms and inhabitants in that environment. If not equipped to break down pollution, researchers can help redesign them. This review aims to summarize the molecular aspects of bioremediation microbes, information gained via genome sequencing, and the phenotype plasticity of some microorganisms. Also, to present redesigned genetically modified microorganisms and specific genes that enabled them to remove target pollution. This data can then be used by researchers and enforced to facilitate decision-making. The natural methods applied in the environmental cleaning and maintenance process promise sustainability in every aspect (Fig. 1).




Fig. 1. Bioremediation towards environmental sustainability.

Results and Discussion

Implication of Microorganisms in Heavy Metal (HM) Removal

The pollution of soil and water caused by the release of heavy metals (HMs), such as chromium (Cr), cadmium (Cd), mercury (Hg), nickel (Ni), zinc (Zn), and lead (Pb), as a consequence, have adverse effects on the environment and human health. The issues related to HM cause nutrient imbalances and reduced soil quality, as well as persistent nature and bioaccumulation. Natural sources, such as minerals, weathering, erosion, and volcanoes, along with anthropogenic sources that include industry, mining, agricultural activities, emissions from improper waste management, etc., make this problem urgent to tackle

[7]. Heavy metals affect genetic material because they compete to bind to the same sites on DNA as essential ions. Therefore, they are considered mutagenic and carcinogenic [47]. Microbial living environments play a significant role in their performance. The pH of each habitat and microorganism is specific. Changed pH can affect the activity of enzymes, HM adsorption in microorganisms, and mobility of heavy metal ions in the soil. Most bacteria's best Pb^{2+} and Zn^{2+} removal performances are in the pH values 2.0-5.5 range. Also, it has been shown that pH below and above these values decreases microorganisms' potential to remove HM. For Cu, Ni, and Pb removal, the best result showed *Bacillus* sp. and *Pseudomonas* sp. at pH 5 and *Bacillus jeotgali* U3 at pH 7 [61]. The temperature also plays a crucial role. Cd^{2+} , Cr^{2+} , and Zn^{2+} removal best capacity for *Thiobacillus ferrooxidans*, *Thiobacillus acidophilus*,

and *Thiobacillus tepidarius* are at 25-27° C, while *Sulfolobus solfa-tataricus* and *Acidianus brierleyi* prefer extremely high temperatures. *Bacillus licheniformis* and *Bacillus jeotgali* U3 showed greater variations from 25 to 35 C [61]. The best capacity for Cu^{2+} and Cr^{6+} removal by *Ochrobactrum intermedium* LBr and *Cupriavidus metallidurans* CH34 was in the pH 6-7 range. While the temperatures differ, 37° C and 27°C, respectively [62].

The substrate concentration is another important factor. Different microorganisms remove Pb at different environmental factors. Temp (°C), optimal pH, initial Pb concentration (mg L^{-1}), Pb removal rate (%), or max Pb uptake (mg g^{-1}) by *Pseudomonas stutzeri*, *Enterobacter cloacae*, *Serratia marcescens*, *Alishewanella* sp. WH16-1 *Microbacterium oxydans* CM3, *Pseudomonas* sp., and *Bacillus subtilis* X3 was for 30°C, pH 7, 103.5 (mg L^{-1})>99.0%; 30°C pH 7, 7.2 (mg L^{-1}), 68%; 28°C, pH 7, 100 (mg L^{-1}), 97.57%; 37°C pH, 6 100 (mg L^{-1}) 84.13%; 30°C, pH 7.59, 400 (mg L^{-1}), 58.0%; 20°C, pH 7, 50 (mg L^{-1})>90.0%, 37°C, pH 4, 300 (mg L^{-1}) 192.05 mg g^{-1} , respectively. In Fungus Temp (°C), optimal pH, initial Pb concentration (mg L^{-1}), Pb removal rate (%) or max Pb uptake (mg g^{-1}) were: for *Bacillus cereus* 30°C, pH 5, 100 (mg L^{-1}), 75.6%; *Penicillium polonicum* 30°C, pH 5, 828.8 (mg L^{-1}), 90.3%; *Spergillus tubingensis* 30°C, pH 5, 828.8 (mg L^{-1}), >90.0%; *Aspergillus niger* 30°C, pH 5, 828.8 (mg L^{-1}), 97.0%; *Trichoderma asperellum* 30°C, pH 7, 250 (mg L^{-1}), 18.71%; *Rhizopus oryzae* 30°C, pH 7, 450 (mg L^{-1}) 33.76%; *Mucor irregularis* 30°C, pH 7, 100 (mg L^{-1}) 17.37% [63]. Heavy metals affect genetic material because they compete to bind to the same sites on DNA as essential ions. Therefore, they are considered mutagenic and carcinogenic [64].

Microorganisms are recognized as a powerful tool in bioremediation because of their natural and upgraded ability to degrade environmental pollutants via biochemical pathways. The genetic determinants involved in HM removal are presented in Table 1.

Chromium (Cr) Pollution

Chromium appears in the environment via natural and anthropogenic pathways. The two forms with positive and negative impacts on life forms are Cr(III) and Cr(VI) [65]. Industrial waste is a known source of toxic heavy metals such as hexavalent chromium. Because of high solubility and toxicity, Cr(VI) needs to be transformed into less toxic and soluble Cr(III). Chemically, graphene oxide (GO) polymers are found to be useful not only in absorption but also in reduction from Cr(VI) into Cr(III) [66]. The conversion of the highly toxic compound hexavalent chromium Cr(VI) into immobile and less toxic trivalent chromium Cr(III) can occur naturally. The chemical reduction process demands high energy, while biologically, the reduction process occurs in microorganisms and does not require high energy input [67]. Many species of microorganisms, *Arthrobacter*, *Pseudomonas*, *Pannonibacter*,

Escherichia, *Enterobacter*, *Bacillus*, *Shewanella*, etc., have been reported to reduce Cr(VI) [67-69].

Although Cr (VI) enzymes reductase are divided into two classes, class I has a role in chromium reduction, while class II relates to other pathways. Class I is encoded by ChrR and YieF genes, and Class II by NfsA (NiR genes) [69]. *Pannonibacter phragmitetus* BB tolerates high concentrations of Cr(VI) because of the presence of chrA1 and chrA2 that encode for two chromate transporters (ChrA1 and ChrA2) and nitR genes that encode a chromate reductase [69]. *Bacillus* sp. SFC 500-1E can tolerate high concentrations of hexavalent chromium, which is related to the chrA gene that encodes for the chromate ion transporter protein (ChrA) belonging to the CHR protein family [70]. It was proven that the modified *B. cereus* NWUAB01 chromate transport protein encoded by chrA has a role in chromate reduction and transport. Similarly, the *Bacillus* sp. strain DHS-12(7) chromate reductase (ChrR) gene encodes chromate reductase protein [71]. Also, it has been revealed that the coexpression of ChrA and ChrT in ChrAT-modified bacteria is linked to enzymatic activity via NADH and NADPH [71, 72]. The ChrA transporter, a membrane protein, has also been detected in *Bacillus cereus* strain TCL [72, 73]. Interestingly, lowering the concentration of Cr(VI) within a cell without changing its valence state (two chromate transporters) only protects cells/microorganisms but does not remove environmental toxins. Therefore, chromate reductase reduces heavy metal pollution in the environment. Also, the locations of genes that encode the chromate reductases (ChrR, YieF, NfsA, NemA, and LpDH) differ. Thus, in the cytoplasm (soluble chromate reductases) and cell membrane (membrane-bound chromate reductases), the reducing conditions are related to aerobic and anaerobic circumstances [69]. Furthermore, chrA genes can be found on plasmids as well as in chromosomal DNA. Therefore, as expected, they are organized in operons with other chr genes; for instance, in *C. metallidurans*, they are organized in the chrFECAB cluster as a part of a plasmid, or they are a part of the chromosome as the chr2 cluster (chrB2, chrA2, chrF2 genes). Furthermore, chr genes, chrB, chrA, chrC, and chrF found in a transposable operon (the chrBACF cluster) in *Ochrobactrum tritici* 5bv11 are involved in Cr resistance [5, 75]. A ChrR gene in *Stenotrophomonas maltophilia* encodes a protein Chromate Reductase (ChrR) homologous to other chromate reductase enzymes [76] that also provides resistance and high Cr(VI) reduction in the natural environment. Furthermore, *Stenotrophomonas maltophilia*, at the optimum 37°C at pH 7, showed the ability to remove and reduce up to 100%, making it a suitable candidate for bioremediation. The Cr(VI) concentration between 10 and 2000 mg/L showed 100% removal at concentrations in the 10-70 to 92% at 500 mg/L [76]. The transportation and reducing ability of transport proteins and Cr-reductase have been used to modify microorganisms that live in contaminated sites but have no Cr (VI)

Table 1. The molecular aspect of HM removal.

Gene/genetic determinant	Microorganism	Protein families	Effect	Reference
<i>ChrR</i> gene	<i>Bacillus</i> sp. strain <i>DHS-12(7)</i>	Chromate Reductase	Chromate reduction and transport	[70]
<i>chrA</i> and <i>chrT</i>	<i>ChrAT</i> engineered bacteria (<i>E. coli</i> BL21)	The CHR protein family, chromium resistance protein ChrA, and chromium reductant protein ChrT.	Chromate reductase	[71]
<i>chrA</i>	<i>Pseudomonas aeruginosa</i> and <i>Cupriavidus metallidurans</i>	The CHR protein family	Chromate transport	[70, 71]
<i>chrA1</i> and <i>chrA2</i> genes	<i>Pannonibacter phragmitetus</i> BB	Chromate Reductase	Chromate transporters (ChrA1 and ChrA2)	[73]
<i>chr2</i> cluster (<i>chrB2</i> , <i>chrA2</i> , and <i>chrF2</i> genes).	<i>Cupriavidus metallidurans</i>	The CHR protein family	Chromate transport	[74]
<i>chrBACF</i> cluster (<i>chr</i> genes-) <i>chrB</i> , <i>chrA</i> , <i>chrC</i> , and <i>chrF</i>)	<i>Ochrobactrum tritici</i> 5bv11			
<i>ChrR</i> gene	<i>Stenotrophomonas maltophilia</i>	Chromate Reductase (ChrR)	Resistance and high Cr (VI) reducing ability	[75]
<i>chrI</i> , <i>chrB</i> , <i>chrA</i> , <i>chrC</i> , <i>chrE</i> , <i>chrF</i> genes	<i>Ralstonia metallidurans</i> plasmid <i>pMOL28</i>	The CHR protein family	Chromate transport	[79]
<i>chrA_{so}</i> gene	<i>Shewanella oneidensis</i> MR1 and Genetically modified <i>E. coli</i>	Chromate efflux pump	Extracytoplasmic chromate reduction	[77]
<i>nitR</i> gene	<i>Ectomycorrhizal Fungus Pisolithus</i> sp.1	Class II chromate reductases	Chromium biotransformation	[77]
<i>NfsA</i> gene	<i>E. coli</i>	Class II chromate reductases	Chromium biotransformation	[77, 78]
<i>nikABCDE</i> operon	Coli	ABC transporter family	Protein related to Ni resistance and translocation (ABC-type nickel permease)	[81-83]
<i>nikABCDE</i> operon	<i>Bacillus altitudinis</i> MT422188 and <i>Bacillus altitudinis</i> FD48	ABC transporter family	Protein related to Ni resistance and translocation	[86-88]
<i>cadA</i>	<i>Microbacterium paraoxydans</i>	P-type ATPase	Cell detoxification and transporting (Zn^{2+}/Cd^{2+}) heavy metal ions across cellular membranes	[89]
<i>CDSs</i> genes	<i>Cupriavidus</i> sp.	CnrA, CnrB, CnrY, NccB, NccC, and NccN proteins	Proteins related to Ni resistance	[91]
		NccX protein	Protein related to Ni resistance	
<i>PbrR</i> operon	<i>Cupriavidus metallidurans</i> CH34	MerR family	Control heavy metal Pb toxicity	[98]
<i>pbrA</i> gene	<i>Pseudomonas fluorescens</i> M114	Pseudobactin M114	Reduce Pb overload in cell/Pb chelation	[100]
	<i>Pseudomonas</i> strain B10 and <i>Pseudomonas aeruginosa</i> PAO1	Pseudobactin B10		[101]

<i>psbA</i> gene	<i>Pseudomonas aeruginosa</i> mutant <i>PALS124</i>	Pseudobactin B10	Pb chelation	[101]
<i>NRPS</i> genes <i>mchA</i> and <i>mchB</i>	<i>Pseudomonas</i> sp. <i>NCIMB 10586</i>	Mupirochelin siderophore	Pb chelation	[102]
<i>CzcD</i> genes	<i>Streptococcus agalactiae</i>	Efflux proteins	Transport / Resistance Zn^{2+}	[107]
<i>smtA</i> and <i>smtB</i>	<i>Synechococcus</i> sp. strain <i>RRIMP N1</i> and <i>Synechococcus</i> <i>TX-20</i>	MT II class	Zn^{2+} and Cd^{2+} chelation in a cytosol	[123]
<i>SmtB</i>	<i>Escherichia coli</i> , <i>Staphylococcus aureus</i> , and <i>Staphylococcus xylosum</i>	MT II class	Zn^{2+} and Cd^{2+} chelation in a cytosol	[123, 124]
<i>bmtA</i> and <i>bxaI</i>	<i>Oscillatoria brevis</i>	A metallothionein and CPx-ATPase	Zn^{2+} and Cd^{2+} chelation in a cytosol and Export of Zn^{2+} and Cd^{2+} ions out of the cytomembrane	[109, 110]
<i>merA</i> and <i>merB</i> genes	Gram-negative and Gram-positive bacteria	Mercuric reductase and organomercurial lyase	reduction of Hg^{2+} to Hg^0 Lysis of C-Hg bond	[115, 116]
<i>merT</i> and <i>merP</i> with <i>merF</i>	Gram-negative and Gram-positive bacteria	Transporter proteins	Hg^{2+} transport into the cytoplasm	[115, 116]
<i>merR</i>	Gram-negative and Gram-positive bacteria	Regulator protein	Positive regulation of the mer operon	[116]
<i>merD</i>	Gram-negative and Gram-positive bacteria	Regulator protein	Negative regulation of the mer operon	[116]
<i>hgcAB</i> genes	<i>Nitrospirae</i> and <i>Chloroflexi Nitrospirae</i> and <i>Chloroflexi</i> (<i>Dehalococcoides</i>)	Hg methylation, corrinoid protein, HgcA, and a ferredoxin, HgcB	Mercury methylation/forming methylmercury (MeHg)	[118-121]
<i>merC</i>	Gram-negative and Gram-positive bacteria, Genetically Engineered <i>Acidithiobacillus ferrooxidans</i>	Transporter proteins	Transporter of mercuric ions	[116, 143]
<i>zntA</i>	<i>E. coli</i>	P-type (PIB) ATPases	Cell detoxification and transporting heavy metal ions across cellular membranes	[123, 126]
<i>copA</i>	<i>Legionella pneumophila</i>	P-type (PIB) ATPases	Cell detoxification and transporting (Zn^{2+}/Cd^{2+}) heavy metal ions across cellular membranes	[123, 126]
<i>zntA</i> ,	<i>Burkholderia</i> sp.	Zn^{2+}/Cd^{2+} exporting ATPase	Metals exporting	[131]
<i>czcA</i> ,		Heavy metal efflux system protein		
<i>czcB</i> ,		Membrane fusion protein		
<i>czcC</i>		Outer membrane protein		
<i>CzcD</i>	<i>Ralstonia metallidurans</i>	The cation diffusion facilitators (CDF family)	Decrease the concentration of Zn^{2+} and Cd^{2+} in cytoplasm	[136]
<i>czcA1 A.f. czcA2 A.f. czcB1 A.f. and czcC1 A</i>	<i>Acidithiobacillus ferrooxidans</i>	The cadmium efflux system (the Czc system)	Efflux pump protein	[137]

<i>czcCBA1</i> and <i>czcCBA2</i>	<i>Pseudomonas putida</i> KT2440	Czc system	Efflux pump protein	[138]
<i>Cad</i>	<i>Bacillus cereus</i> strain NWUAB01	Cadmium transporting ATPase, cadmium efflux system accessory protein, and cadmium resistance transporter	Cd transport protein	[139]
<i>The NR genes and the NiR genes</i>	<i>Ectomycorrhizal Fungus (EMF) fungi, such as Hebeloma sp., Tuber borchii, Laccaria bicolor, and Wilcoxina mikolae var. mikolae, Pisolithus sp., Paxillus ammonia virescens</i>	Class I reductase-Nitrate reductase (NR) and Class II reductase-nitrite reductase (NiR)	Chromium biotransformation	[142]
<i>MerH</i>	Genetically engineered <i>Deinococcus radiodurans</i>	Transporter protein	Resistance against mercury	[143]
<i>nixA</i>	Genetically engineered <i>E. coli</i> SE5000 strain	the NixA nickel permease	Degradation of nickel in water	[143]
<i>CzcR and CzcD genes</i>	<i>Alcaligenes eutrophus</i>	CzcR and CzcD proteins	Transport/Resistance to Co^{2+} , Zn^{2+} , and Cd^{2+} (czc system)	[144]

resistance. Transfer of Cr-transporter *chrA* genes and Cr-reducing *yieF* genes to *Comamonas aquatica* makes this genetically modified strain a very effective bioremediation microorganism [77]. Also, in chromium removal, there are enzymes nitrate reductase (NR) and nitrite reductase (NiR), which are related to N cycles in microorganisms. These enzymes, along with the nitrate and nitrogen intake in the NADP and NADPH domains, reduce hexavalent chromium and have been found in different microorganisms. The genes that encode for these enzymes are found in NR (*Pisolithus* sp. 1 NR) and NiR (*Pisolithus* sp. 1 NiR) of *Pisolithus* sp. 1 [78]. The text below contains information about identifying specific genes in *Vibrio harveyi* KCTC 2720. It also discusses the similarity of the reducing capacity of enzymes encoded by these genes to the *nfsA* gene found in *E. coli* [79]. Additionally, it mentions that some genes enable resistance to Cr(VI) in *Ralstonia metallidurans* *chr2*, specifically within the *chrB2* and *chrA2* genes, the *niaD* gene is the first to encode Cr(VI) reductase, and it has been identified in *Vibrio harveyi* KCTC 2720. Moreover, enzymes encoded by genes related to Nassimilation showed the same reducing capacity, such as the *nfsA* gene found in *E. coli* [79]. Apart from reducing capacity, some genes enable Cr(VI) resistance. In *Ralstonia metallidurans*, *chr2*, as a part of *chrB2*, *chrA2*, and *chrF2* gene clusters, has a role in chromate resistance. The chromate efflux pump encoded by the *chrA* gene, the previously known *chr₁*, belongs to *Ralstonia metallidurans* plasmid pMOL28 genes. These genes, along with *chrI*, *chrB₁*, *chrA₁*, *chrC*, *chrE*, *chrF₁*, and chromosomal *chr₂* cluster (genes *chrB₂*, *chrA₂*, *chrF₂*) enable Cr resistance [80]. Another promising bacterium, *Shewanella oneidensis* MR1, with chromosomal gene *chrA_{SO}*, encodes the chromate efflux pump protein. Furthermore, there are genetically modified strains, such as modified *E. coli* (heterologous expression techniques), with increased Cr(VI) reduction [81].

Nickel (Ni) Pollution

Nickel is an essential transitional trace heavy metal; being a micronutrient, it is found abundantly in the environment. Anthropogenic activities opposite from natural appearance have led to toxic levels of Ni in ecosystems. Then, it becomes lethal and immunosuppressive instead of essential for the metabolic activities in living cells. *Bacillus thuringiensis* KUNi1 was found to be resistant to Ni up to 82%. Furthermore, this strain is also resistant to zinc (Zn), copper (Cu), and cobalt (Co) at the same level of resistance, while in cadmium (Cd), small activity is shown [82]. Exploring the potential of *E. coli* in the Ni resistance mechanism, the *nik* operon has been identified. The *nik* operon consists of 6 genes (*nikABCDE*) that regulate Ni cell concentration. The first five encode for the ABC transporter system (ABC-type nickel permease) linked to translocation,

whereas genes *nikB* and *nikC* encode for proteins linked to the chemiosmotic gradient, while *nikD* and *nikE* encode for proteins that need energy provided by hydrolysis of ATP to translocate Ni. Interestingly, the first gene in the cluster, *nikA*, relates to chemotaxis [82-84]. Another gene, also part of this cluster, steps out, a *nikR* gene that can function as a repressor of the *nik* operon and as an activator of gene transcription [85, 86]. The transporter family encoded by *nikABCDE* in *Bacillus altitudinis* MT422188 is regulated by the ABC transporter family, ATP independent or ATP dependent [87], also known as ATP-binding cassette systems encoded by the same *nikABCDE* operon in *Bacillus altitudinis* FD48 [88, 89]. Furthermore, in Ni^{2+} resistance, secondary permeases of the NiCoT family (Ni-tolerant genes) were found in *Microbacterium oxydans* strains CM3 and CM7 and *Microbacterium paraoxydans*. The *M. paraoxydans* Ni resistance is linked to three members of Ni-binding proteins with different locations. The *nikABCDE* system includes *NikA* (periplasmic), *NikB*, and *NikC* (inner membrane) proteins involved in Ni uptake and bioaccumulation. Additionally, in *Microbacterium paraoxydans*, genes involved in Zn tolerance include *nikC*, an inner membrane protein, then *znuB* integral membrane protein, and *zur* negative regulator [90]. Engineered *Escherichia coli* bioaccumulate high concentrations of Ni by its native *NikABCDE* transporter and a heterologous, cysteine-rich metallothionein from the plant *Pisum sativum* [91]. Another group of genes (CDSs) showed involvement in Ni resistance. In the *Cupriavidus* sp. Ni-2 genome, 12 CDS genes encoding *CnrA*, *CnrB*, *CnrY*, *NccB*, *NccC*, and *NccN* proteins related to Ni resistance, and *NccX2+* proteins related to Ni transport were identified [92]. In addition, various members of the HoxN-family transporters proteins are involved in the Ni^{2+} resistance mechanism. For instance, HoxN is found in *Ralstonia eutropha* and used in *Streptomyces* sp., which are very effective microorganisms for high Ni concentration removal [93]. Actinomycetes showed a high ability to remove organic and inorganic waste from polluted soil and water. The focus can be directed towards the *Rhodococcus* genus because of its many-sided ability to produce enzymes and biosurfactants and degrade a wide range of pollutants. *Rhodococcus opacus* and *Rhodococcus rhodochrous* were identified as successful in Ni^{2+} and Pb^{2+} removal [94].

Lead (Pb) Pollution

Anthropogenic activities related to waste batteries, pigments, lead arsenate insecticides, lead water pipes, manufacturing, mining, industrial wastewater, pesticides, etc., release high amounts of lead into the environment [95]. Lead is widely recognized as a ubiquitous and very toxic metal, proven to inhibit enzyme activity, damage DNA, and disrupt cell membrane permeability in microorganisms. Metallothioneins, cysteine-rich proteins encoded by plasmid genes, are involved in the

sequestration or bioaccumulation of toxic metals within the cell [96]. Bacteria exhibit several detoxification mechanisms to reduce the harmful effects of heavy metals and protect themselves [97]. Several lead-resistant bacteria, including *Klebsiella michiganensis* R19, *Providencia rettgeri* L2, *Raoultella planticola* R3, *Serratia* sp. L30, *Pseudomonas pseudoalcaligenes*, and *Micrococcus luteus* are found to be useful in Pb removal [98]. A genetic determinant, a *PbrR* operon, activates a group of transcriptional regulators called the *MerR* family that controls heavy metal toxicity. This was found in *Cupriavidus metallidurans* CH34, which showed resistance in the presence of lead (II) [99, 100]. Another cluster of the *pbrABCD* genes also codes for Pb, Cd, and Zn transport proteins. Thus, *pbrAB* genes code for *PbrA* protein, a type of ATPase protein, and *PbrB*, an undecaprenyl pyrophosphate phosphate, with a role in lead sequestration. Notably, the *Pseudomonas fluorescens* M114 *pbrA* gene is responsible for the cellular response to iron via pseudobactin M114 synthesis [101]. A high-affinity iron-chelating Pseudobactin B10 regulated by the *pbrA* gene is also identified in *Pseudomonas* strain B10 and *Pseudomonas aeruginosa* PAO1, while in *Pseudomonas aeruginosa* mutant *PALS124* (*pvdA*), it is encoded by the *psbA* gene [102]. The soil bacterium *Pseudomonas* sp. *NCIMB10586* showed strong iron toxicity resistance, producing two mupirochelin siderophores by *NRPS* genes *mchA* and *mchB* [103]. Apart from bacteria, some microalgae are also known as organisms with high heavy metal tolerance. For instance, the genera *Chlorella* and *Scenedesmus* with certain species (*C. reinhardtii*, *C. sorokiniana*, and *S. obliquus*) can tolerate to some extent Pb(II), Cd(II), Cu(II), and Cr(VI), depending on the cell structure and metabolism [104]. Several transport protein groups enable the influx and efflux of different metal ions. This mechanism protects cells from overtoxicity. The study related to the Pb removal taxon, Chlorophyta with *Chlamydomonas reinhardtii*, *Chlorophyta*, *Chlorella vulgaris*, *Oedogonium* sp., and *Pseudochlorococcum typicum* has shown promising results [105]. Also, *C. acidophila* PM01 showed substantial tolerance to high concentrations of Cd, Cu, and Zn, with a promising increase in bioaccumulation of Cu due to metabolic adaptation [106].

Zinc (Zn) Pollution

Zinc is an essential element due to its role in bacterial resistance related to plasmids. The resistant strategy mechanism in Gram-negative and positive bacteria includes efflux, sequestration, chelation, and metal uptake. Bacteria's main efflux proteins are the Cation Diffusion Facilitator (CDF), Resistance Nodulation Division (RND) efflux pumps, and P-type ATPase families [107]. A Zn resistance can be linked with three structural genes (*czcABC*) that encode RND (*CzcA*, *CzcB*) and CDF (*CzcD*) efflux pumps for Zn^{2+} transport out of the cell, and they are found in both Gram-positive

and negative bacteria. In *Streptococcus agalactiae*, resistance is related to *czcD* genes that encode an efflux protein [108], as well as the CPx-type (CadA) efflux proteins found in Gram-positive bacteria. For the metal Zn uptake, Gram-positive bacteria have P-type ATPase (ZosA), *znuABC*, and membrane proteins (YciABC, Ycd H1). However, the P-type ATPase protein ZntA participates in the efflux of Zn^{2+} in both Gram-positive and negative bacteria [108]. Furthermore, the ABC transporter family belongs to the high-affinity Zn^{2+} uptake systems and presents members of the ATP-binding cassette, with *znuA*, *znuC*, and *znuB* proteins encoded by *znuABC* genes [109]. In sequestration, the role has periplasmic or cytoplasmic proteins [108]. In *Escherichia coli*, Zn detoxification is related to the CDF family with the ZitB and YiiP proteins, one P-type ATPase, and the ZntA protein. In *Staphylococcus aureus*, resistance enables the CDF family. In *Caulobacter crescentus*, *czrCBA* and *nczCBA* belong to the RND efflux systems. *C. crescentus*, CzcCBA system exports Zn ions from the cytoplasm of cells, similar to the CzcABC system, while in *Ralstonia metallidurans*, there is the CzcS/CzcR two-component regulatory system. In *Streptococcus pneumoniae*, that is CzcD, a member of the CDF family [109]. Bacteria possessing MTs are an ideal tool for the Zn and Cd bioremediation. The marine cyanobacterium, *Synechococcus* sp. strain RRIMP N1, and a freshwater *Synechococcus* TX-20 can remove Zn by MT class II encoded by the *smtA* gene. Another important gene is the *smtB* gene, which plays a role in the metal-dependent repression of the *smtA* gene. In Zn and Cd removal, *Escherichia coli*, *Staphylococcus aureus*, and *Staphylococcus xylosus* SmtB gene and ArsR proteins are also involved. Also, *bmtA* and *bxaI* genes of *Oscillatoria brevis* encode metallothionein (BmtA) and CPx-ATPase (BxaI) [110, 111]. A CPx-ATPase-like ABC transporter, CDF families, etc., export metal ions out of cytotcells [111]. Thus, these microorganisms are capable of chelating and exporting metal ions, therefore protecting themselves and their habitat.

Mercury (Hg) Pollution

Mercury (Hg), previously known as hydragyrum, stands out with a unique ability to keep liquid form at Earth's temperature and pressure [112]. Recognized as a metal of great toxicity, its removal from the environment is an issue to be tackled. In addition, Hg has no identified biological role opposite to Fe, Cu, Zn, Se, Cr, etc. [113]. The ability of some microorganisms to live in the presence of high concentrations of Hg includes some chemical or biological change of mercury [114]. One of the mechanisms is the reduction of Hg^{2+} to Hg^0 , which is controlled by the *mer* operon, a cluster of genes involved in Hg transport, reduction, and volatilization [115]. A cytoplasmic enzyme, mercuric reductase, and an organomercurial lyase are encoded by *merA* and *merB* genes, respectively [116]. Besides *merA* and

merB genes, identified *mer* operons with *merR*, *merP*, *merT*, *merD*, *merF*, and *merC* encode for different inner-membrane proteins with different roles in Hg transport and metabolic pathways. Namely, *merR* encodes for MerR, the metalloregulatory protein, a regulator of the operon transcription in the presence/absence of Hg^{2+} ; *merD* encodes for the MerD regulator protein, a secondary regulator protein with a similar but weaker influence on the operon transcription [116]. Other important genes are *merT* and *merP*, and sometimes with *merF*, help Hg^{2+} transport into the cytoplasm [117, 118]. Hg^{+2} to Hg^0 reduction mechanism can be independent of the *mer* system and has been found in *Geobacter sulfurreducens* PCA, *Geothrix fermentans* H5, *Cupriavidus metallidurans* AE104, *Shenwellia oneidensis* MR-1, and *Geobacter metallireducens* GS-15. The primary drivers of Hg methylation are the corrinoid protein, HgcA, and a ferredoxin, HgcB, encoded by *hgcAB* genes [119]. The diversity of HgcAB-encoding microbes is reflected in aerobic nitrite-oxidizing bacteria *Nitrospirae* and *Chloroflexi* (Dehalococcoides) [120, 121]. Important methylating genes, *hgcAB*, are also found in mercury methylators from *Deltaproteobacteria*, *Firmicutes*, *Chloroflexi*, *Phycisphaerae*, *Aminicenantes*, *Spirochaetes*, *Elusimicrobia*, *Lentisphaerae*, *Bacteroidetes*, and *Atribacteria* [121, 122].

Cd Pollution

Among heavy metals, Cd is classified as highly toxic and difficult to degrade; thus, pollution with this metal is serious. Its effect on living organisms is reflected in reducing the biological activity of soil microorganisms and plants, which can be linked to animal and human poisoning by the food chain [123]. Many microorganisms have developed resistance to this metal, and their activity is not disturbed [123]. *Staphylococcus aureus* strain is known for its natural Cd resistance; *Cupriavidus metallidurans*, *Escherichia coli*, and *Bacillus subtilis* are also applied to Cd removal. The basis for the resistance mechanism lies in the presence of specific resistance genes or operons like *cad*, *czc*, and *znt*. These genes encode for P-type ATPases and cation diffusion facilitators (CDF) that transport Cd from the cytoplasm to the periplasm and RND-driven efflux systems that transport Cd from the periplasm to outside the cells. Therefore, export from the cell is complete [124]. Some studies indicate the role of strain Y8/genome with more than 75% guanine-cytosine content (CG). According to Chen et al. (2019a, b), metal-resistant strains such as *Cupriavidus*, *Thiobacillus*, *Deinococcus*, and *Comamonas* spp. have an average GC content of 66.2%, 62.6%, 67.3%, and 61.5%, respectively [125, 126]. Genes identified as Cd metal resistance genes *zntA* of *E. coli* and *copA* of *Legionella pneumophila*, along with the ACR3 family genes, encode for two of four heavy-metal transporting P-type (PIB) ATPases [124]. These transporters are essential in cell detoxification,

transporting heavy metal ions across cellular membranes [127]. Interestingly, genes *czcD* of *Cellulomonas metallidurans*, *cutC* of *Enterococcus faecalis*, and *znuA* of *E. coli* are homologous to genes identified in Y8 strains [124]. Specifically, *zntAY8* and *czcDY8* in Y8's *E. coli* *zntAY8* and the genus *Cellulomonas* with (*zntAY8*, *copAY8*, *HMTY8*, and *czcDY8*). Furthermore, the recombinant *E. coli* metallothionein gene, cloned from *Neurospora crassa* belonging to the fungal class Ascomycetes, is linked with Cd uptake. Furthermore, *cadA* and *norVW* operons encode for *CadA* and *NorVW* involved in Cd resistance in the highly cadmium-resistant strain *Enterobacter cloacae* LPY6/ domesticated *E. cloacae* Cu6 [128]. Genetically modified *Pseudomonas putida* KT2440 showed great ability in Cd removal. The PC synthase gene (PCS) originated from the fungus *Schizosaccharomyces pombe* and was cloned and expressed in *P. putida* KT2440. The PC synthase gene (PCS) encodes for phytochelatins (PCs), an enzyme rich in the cysteine thiolate groups, the primary binding sites for heavy metals [129, 130]. *Pseudomonas putida*'s ability to remove copper and cadmium lies in plasmid-encoded heavy metal ion resistance [131]. Furthermore, the *Microbacterium paraoxydans* gene *cadA* encodes for Cadmium P-type ATPase [73]. *Burkholderia* sp. SRB-1 genome contains metal-exporting genes *zntA*, *czcA*, *czcB*, and *czcC*, and they encode for Zn^{2+} / Cd^{2+} -exporting ATPase, heavy metal efflux system protein, membrane fusion protein, and outer membrane protein, respectively. Additionally, detoxification *dsbA* and *cysM* genes encode for the proteins dithiol oxidoreductase and S-sulfo-L-cysteine synthase, and genes *katE*, *katG*, and *SOD1* encode for catalase, catalase-peroxidase, and superoxide dismutase, Cu–Zn family involved in Cd antioxygenation/antioxygenative enzyme activity [132]. The phosphate-mineralizing bacteria *Enterobacter* sp. shows high Cd resistance via mechanisms that include Cd ion transport and efflux, as well as the reduction and antioxygenation potential. Genes identified in PMB bacteria encode proteins of the Cd resistance mechanisms. *Enterobacter* sp. genes *czcD*, *MgtA*, and *zntB* encode for ion metal transport protein, *SOD2*, and *katE*, encode for enzymatic detoxification and *dsbA*, *cysM* antioxidant enzymes [133]. Different efflux pump proteins encoded by *cadA* and *cadB* plasmid R genes belonging to *Staphylococcus aureus* also play a crucial role in Cd bacteria resistance [134, 135]. *Alcaligenes eutrophus* and *Ralstonia* sp. CH34 are the *Cad* systems encoded by the *CadAB* operon for cadmium resistance [135, 136]. The cation diffusion facilitators (CDF family) resistance to Zn^{2+} and Cd^{2+} , encoded by *CzcD* from *R. metallidurans* strain CH34, decreases the concentration of Zn^{2+} and Cd^{2+} in the cytoplasm [137]. Similarly, *Acidithiobacillus ferrooxidans* *czcA1* A.f, *czcA2* A.f, *czcB1* A.f, and *czcC1* A.f genes belong to the cadmium efflux system (the *Czc* system) [138]. Furthermore, in *Pseudomonas putida* KT2440, *Cad* and *Czc* systems are also identified in Cd resistance. Namely, a regulator of the *CzcRS3*

system activates *czcCBA1* and *czcCBA2* genes that encode efflux pump protein [139]. Additionally, *Cad* genes of *Bacillus cereus* strain NWUAB01. also encode for cadmium transporting ATPase, cadmium efflux system accessory protein, and cadmium resistance transporter, all belonging to Cd transport proteins. Cytoplasmic copper homeostasis proteins encoded by *cutC* with copper resistance proteins encoded by *CopC* or *CopD* have a role in copper resistance transport. *CorA* genes encode for *CorA* proteins, *CorA*-like magnesium transport proteins, and magnesium transport proteins, all of which are involved in Mg transport. The manganese transport protein encoded by the *MntH* gene has a role in manganese transport. Cobalt–zinc–cadmium resistance protein encoded by the *CzcD* gene has a role in cadmium, cobalt, and zinc transport. Zinc ABC transporter is a periplasmic binding protein encoded by the *ZnuA* gene that has a role in zinc transport. Both *CzcD* and *cadA* operons are energy-dependent efflux systems [139], similar to the *PbrA* efflux protein, which gains energy by ATP hydrolysis and is encoded by the *PbrA* gene [140, 141]. The *CzcR* and *CzcD* genes encode for the *CzcR* and *CzcD* proteins that are implicated in resistance to Co^{2+} , Zn^{2+} , and Cd^{2+} (*czc* system) [142]. *Pseudomonas aeruginosa* T-3 has plasmid-encoded chromate ion resistance [143, 144], as well as *Pseudomonas putida* PhCN for Cd ion. Furthermore, *Pseudomonas aeruginosa* strain ZM130 resistance is linked with the enzyme laccase and nicotinamide adenine dinucleotide (reduced)-dichlorophenolindophenol reductase (NADH-DCIP reductase) [145].

Persistent Organic Pollutants POPs-Polycyclic Aromatic Hydrocarbons (PAHs) and Polychlorinated Biphenyls (PCBs)

Toxic, hydrophobic, and lipophilic persistent organic pollutants (POPs) can be found in soil, water, and air, as well as in living organisms. These include many manufactured chemicals, but the 3 main groups are polycyclic aromatic hydrocarbons (PAHs), polychlorinated biphenyls (PCBs), and pesticides [146]. Besides biochemical pathways, the physicochemical parameters (pH, temperature, and substrate concentration) play a role in microorganism performance. Benzo[a]pyrene (PAHs), degraded by *Sphingobacterium spiritivorum* and *Aspergillus brasiliensis*, was performed at pH 6 and 40 mg/L benzo[a]pyrene concentration and pH 6.41 and 38.8 mg/L benzo[a]pyrene concentration, respectively [147]. Although fungi prefer low acidic and high alkaline environments, it was noticed that a soil pH of 7.5 was the most suitable for the degradation of all the PAHs, reaching 50. *Penicillium* sp. prefers acidic pH, whilst *Aspergillus* sp. was found to be effective at pH 7.5–8 [148]. In POPs degradation, the best-removing capacity is *Pseudomonas putida* F6 at 30°C, optimum pH 7, and *Streptomyces cyaneus* at 60°C, optimum pH 4.5. *Bacillus* sp. showed variations

depending on the organic compound in the temperature between 30-37° C, optimum pH 5-8, and at similar conditions, *Mycobacterium* sp. degrading xenobiotics, *Sphingobacterium* sp. PLA, and *Chromobacterium viscosum* PBC [149]. The widespread use of many of them, as a consequence, has led to bioaccumulation and magnification. The problem needs to be addressed in a way that cleans but does not harm ecosystems; therefore, microorganisms present the best solution for POP removal. The genetic determinants involved in POP bioremediation are presented in Table 2.

Implication of Microorganisms in Polycyclic Aromatic Hydrocarbons (PAH) Removal

Polycyclic aromatic hydrocarbons (PAHs) are a class of organic compounds produced by incomplete combustion or pyrolysis of fossil fuels (oil, petroleum, gas, coal, and wood). PAHs are a type of organic compound that consists of 3 benzene rings (low molecular weight PAHs) or more (high molecular weight PAHs) [150, 151]. There are many methods of removing PAHs from the environment. Chemically, enzymes oxidize PAHs to dihydrodiol, then dioxygenase→intradiol and extradiol cleavage→protocatechuates and catechol [152].

Biologically, specialized microorganisms isolated from their natural habitats successfully break down PAHs in the oxidation process. The outcome is the conversion of hydrocarbon molecules into carbon dioxide, biomass, or other harmful molecules. The catabolic pathway includes catabolic genes that encode enzymes involved in mineralizing persistent organic pollutants. Several microbial communities have been applied in PAH removal strategies with *nidA/B*, *pah*, and *phd* genes, designated as important biomarkers of pyrene metabolism [153]. An abundant pollutant, pyrene, belongs to high molecular weight PAHs with a tetracyclic structure, and bacteria can very successfully remove it from different phyla. For instance, *Mycobacterium* spp. is recognized as highly effective, along with other genera, such as *Pseudomonas* sp. *JPN2*, *Staphylococcus* sp. *BJ06*, *Bacillus vallismortis*, *Diaphorobacter*, and *Pseudoxanthomonas*. Wang et al. (2023) applied metagenomics analysis to identify the degradation pathways of the identified genes and their contribution to this process. The consortium of bacteria involved in pyrene degradation consists of Proteobacteria (99.8%) and Alphaproteobacteria (96.6%). This analysis revealed that *nidA* and *nidB* genes encoding PAH dioxygenase α/β subunits catalyze the first step in the degradation process. Duplication of the dioxygenase *nid* gene *nidA/nidB* and *nidA3/nidB3* plays a key role in the initial degradation of pyrene. The following steps in degradation involve genes such as *phdE*, *phdF*, *phdG*, *phdK* (phenanthrene), *nidD* (pyrene), *phtAa/Ab/Ac/Ad*, *phtB*, *phtC* (phthalate), and *pcaG/H* [154]. Although highly effective, applying only one genus is yet to be proven to be the best bioremediation/solution. For instance, the PAH-removing consortium of anaerobic

Table 2. The molecular aspect of POPs removal.

Gene/genetic determinant	Microorganism	Protein families	Effect	Reference
<i>nidA</i> and <i>nidB</i> <i>nidA3/nidB3</i>	<i>Mycobacterium ghluum</i> PYR. GCK <i>Mycobacterium vanbaalenii</i> PYR-1	PAH dioxygenase α/β subunits	Pyrene degradation	[154-156, 161].
<i>nahAa</i> , <i>nahAb</i> , <i>nahAc</i> , and <i>nahAd</i> <i>nah4cR</i>	<i>Pseudomonas putida</i> G7 <i>Pseudomonas fluorescens</i> AH-40	Naphthalene dioxygenase	Naphthalene degradation	[159-161]
<i>nahA</i>	<i>Pseudomonas putida</i> strain G7 (pNAH7) cis-dihydrodiol naphthalene dehydrogenase 1,2-dihydroxy naphthalene dioxygenase 2-hydroxychromene2-carboxylate isomerase 2-hydroxybenzalpyruvate aldolase salicylaldehyde dehydrogenase	Naphthalene dioxygenase	The naphthalene conversion proceeded via salicylate	[153, 163, 191]
<i>nahB</i>				
<i>nahC</i>				
<i>nahD</i>				
<i>nahE</i>				
<i>nahF</i>				

<i>nahG</i>	<i>Pseudomonas putida</i> strain G7 (pNAH7) catechol 2,3-oxygenase 2-hydroxymuconate semialdehyde dehydrogenase 2-hydroxymuconate tautomerase 4-oxalocrotonate decarboxylase catechol 1,2-oxygenase	salicylate hydroxylase C120, catechol 1,2-oxygenase.	catechol further oxidized by the meta-pathway [163, 191]
<i>nahH</i>			
<i>nahI</i>			
<i>nahJ</i>			
<i>nahK</i>			
<i>nahM</i>			
<i>nahR</i>	<i>Pseudomonas</i> sp.	NahR regulatory protein	Transcriptional regulator for naphthalene degradation [160, 185]
<i>narAa, narAb</i>	<i>Rhodococcus</i> sp.	the α - and β - subunits of the naphthalene dioxygenase (NDO)	Naphthalene degradation [164]
<i>pcaG and pcaH</i>	<i>Pseudomonas</i> sp. strain HR199, <i>Stenotrophomonas maltophilia</i> KB2 <i>Paracoccus denitrificans</i> , <i>Devosia</i> , <i>Pusillimonas caeni</i> , and <i>Eoetvoestia caeni</i>	Two subunits of protocatechuate 3,4-dioxygenase	The enzyme involved in vanillin catabolism [166, 167]
<i>vdh vanAB, pbdA, pbdB, pbdL</i>	<i>Rhodococcus jostii</i> RHAI <i>Amycolatopsis</i> sp. strain ATCC 39116	vanillin dehydrogenase and vanillate <i>O</i> -demethylase	The enzyme involved in vanillin catabolism [177, 169]
<i>phtAc</i>	<i>Arthrobacter keyseri</i> 12B	Oxygenase ferredoxin component	Pyrene degradation and metabolism [154, 170]
<i>phdI, phdF</i>	<i>Mycolicibacterim</i> sp., <i>Mycobacterium vanbaalenii</i> PYR-1	Ring cleavage dioxygenase	
<i>phdJ</i>	<i>Nocardioideis</i> sp. strain KP7	Hydratase-aldolase	
<i>phtC</i>	<i>Nocardioideis</i> sp. strain KP7	Decarboxylase	
<i>xylQ</i>	<i>Arthrobacter keyseri</i> 12B	acetaldehyde dehydrogenase,	
<i>nahE</i>	<i>Sphingomonas paucimobilis</i> 20006FA a putative 2-hydroxybenzylpyruvate aldolase glutathione S-transferase		
<i>bphK</i>			
<i>panABC</i> (gene cluster)	<i>Burkholderia</i> sp. strain RP007	The iron-sulfur protein α and β subunits of a dioxygenase, the dihydrodiol dehydrogenase, and the extradiol dioxygenase	The naphthalene and phenanthrene oxidation [176]
<i>pheA, pheB, and C120 gene</i>	<i>Klebsiella pneumoniae</i> I, <i>Bacillus cereus</i> , <i>Pseudomonas montellii</i> , <i>Bacillus subtilis</i> , <i>Pseudomonas mosselii</i> , <i>Staphylococcus equorum</i> , <i>Bacillus benzoevorans</i> , <i>Bacillus circulans</i> , <i>Pseudomonas fulva</i> , <i>Pseudomonas aeruginosa</i> II, <i>Pseudomonas putida</i> , <i>Burkholderia cepacia</i> , <i>Bacillus cereus</i> , and <i>Klebsiella pneumoniae</i> II, <i>Pseudomonas putida</i>	Phenol hydroxylase, catechol 1,2-dioxygenase, and catechol 2,3-dioxygenase,	Phenanthrene and pyrene degradation [180]

C23O gene	<i>Pseudomonas</i> sp. Strain ND6 EAU29438 <i>Ewingella Americana</i> , <i>Bacillus megaterium</i> strain IV22	The catechol 2,3-dioxygenase (C23O) gene	Naphthalene catabolism, crude oil degradation	[181, 194, 195]
strain C12O genes (<i>catA</i> , <i>catA_I</i> , and <i>catA_{III}</i>)	<i>Pseudomonas</i> sp. Strain ND6	The catechol 1,2-dioxygenase	Naphthalene catabolism	[182]
1,2 and 2,3 CTD genes	The bacterial genera <i>Acinetobacter</i> and <i>Pseudomonas</i>	The catechol dioxygenase	Naphthalene catabolism	[186, 195]
the Bph genes/degrading gene clusters	<i>Geobacillus</i> sp. strain JF8 pBt40, <i>Rhodococcus</i> sp. linear pRHL1, <i>Cupriavidus</i> sp. pSK-4, <i>Pseudomonas putida</i> MB1335	dehydrogenase	Biphenyl catabolic enzymes (co-metabolize certain PCBs into chlorobenzoic acids, and a substitute of chlorine)/metabolism of aromatic compounds	[186-189]
bph gene cluster (<i>bphEGFAI</i> A2A3BCDA4)	<i>Cupriavidus basilensis</i> KF708 <i>Pseudomonas</i> sp. biphenyl dioxygenase			
<i>bphA1A2A3A4</i>				
ICEbph-sal (<i>bhp</i> and <i>sal</i> gene cluster)	<i>Pseudomonas putida</i> KF715	SalR protein	Biphenyl and salicylate catabolism	[190]
<i>salA</i>	<i>Acinetobacter</i> sp. strain ADP1		Salicylate hydroxylase activator	[191]
<i>sal</i> genes	<i>Pseudomonas putida</i> S-1		Conversion of alkyl salicylates into catechol	

bacteria affiliated with *Diaphorobacter* and *Paracoccus* genera showed high removal capacity for acenaphthene and phenanthrene. A myriad of different genes, *pht2/3*, *pht4*, and *pht5* (phthalate degradation) and *ligA/ligB* (protocatechuate degradation) was detected in *Achromobacter spanius Devosia* sp., *Pusillimonas caeni*, *Eoetvoesia caeni*, and *Bradyrhizobium* sp., *Bosea* sp., *Afipia* sp., *Steroidobacter* sp., and *Mesorhizobium* sp. [155]. The newly discovered strain *Pseudomonas veronii* SM-20 showed remarkable results in degrading/oxidizing phenanthrene [156]. Apart from genera (genomes) that naturally degrade aromatic compounds, horizontal gene transfer enables the employment of different bacterial consortiums. Similarly, the presence and influence of *nidA* in *Mycobacterium gilvum* PYR. GCK [157], the *nidA/B* gene in *Mycobacterium vanbaalenii* PYR-1 [158], and the *nidA*, *nidA3*, *pdoA2*, and *pcaH* genes were identified [159]. The *Pseudomonas putida* G7 (*nahAa*, *nahAb*, *nahAc*, and *nahAd*) *nahAb* gene that encodes for the naphthalene dioxygenase protein is involved in naphthalene degradation and is typical for all *nah*-type degrading bacteria. The *NahR* protein is a transcription factor encoded by the *NahR* gene that activates the *NAH7* plasmid operons [160]. Another strain, *Pseudomonas fluorescens* AH-40 with the *nahAcR* gene, was involved in naphthalene degradation [161]. *P. putida* strain G7 (pNAH7), where the naphthalene conversion proceeds via salicylate (*nah*-operon, *nahABCDEF*) and catechol is further oxidized by the meta-pathway (*sal*-operon, *nahGHIJKLM*). *Pseudomonas*-like organisms with a *nah*-like genotype were identified using genetic markers with *nah*-like sequences. In this way, markers *nahAc* from *P. putida* G7, *phnAc* from *Burkholderia* RP007, and the GST-encoding gene from *Sphingomonas* WP01 were revealed. The analogs of the bacterial glutathione S-transferase GST-encoding gene from *Sphingomonas paucimobilis* EPA505 were also identified [162]. Furthermore, *P. putida* KT2440 *NAH7* plasmid genes are under salicylate control, thus encoding for enzymes involved in naphthalene (PAH) mineralization/degradation [163]. In naphthalene degradation, two *NAH7* operons (gene clusters) participate with genes *nahA-F* and *nahG-M* that encode enzymes for the metabolism of naphthalene to salicylate and salicylate to tricarboxylic acid, respectively [164]. In *Mycobacterium vanbaalenii* PYR-1, *nidB2* and *nidB-nidA* gene clusters were similar to *phtAa* and *phtAb*. They encode α/β -subunits of phthalate dioxygenase, involved in the catabolic pathway of phenanthrene and pyrene. Furthermore, *Mycobacterium* sp. 6PY1 *pdoA1/pdoB1* and *pdoA2/pdoB2* genes are found to be similar to *Mycobacterium vanbaalenii* PYR-1 *nidA* and *nidB* genes. The gene cluster, including *nidB2*, *nidB/nidA*, and *nidA3/nidB3*, also encodes pyrene/fluoranthene/naphthalene/phenanthrene ring-hydroxylating dioxygenase α - and β -subunits [165]. In the same fungi, for instance, in *Rhodococcus* sp., naphthalene degradation participates only in the *narAa*, *narAb*,

and narB genes, which act as single units not organized in clusters, and they are regulated by narR1, narR2, and GntR-like transcriptional regulators. The narAa and narAb genes code for the α - and β - subunits of the naphthalene dioxygenase (NDO) with no similarity to genes in *P. putida* that encode NDO subunits [166]. Furthermore, *Rhodococcus* sp. P14 is mentioned in several polycyclic aromatic hydrocarbons via serine hydrolase, hydratase, alcohol dehydrogenase, protocatechuate 3,4-dioxygenase, β -ketoadipate CoA transferase, and β -ketoadipyl CoA thiolase. These enzymes are linked to ring cleavage, side chains, and central aromatic processes. Eight of the thirteen MAGs were confirmed to carry genes for PAH degradation and were affiliated mainly with *Actinobacteriota* and *Proteobacteria*. For instance, genes pcaG and pcaH are encoded for two subunits of protocatechuate 3,4-dioxygenase, the enzyme essential in vanillin catabolism in *Pseudomonas* strain HR199 [167], and in many xenobiotic metabolic reactions in *Stenotrophomonas maltophilia* KB2 [168]. These genes were also found in three other retrieved MAGs affiliated with *Paracoccus denitrificans*, *Devosia*, *Pusillimonas caeni*, and *Eoetvoesia caeni*. Also, genes vdh and vanAB, pbdA, pbdB, and pcaL encoding vanillin dehydrogenase and vanillate O-demethylase participate in vanillin catabolism, and they were found in *Rhodococcus jostii* RHA1 [169]. In addition to pcaG and pcaH, genes phdI, phdJ, phtAc, and phtC were identified in *Mycolicibacterium* sp. and *Mycobacterium vanbaalenii* PYR-1, encoding for enzymes that participate in pyrene degradation and metabolism [170]. In pyrene degradation and metabolism, several enzymes are encoded by previously mentioned genes isolated from different organisms: oxygenase ferredoxin component (phtAc) from *Arthrobacter keyseri* 12B, ring cleavage dioxygenase (phdI) from *Nocardioides* sp. strain KP7, and phdF from *Sphingomonas* sp. strain KA1, hydratase-aldolase (PhdJ) from *Nocardioides* sp. strain KP7, and decarboxylase (phtC) from *Arthrobacter keyseri* 12B [171]. An *S. paucimobilis* 20006FA shows similarities in the gene clusters bph and xyl with the plasmid pNL1. This plasmid with the target genes is found in the *Sphingomonas* sp. and *Sphingobium* sp. C100 [172] and *Sphingobium yanoikuyae* B1 genomes [173, 174]. Genes belonging to *Sphingomonas paucimobilis* 20006FA, xylQ, nahE, and bphK encode for enzymes involved in phenanthrene degradation, namely, acetaldehyde dehydrogenase, a putative 2-hydroxy benzyl pyruvate aldolase, and glutathione S-transferase, respectively [175]. Besides, cytochrome P450 monooxygenases (CYPs) and laccase (the MtL enzyme) show a high ability for PAH degradation. The MtL oxidoreductase enzyme belongs to ligninolytic enzymes and is characteristic of fungi. For bioremediation, originally found in *Aspergillus oryzae*, the laccase MtL gene was transferred to *Myceliophthora thermophila* [176]. Apart from the classical catabolic nah-like genes, the phn genes are detected on the phn

locus in *Burkholderia* sp. strain RP007. The cluster phnABC was under the regulatory control of PhnR and PhnS, encoding for different enzymes. While *Burkholderia* sp. strain RP007 phnABC cluster showed selective oxidation of either naphthalene or phenanthrene, transgenic *E. coli* phn enzymes oxidized both naphthalene and phenanthrene [177]. Furthermore, alkB and alkH, besides C12O and C23O, showed high catabolic potential in PAH removal. Bacterial strains *Enterococcus faecalis* (MK298392.1), *Bacillus albus* (MT325971), *Enterococcus faecalis* (MT325972), *Providencia vermicola* (MT325973), *Enterococcus faecalis* (MT345788), *Carnobacterium gallinarum* (MT350233), and *Pseudomonas aeruginosa* (JF756593.1) isolated from landfill soil artificially polluted with waste engine oil showed increased alkane hydroxylase gene (alkB) and aldehyde dehydrogenase gene (alkH) activity [178].

Polychlorinated Biphenyls (PCBs)

Polychlorinated biphenyls (PCBs) are aromatic synthetic organochlorine chemicals with widespread use in transformers and electrical capacitors. PCBs are also used as hydraulic fluid, oil additives for paints, etc. As the name indicates, this biphenyl molecule consists of two six-carbon rings linked by a single carbon-carbon bond. In the PCB molecule, hydrogen is substituted with chlorine and bonds to 12 carbon atoms [179]. The phenol degradation pathway includes the activity of estradiol-cleaving and estradiol-cleaving enzymes encoded by plasmid genes. Some of the enzymes involved in PBC degradation are catechol 2, 3-dioxygenase (meta-fission), catechol 1,2-dioxygenase (ortho-fission), and phenol hydroxylase. Namely, phenol hydroxylase encoded by pheA and pheB was found in *Klebsiella pneumoniae* II, *Pseudomonas putida*, and *Bacillus subtilis*. Furthermore, when testing enzymatic activity, it was found that the highest performance had phenol hydroxylase compared to both catechol 1, 2-dioxygenase and catechol 2,3-dioxygenase with the same enzyme activity [180]. The degradation pathway of phenol in the *Marinobacter* HA2 strain pathway of phenol degradation can be described via the 2,3 ortho-cleavage reaction, while the *Halomonas* HA1 strain involves the 1,2 phenol meta-cleavage mechanism. In bacteria, the major catabolic aerobic pathway of PCBs is an estradiol cleavage of the aromatic ring with the production of chlorobenzoate. Several genes, usually designated as bph, encode the key enzymes involved in this pathway. A mixed microbial population (7 bacteria and 1 fungus) was isolated from a soil sample of an unexposed natural environment. After enrichment and establishment in a defined medium containing p-CB (p-chlorobiphenyl) as the sole carbon source, this synthetic organic compound could be utilized. One of the microorganisms was engineered *P. putida* (pUC18) with the pUC18-nahH gene with catechol 2,3-dioxygenase (C23O), an enzyme able to eliminate phenanthrene and pyrene compounds

[181]. The catechol 2,3-dioxygenase (C23O) gene isolated from *Pseudomonas* sp. strain ND6 showed high similarity with the *Pseudomonas putida* G7 NAH 7 plasmid, 93% in amino acid sequence, and with the *Pseudomonas stutzeri* AN10 chromosome, 89% in amino acid sequence [182]. In the same strain, the C12O genes (*catA*, *catA_I*, *catA_{II}*, and *catA_{III}*) showed involvement in naphthalene catabolism, with the catechol 1,2-dioxygenase isoenzymes encoded by these genes [183]. Interestingly, some bacteria had no genetic basis for the naphthalene-degradative pathway. The main role in this process was played by primers for 1,2 and 2,3 CTD genes, encoding for catechol dioxygenase, bacterial iron-containing enzymes [183]. Some research indicates the involvement of *Pseudomonas* sp. dioxygenases enzymes in ortho (intradiol) cleavage and meta (extradiol) cleavage. Also, the catechol 1,2-dioxygenase (C12O) has been identified in *Achromobacter xylosoxidans* DN002, also involved in the ortho-cleavage pathway [184, 185]. The *Pseudomonas stutzeri* DJP1 strain degradation system consisted of PHE, seven metabolic intermediates, and PYR. The activity of key enzymes such as ring-cleaving dioxygenase, acetaldehyde dehydrogenase, and catechol 12 dioxygenase encoded by *PhdF*, *NidD*, and *CatA* genes, respectively, was recognized in cometabolic degradation and removal of high molecular weight polycyclic aromatic hydrocarbons (HMW-PAHs). Furthermore, *Pseudomonas* sp. plasmid encodes not only for catechol 1, 2-dioxygenase but also 3, 4-PCD with the same role, aromatic ring degradation [186]. Similar to lignin degradation, biphenyl-degrading bacteria can be useful and degrade xenobiotics. Located on MGE, horizontal gene transfer enables gene sharing and evolution, so these can be found in different biphenyl-degrading bacteria. Therefore, the Bph genes/degradation gene clusters for biphenyl/PCB are located on plasmids *Geobacillus* sp. strain JF8 pBt40, *Rhodococcus* sp. linear pRHL1, *Cupriavidus* sp. pSK-4, and *Pseudomonas putida* MB1335 [187]. Furthermore, biphenyl-degrading bacteria can be useful and degrade xenobiotics. The xenobiotics, PCBs with up to 10 chlorine atoms attached to two aromatic benzene rings, are highly hydrophobic and toxic compounds. Furthermore, *Cupriavidus basilensis* KF708 genome sequencing revealed 207 genes involved in the metabolism of aromatic compounds belonging to the bph gene cluster (bphEGFA1A2A3BCDA4) [188]. In general, bph genes encode dehydrogenases, namely bphA1A2A3A4, for the biphenyl dioxygenase, a multi-component enzyme. The bph genes beside the plasmid have been found on ICEs in *C. oxalaticus* A5 (formerly *Ralstonia* sp. strain A5) and *Acidovorax* sp. KKS102 [189]. *Pseudomonas* sp. strain shows adaptability, expressing biphenyl and salicylate metabolism gene clusters (the bph-sal element). Transfer of the bph-sal element in *Pseudomonas putida* KF715 enabled genome plasticity and genetic variability to withstand environmental challenges [190]. The role of biphenyl catabolic enzymes is to co-metabolize certain PCBs into chlorobenzoic acids and to substitute

chlorine. Another group of PCB genes is salicylate catabolic sal genes. In 9 of 10 biphenyl/PCB bacterial strains, bph and sal gene clusters were transferred by integrative conjugative elements (ICEs), named *ICEbph-sal*, in *Pseudomonas putida* KF715. The SalR protein of *Acinetobacter* sp. strain ADP1 encoded by *salA* is a salicylate hydroxylase activator, while the same SalR protein, the *Pseudomonas putida* S-1 SalR protein encoded by sal genes, regulates salicylate degradation. The Sal gene encodes for LysR-Type Transcriptional Regulator family proteins - SalR is involved in converting alkyl salicylates into catechol [191]. Some cyanobacteria are involved in organic pollutant degradation while exhibiting similar mechanisms. *Leptolyngbya* sp. can be very effective in phenol degradation, while *Spirulina* CPCC-695 degrades estrone using it as a carbon source [192, 193]. Besides point mutations and mobile genetic elements, metagenomic tandem repeats have shown potential in engineered organisms. A metagenomic tandem repeat sequence of catechol 2,3-dioxygenase (C23O)-encoding genes in genetically modified *E. coli* showed increased C23O enzyme activity [194]. A catechol 2,3-dioxygenase (C23O)-encoding gene sequence plays a crucial role in crude oil degradation. The most degrading activity had EAU29438 *Ewingella americana* 16S rRNA gene and partial sequence (CFfab 14) and *Bacillus megaterium* strain IV22 16S ribosomal RNA gene, partial sequence (CFfab15) [195].

Benefits and Drawbacks of the Introduction of GMOs/GEMs in Bioremediation

The bioremediation technique is an old/new approach to restoring the harmed environment. New methods include the introduction of genetically modified microorganisms that advance pollution elimination without forming partially degraded intermediates, which are more toxic than the prime pollutants, which is the opposite of conventional procedures. Biological methods are given an advantage because chemical and physical remediation approaches are expensive and have adverse effects on the environment. Thus, there has been a rise in the use of genetically modified/engineered microorganisms and biotechnology or genetic engineering. GMOs refer to all genetically changed living organisms. In contrast, the term "genetically engineered organisms" (GEMs) refers to microorganisms (bacteria, fungus, and yeast, among others) genetically modified by synthetic biology (removing and inserting genes) [38]. These designed organisms are tailored to have specific metabolic activities, the ability to overcome environmental stress, adjust to different abiotic and biotic factors, stronger structuring of genetic variation retained in the population, and the tracing of GEMs with molecular techniques. The enzyme 2,3-dihydroxy biphenyl 1,2-dioxygenase is encoded by genes bphC and etbC present in the *Rhodococcus* strain transferred from *R. corallines* to *Rhodococcus* TE1. Thus, the recombinant

Rhodococcus achieved complete degradation of atrazine to cyanuric acid [196]. Manipulation with genetic material brings environmental concern.

In nature, trait variation within bacteria happens through gene transfer (conjugation, transformation, and transduction). Laboratory-engineered microorganisms can, with target genes, introduce unwanted genes into the environment (antibiotic resistance genes). However, the risk can be avoided by introducing technical safeguards and innovation and applying laws and regulations that are characteristic of every region. The plasmids as cloning vectors can be substituted with mini-transposons to clone and insert target sequences with non-antibiotic resistance genes [197]. The natural ability of bacteria's toxin-antitoxin system to program their suicide under stress can also be activated after the depletion of pollutants. The interaction between microorganisms in environmental conditions is always uncertain; thus, field experiments must be conducted. Furthermore, remediation of the environmental damage caused by pollution and the potential threats of introducing GEMs can be leveled by a detailed risk assessment and constant monitoring after release. Also, the manipulation of genetic material must have boundaries by strict regulations, standards, and laws [198]. Thus, the introduction of rDNA into ecosystems is still restricted.

Other Aspects of Bioremediation without Using Biological Agents

The bioremediation technique that uses biological agents is not without any limitations. There are high molecular PAHs that cannot be broken down using a microorganism or incomplete degradation of pollutants, creating bigger treated, more toxic, and mobile intermediates. Also, unlike chemical and physical techniques, bioremediation takes a lot of time to remove toxins. Remediation with superoxidation products is based on known Fenton and Haber-Weiss reactions. The EU-funded HAC project targets the application of this advanced technique to clean water and soil from a wide range of contaminants such as PAHs, PBCs, CHCs, benzene, toluene, petroleum hydrocarbons, crude oil, and pesticides [199].

PAH degradation by chemical oxidation was shown to be very successful, namely modified Fenton's reagent, hydrogen peroxide, and potassium permanganate degradation up to 95% light PAHs up to 1600 mg/kgSS, heavy PAHs up to 1200 mg/kgSS, residues were below 100 mg/kgSS. [200]. Some other studies compared benzo[a]pyrene (BaP) degradation by potassium permanganate (KMnO_4), Fe^{2+} + sodium persulfate ($\text{Fe}^{2+} + \text{PS}$), Fenton's reagent ($\text{Fe}^{2+} + \text{H}_2\text{O}_2$), and hydrogen peroxide (H_2O_2). The higher removal efficiency showed KMnO_4 [201]. Oxidation with persulfate (PS) was less effective (up to 43%) than with permanganate (PM) (up to 98%), while treatment with modified Fenton's reagent (F) showed the lowest degradation yields [202].

Stabilization/solidification (S/S) is considered the most appropriate method of converting contaminants into their least soluble/toxic. This is a physical and chemical reaction during which insoluble metal hydroxides are formed. Carbon binders are applied to chemically change and mechanically encapsulate the material. The calcined clay (CC) and limestone (LS) showed good Zn, As, and Pb immobilization results. Additionally, limestone-calcined clay cement (LC3) is also an effective Zn and Pb binder. Further, the reaction between LS and CC results in primary and secondary hydration products such as calcium silicate hydrate, calcium hydrate, calcium aluminate silicate hydrate, and calcium aluminate hydrate [203]. Geopolymer binders are considered eco-friendly and safer than conventional ones. Alkaline-activated silica-rich wastes, 3 aluminum silicate hydrates, showed binding capacity with 20% fly ash, 15% silica fume, and rice husk ash, all three with 6.5 molar of NaOH [204].

Advanced oxidation processes include chemical processes (Fenton's process and Peroxone process), photochemical (photodecomposition (H_2O_2 /UV/ O_3 /UV), photocatalysis (TiO_2 , ZnO), Photo- Fenton process ($\text{H}_2\text{O}_2/\text{Fe}^{2+}$ /UV), electrochemical (Electro-Fenton process) [205]. POP degradation using the TiO_2 /UV photocatalytic system in wastewater was efficient. Namely, the TiO_2 /UV photocatalytic system effectively degraded pollutants with constant hydrogen production. In addition, toxic compounds from textiles, such as suspended solids, toxic chemicals, azo dyes, color, etc., can be degraded by adding strong oxidants such as hydroxyl radical ($\cdot\text{OH}$) [206]. The introduction of novel materials that can photocatalytically degrade malachite green (MG) under the UV or VIS has promising results. Zinc oxide nanocomposites impregnated with nickel (Ni/ZnO) were used in malachite green (MG) degradation. The highest removal efficiency of 99.4% was detected at pH 8 under VIS light [207].

Nanomaterials (nanomembranes, nanotubes, and nanoparticles) have wide use in drinking water purification, being able to remove toxic heavy metals such as Ni(II), Pb(II), Hg(II), Cu(II), and Zn(II) and POPs such as antibiotics, pesticides, and biological toxic components such as viruses, organic materials, bacteria, micronutrients, and algae [208]. Immobilization of pollutants can be achieved using carbon-based nanomaterials. Engineered nanomaterials carbon nanotube (CNT)- Fe_xO_y composites and (CFs) carbon nanotubes absorb and immobilize Cd and As. Also, Fe nanoparticles showed 88.34% successful Total Petroleum Hydrocarbons (TPHs) removal capacity from water. NM-based modifications act via reductive reactions, affecting the mobility of pollutants. Sodium carboxymethyl cellulose-stabilized nZVI could remove about 80% of Cr (VI) from soil. However, using these techniques has side effects, such as photocatalysis, the high level of mineralization, toxic postreaction products, photocorrosion, and light irradiation, which cause photocatalysts to become unstable; advanced oxidation

processes (AOP) change pollutants, affect oxidants and catalysts, change configuration and form, and release toxic byproducts; the Fenton process applied at large treatments mishandles hydrogen peroxide [209]. Nitrogen and ammonium nitrogen in wastewater can be removed by algal and bacterial consortiums [210]. The problem of drinking water can be resolved through electro-bioremediation. Research funded by the European Union's Horizon 2020 project ELECTRA within the EU Bioremediation cluster NYMPHE project showed high nitrate removal and chlorine production at the same time, eliminating pathogens such as *T. coliforms*, *E. coli*, and *Enterococcus* from drinking water. The project was conducted per the European Directive 2020/2184 [211]. Also, under this project, research conducted in a real environment showed that electro-bioremediation associated with Pseudomonadaceae, Rhizobiaceae, Gallionellaceae, and Xanthomonadaceae completely reduced NO_3^- to N_2 [212]. Therefore, further research in this direction is needed.

Conclusions

Soil remediation methods may be applied in situ, on-site, and off-site. Within R&I, on-site methods in situ occur directly at the field, leaving the soil intact and clean. Therefore, on-site applied green techniques are within biological methods, bioremediation, and phytoremediation, or their combination. For instance, bacteria like *Bacillus*, *Enterobacter*, *Escherichia*, and *Pseudomonas* help remediate metal-contaminated soil.

Water pollution can be addressed by the algal-bacteria consortium that applies to the bioremediation of wastewater. This application is climate-friendly because the algae produce oxygen for the bacteria while using carbon dioxide in photosynthesis [213].

According to this review, the implementation of successful bioremediation requires a functional genomics approach. Being capable of adjusting to any environment, microorganisms' tolerance detoxifies and removes the toxic metals. Mechanisms developed include different efflux transporter proteins. This review presents a variety of proteins, including P-type ATPases, RND (resistance, nodulation, cell division), CDF (cation diffusion facilitator), ATP binding cassette (ABC) transporters, MFS family proteins, and more, encoded by revealed and new genes and operons [214]. Therefore, many different biodegradation pathways have been discovered; however, new genes, their relationships, and their positions in the genome are receiving increasing attention. In addition to native genes, genetic engineering enabled various microorganisms to remove highly toxic substances from the environment. Numerous chemical, physical, and biological factors influence the course and success of bioremediation: pH, temperature, water content, nutrient availability, redox potential, and concentration of pollutants. Namely, many microorganisms synthesize enzymes

only in the presence of a high concentration of toxic substances. Therefore, significance is in the novel gene sequence that will enable removal at low pollutant concentrations. Throughout this review, many native and engineered microorganisms showed a high ability to deal with both inorganic and organic pollution. Moreover, the same bacterial strain genome exhibits HMs and POPs removal potential. For instance, *Pseudomonas* sp., *Cupriavidus* sp., *Burkholderia* sp., and fungi *Rhodococcus* sp. were shown to be useful in the removal of a wide variety of pollutants. In general, microorganisms are handy tools for removing pollution, and offering insight into the mechanism of their action is essential for all researchers. Lab-scale biodegradation studies to field-scale trial insights can be conducted and evaluated by applying Molecular Biological Tools (MBTs) such as qPCR and 16S gene amplicon rRNA sequencing. Thus, the site abundance of functional genes can be detected. In enhanced bioremediation field trials, microbial consortia linked to PHC and TCE biodegradation include *Dechloromonas*, *Deinococcus*, *Clostridium*, *Geobacillus*, *Syntrophus*, *Enterobacter*, as well as *Methanobacteria* and *Methanomicrobia* that degrade HC to CH_4 [215]. Furthermore, *in situ* bioremediation reduced TPHs in soil by about 88%, introducing *Bacillus* sp., *Pseudomonas* sp., *Arthrobacter* sp., *Acinetobacter* sp., *Alcaligenes* sp., and *Brevibacillus* sp. Some other strains, *Acinetobacter baumannii* (S19, S26, S30), *Burkholderia cepacia* (P20), and *Pseudomonas* sp. (S24), also showed the *in situ* potential of 90% in TPH removal. Further, *Bacillus*, *B. thuringiensis* B3, and *B. cereus* B6 reduced diesel and crude oil from crude oil-polluted fields between 28% and 84% [216]. In field trials for removing heavy metals from water and soil, *Bacillus* sp. and *Pseudomonas* sp. applications have shown promising results. Furthermore, *Geobacteraceae* species were found to be the most abundant in reducing Fe^{3+} . Intrinsic bioremediation, bioaugmentation, and engineered bioremediation help restore polluted soil. However, from lab to field applications, there need to be considerations for monitoring after the release of GEMs and factors such as environmental challenges, including temperature, humidity, pH, rather than the competition and survival of microorganisms [216, 217]. The findings of these studies imply that microorganisms are suitable candidates for the bio-removal of a wide range of inorganic and organic pollutants, as shown in Table 1. and Table 2. Research results are vital in implementing preventive and protective actions. The implication of this research is to identify genetic determinants of prime bioremediation as an advantage in target pollutant removal and to direct researchers toward genetically modified organisms of interest with respect to potential drawbacks. Therefore, this research is relevant for researchers and stakeholders participating in living labs and lighthouses as open innovation platforms for the transdisciplinary conservation, environmental, and agricultural sustainability fields.

Further Perspectives

The greatest worry in recent years is human-induced climate change. As a consequence, contaminant release, pathways, and bioavailability in terrestrial and aquatic environments are changed [218]. To tackle this issue, future bioremediation focus must be shifted to genetic microbial community modeling, pollutant-microbe interaction modeling, and site-specific pollution forecasting. The biggest problem in successful bioremediation is environmental changes that occur faster than wanted. Therefore, predictive ecology must tailor effective microbial consortia.

Future research should focus on forecasting models based on improved genomic techniques and developing precision-designed synthetic microbial communities that won't be affected by environmental factors. Consequently, climate reshapes the landscape and influences environmental factors and biocenosis; therefore, advancing genetic encoding of pollutant pathways via DNA nanotechnology can give precise pollutant-microbe interaction patterns in changed ecosystems.

The limiting factor is the concentration of the contaminants in the environment. Toxins in high and low concentrations directly affect microbial activity. Thus, lower pollution concentration decreases degrading enzyme production by microorganisms. Therefore, the decomposition rate of catabolic enzymes can be increased by genetic engineering methods, increasing the yield of enzyme production. For instance, organophosphate degrading enzymes were overexpressed in recombinant *E. coli* BL21, increasing the pesticide degrading rate compared with non-recombinant enzymes [219]. The aspects of assisted evolution also should be considered as a new direction.

Conflict of Interest

The authors declare no conflict of interest.

References

1. EUROPEAN COMMISSION. Environment for Europeans. No 65, March 2018. Available online: <https://op.europa.eu/en/publication-detail/-/publication/d499cc57-4768-11e8-beld-01aa75ed71a1> (accessed on 22.8.2024). **2018**.
2. RAVEN P., WACKERNAGEL M. Maintaining biodiversity will define our long-term success. *Plant Diversity*. **42** (4), 211, **2020**.
3. EDO G.I., SAMUEL P.O., OLONI G.O., EZEKIEL G.O., IKPEKORO V.O., OBASOHAN P., AGBO J.J. Environmental persistence, bioaccumulation, and ecotoxicology of heavy metals. *Chemistry and Ecology*. **40** (3), 322, **2024**.
4. SALES DA SILVA I.G., GOMES DE ALMEIDA F.C., PADILHA DA ROCHA E SILVA N.M., CASAZZA A.A., CONVERTI A., ASFORA SARUBBO L. Soil bioremediation: Overview of technologies and trends. *Energies*. **13** (18), 4664, **2020**.
5. SAHA A., MUKHERJEE P., ROY K., SEN K., SANYAL T. A review on phyto-remediation by aquatic macrophytes: a natural promising tool for sustainable management of ecosystem. *International Journal of Experimental Research and Review*. **27**, 9, **2022**.
6. NIKOLIĆ M., STEVOVIĆ S. Family Asteraceae as a sustainable planning tool in phytoremediation and its relevance in urban areas. *Urban Forestry & Urban Greening*. **14** (4), 782, **2015**.
7. NIKOLIĆ M., TOMAŠEVIĆ V. Implication of the plant species belonging to the brassicaceae family in the metabolization of heavy metal pollutants in urban settings. *Polish Journal of Environmental Studies*. **30** (1), 523, **2020**.
8. NIKOLIĆ M., TOMASEVIĆ V., UGRINOV D. Energy plants as biofuel source and as accumulators of heavy metals. *Hemijaska Industrija*. **76** (4), 209, **2022**.
9. YAASHIKAA P.R., KUMARP.S. Bioremediation of hazardous pollutants from agricultural soils: A sustainable approach for waste management towards urban sustainability. *Environmental Pollution*. **312**, 120031, **2022**.
10. SHACKIRA A.M. JAZEEL K. PUTHUR J.T. Phycoremediation and phytoremediation: Promising tools of green remediation. In: Sustainable environmental clean-up. 1st ed.; Mishra, K.V, Kumar A. (eds). Elsevier, Amsterdam, pp. 273, **2021**.
11. WU P., ZHANG Z., LUO Y., BAI Y., FAN J. "Bioremediation of phenolic pollutants by algae-current status and challenges." *Bioresource Technology*. **350**, 126930, **2022**.
12. SINGH A., PAL D.B., KUMAR S., SRIVASTVA N., SYED A., ELGORBAN A.M., SINGH R., GUPTA V.K. Studies on Zero-cost algae based phytoremediation of dye and heavy metal from simulated wastewater. *Bioresource Technology*. **342**, 125971, **2021**.
13. DINAKARKUMAR Y., GNANASEKARAN R., REDDY G.K., VASU V., BALAMURUGAN P., MURALI G. Fungal bioremediation: An overview of the mechanisms, applications and future perspectives. *Environmental Chemistry and Ecotoxicology*. **6**, 293, **2024**.
14. GU J.D. On environmental biotechnology of bioremediation. *Applied Environmental Biotechnology*. **5**, 3, **2021**.
15. SHARMA B., DANGI A.K., SHUKLA P. Contemporary enzyme based technologies for bioremediation: a review. *Journal of Environmental Management*. **210**, 10, **2018**.
16. MILIĆ J., AVDALOVIĆ J., KNUDSEN T.Š. Microbial bioremediation of the oil polluted environment and the sustainable development goals of pillar Planet of the Agenda 2030. *Environment, Development and Sustainability*. **23**, **2024**.
17. PURVIS B., MAO Y., ROBINSON D. Three pillars of sustainability: in search of conceptual origins. *Sustainability science*. **14**, 681, **2019**.
18. Europe's 2030 climate and energy targets Available online: <https://op.europa.eu/en/publication-detail/-/publication/1c8ab88a-e44d-11eb-895a-01aa75ed71a1/language-en/format-PDF/source-219125747> (accessed on 11.07.2024). **2024**.
19. The European Green Deal Available online: https://commission.europa.eu/strategy-and-policy/priorities-2019-2024/european-green-deal_en (accessed on 11.07.2024). **2024**.
20. The Horizon Europe, Available online: <https://research-and-innovation.ec.europa.eu/funding/funding->

- opportunities/funding-programmes-and-open-calls/horizon-europe_en (accessed on 11.07.2024).
21. Horizon Europe Work programme (2023-24) - Cluster 6. Available online: https://ec.europa.eu/info/funding-tenders/opportunities/docs/2021-2027/horizon/wp-call/2023-2024/wp-9-food-bioeconomy-natural-resources-agriculture-and-environment_horizon-2023-2024_en.pdf (accessed on 11.07.2024). **2024**.
 22. Biosysmo. Available online: <https://www.biosysmo.eu/> (accessed on 25.12.2024). **2024**.
 23. DVOŘÁK P., GALVÃO T.C., PFLÜGER-GRAU K., BANKS A.M., DE LORENZO V., JIMÉNEZ J.I. Water potential governs the effector specificity of the transcriptional regulator XylR of *Pseudomonas putida*. *Environmental Microbiology*. **25** (5), 1041, **2023**.
 24. AVENDAÑO R., MUÑOZ-MONTERO S., ROJAS-GÄTJENS D., FUENTES-SCHWEIZER P., VIETO S., MONTENEGRO R., SALVADOR M., FREW R., KIM J., CHAVARRÍA M., JIMÉNEZ J.I. Production of selenium nanoparticles occurs through an interconnected pathway of sulphur metabolism and oxidative stress response in *Pseudomonas putida* KT2440. *Microbial Biotechnology*. **16** (5), 931, **2023**.
 25. GREENER. Available online: <https://www.greener-h2020.eu/en/normal/home> (accessed on 29.12.2024). **2024**.
 26. MARTÍNEZ-CUESTA R., CONLON R., WANG M., BLANCO-ROMERO E., DURÁN D., REDONDO-NIETO M., DOWLING D., GARRIDO-SANZ D., MARTIN M., GERMAINE K., RIVILLA R. Field scale biodegradation of total petroleum hydrocarbons and soil restoration by Ecopiles: microbiological analysis of the process. *Frontiers in Microbiology*. **14**, 1158130, **2023**.
 27. VERDEL N., RIJAVEC T., RYBKIN I., ERZIN A., VELIŠČEK Ž., PINTAR A., LAPANJE A. Isolation, Identification, and Selection of Bacteria With Proof-of-Concept for Bioaugmentation of Whitewater From Wood-Free Paper Mills. *Frontiers in Microbiology*. **12**, 758702, **2021**.
 28. GARRIDO-SANZ D., REDONDO-NIETO M., MARTÍN M., RIVILLA R. Comparative genomics of the *Rhodococcus* genus shows wide distribution of biodegradation traits. *Microorganisms*. **8** (5), 774, **2020**.
 29. ARRIDO-SANZ D., SANSEGUNDO-LOBATO P., REDONDO-NIETO M., SUMAN J., CAJTHAML T., BLANCO-ROMERO E., MARTIN M., UHLIK O., RIVILLA R. Analysis of the biodegradative and adaptive potential of the novel polychlorinated biphenyl degrader *Rhodococcus* sp. WAY2 revealed by its complete genome sequence. *Microbial genomics*. **6** (4), 000363, **2020**.
 30. Mibirem. Available online: <https://www.mibirem.eu/approach/> (accessed on 29.12.2024). **2024**.
 31. ZAGHLOUL A., SABER M., GADOW S., AWAD F. Biological indicators for pollution detection in terrestrial and aquatic ecosystems. *Bulletin of the National Research Centre*. **44**, 1, **2020**.
 32. KUPPAN N., PADMAN M., MAHADEVA M., SRINIVASAN S., DEVARAJAN R. A comprehensive review of sustainable bioremediation techniques: Eco friendly solutions for waste and pollution management. *Waste Management Bulletin*. **2** (30), 154, **2024**.
 33. SALES DA SILVA I.G., GOMES DE ALMEIDA F.C., PADILHA DA ROCHA E SILVA N.M., CASAZZA A.A., CONVERTI A., ASFORA SARUBBO L. Soil bioremediation: Overview of technologies and trends. *Energies*. **13**(18), 4664, **2020**.
 34. LACALLE R.G., BECERRIL J.M., GARBISU C. Biological methods of polluted soil remediation for an effective economically-optimal recovery of soil health and ecosystem services. *Journal of Environmental Science and Public Health*. **4** (2), 112, **2020**.
 35. KUMAR V.V. Microbial Remediation: A Natural Approach for Environmental Pollution Management. In: *Mycoremediation and Environmental Sustainability. Fungal Biology*. 1st ed.; Prasad R., Nayak S.C., Kharwar R.N., Dubey N.K. (eds) Springer, Cham. **3**, 171, **2021**.
 36. JOSHI S.S., JADHAV M.P.K., CHOURASIA N.K., JADHAV M.P.J., PATHADE K.N. Bioremediation Strategies for Soil And Water Pollution Harnessing The Power Of Microorganisms. *Migration Letters*. **2** (S8), 1, **2024**.
 37. PERERA I.C., HEMAMALI E.H. Genetically modified organisms for bioremediation: Current research and advancements. *Bioremediation of Environmental Pollutants: Emerging Trends and Strategies*. 163, **2022**.
 38. RAFEEQ H., AFSHEEN N., RAFIQUE S., ARSHAD A., INTISAR M., HUSSAIN A., BILAL M., IQBAL H.M. Genetically engineered microorganisms for environmental remediation. *Chemosphere*. **310**, 136751, **2023**.
 39. SAXENA G., KISHOR R., SARATALE G.D., BHARAGAVA R.N. Genetically Modified Organisms (GMOs) and Their Potential in Environmental Management: Constraints, Prospects, and Challenges. In: *Bioremediation of Industrial Waste for Environmental Safety*. 1st ed.; Bharagava, R., Saxena, G. (eds), Springer, Singapore. **2**, 1, **2020**.
 40. SHARMA P., BANO A., SINGH S.P., SHARMA S., XIA C., NADDA A.K., LAM S.S., TONG Y.W. Engineered microbes as effective tools for the remediation of polycyclic aromatic hydrocarbons and heavy metals. *Chemosphere*. **306**, 135538, **2022**.
 41. EUROPEAN COMMISSION. Cost-effective solutions to saline wastewater. Available online: <https://projects.research-and-innovation.ec.europa.eu/en/projects/success-stories/all/cost-effective-solutions-saline-waste-water> (accessed on 03.09.2024). **2018**.
 42. EU Mission: A Soil Deal for Europe. Available online: https://research-and-innovation.ec.europa.eu/funding/funding-opportunities/funding-programmes-and-open-calls/horizon-europe/eu-missions-horizon-europe/soil-deal-europe_en (accessed on 03.09.2024). **2024**.
 43. EEU Mission: Restore our Ocean and Waters. Available online: https://research-and-innovation.ec.europa.eu/funding/funding-opportunities/funding-programmes-and-open-calls/horizon-europe/eu-missions-horizon-europe/restore-our-ocean-and-waters_en (17.09.2024). **2024**.
 44. XNATURA. Bioremediation as an effective tool for a clean environment, Available online: <https://circulareconomy.europa.eu/platform/en/news-and-events/all-events/bioremediation-effective-tool-clean-environment> (17.09.2024). **2024**.
 45. SANTIAGO E., MORENO D.F., ACAR M. Phenotypic plasticity as a facilitator of microbial evolution. *Environmental Epigenetics*. **8** (1), 1, **2022**.
 46. HUSSAIN S., KHAN M., SHEIKH T.M.M., MUMTAZ M.Z., CHOCHAN T.A., SHAMIM S., LIU Y. Zinc essentiality, toxicity, and its bacterial bioremediation: A comprehensive insight. *Frontiers in Microbiology*. **13**, 900740, **2022**.
 47. ELYAMINE A.M., KAN J., MENG S., TAO P., WANG H., HU Z. Aerobic and anaerobic bacterial and fungal degradation of pyrene: mechanism pathway including

- biochemical reaction and catabolic genes. *International Journal of Molecular Sciences*. **22** (15), 8202, **2021**.
48. SHAKERIAN F., ZHAO J., LI S.P. Recent development in the application of immobilized oxidative enzymes for bioremediation of hazardous micropollutants—A review. *Chemosphere*. **239**, 124716, **2020**.
 49. ONTAÑÓN O.M., FERNANDEZ M., AGOSTINI E., GONZÁLEZ P.S. Identification of the main mechanisms involved in the tolerance and bioremediation of Cr (VI) by *Bacillus* sp. SFC 500-1E. *Environmental Science and Pollution Research*. **25**, 16111, **2018**.
 50. SANTERO E., DÍAZ E. Genetics of biodegradation and bioremediation. *Genes*. **11** (4), 441, **2020**.
 51. NAG A. Understanding the Environment and Sustainability with Molecular Approaches. In: *Biotechnological Innovations for Environmental Bioremediation*. 1st ed; Arora S., Kumar A., Ogita S., Yau Y.Y. (eds), Springer, Singapore, pp 895, **2022**.
 52. MALIK G., ARORA R., CHATURVEDI R., PAUL M.S. Implementation of genetic engineering and novel omics approaches to enhance bioremediation: a focused review. *Bulletin of Environmental Contamination and Toxicology*. **108**, 443, **2021**.
 53. SHEKHAR S.K., GODHEJA J., MODI D.R. Molecular technologies for assessment of bioremediation and characterization of microbial communities at pollutant-contaminated sites. In *Bioremediation of Industrial Waste for Environmental Safety: Volume II: Biological Agents and Methods for Industrial Waste Management*, 1st ed; Bharagava R., Gaurav Saxena G. (eds), Springer, Singapore, pp 437, **2020**.
 54. DANOUCHE M., EL GHACHTOULI N., EL ARROUSSI H. Phycoremediation mechanisms of heavy metals using living green microalgae: physicochemical and molecular approaches for enhancing selectivity and removal capacity. *Heliyon*. **7** (7), e07609, **2021**.
 55. LIU S.H., ZENG G.M., NIU Q.Y., LIU Y., ZHOU L., JIANG L.H., CHENG M. Bioremediation mechanisms of combined pollution of PAHs and heavy metals by bacteria and fungi: A mini review. *Bioresource Technology*. **224**, 25, **2017**.
 56. KAUR S., ROY A. Bioremediation of heavy metals from wastewater using nanomaterials. *Environment, Development and Sustainability*. **23** (7), 9617, **2021**.
 57. BIELICKA A., BOJANOWSKA I., WISNIEWSKI A. Two Faces of Chromium-Pollutant and Bioelement. *Polish Journal of Environmental Studies*. **14** (1), 5, **2005**.
 58. GUO C., MA X., GAO F., GUO Y. Off-target effects in CRISPR/Cas9 gene editing. *Frontiers in Bioengineering and Biotechnology*. **11**, 1143157, **2023**.
 59. JAISWAL S., SINGH D.K., SHUKLA P. Gene editing and systems biology tools for pesticide bioremediation: a review. *Frontiers in Microbiology*. **10**, 87, **2019**.
 60. CHAUDHARI M., SOLANKI K. Role of Crisper-Cas Technique in Bioremediation of Pesticides. *International Journal of Current Microbiology and Applied Sciences*. **12** (08), 82, **2023**.
 61. PICKAR-OLIVER A., GERSBACH C.A. The next generation of CRISPR–Cas technologies and applications. *Nature Reviews Molecular Cell Biology*. **20** (8), **2019**.
 62. JIN Y., LUAN Y., NING Y., WANG L.J. Effects and mechanisms of microbial remediation of heavy metals in soil: a critical review. *Applied Sciences*. **8** (8), 1336, **2018**.
 63. FAN J., OKYAY T.O., RODRIGUES D.F. The synergism of temperature, pH and growth phases on heavy metal biosorption by two environmental isolates. *Journal of Hazardous Materials*. **279**, 236, **2014**.
 64. SHAN B., HAO R., ZHANG J., LI J., YE Y., LU A. Microbial remediation mechanisms and applications for lead-contaminated environments. *World Journal of Microbiology and Biotechnology*. **39** (2), 38, **2023**.
 65. BANDARA P.C., PEÑA-BAHAMONDE J., RODRIGUES D.F. Redox mechanisms of conversion of Cr (VI) to Cr (III) by graphene oxide-polymer composite. *Scientific Reports*. **10** (1), 9237, **2020**.
 66. SURESH G., BALASUBRAMANIAN B., RAVICHANDRAN N., RAMESH B., KAMYAB H., VELMURUGAN P., RAVI A.V. Bioremediation of hexavalent chromium-contaminated wastewater by *Bacillus thuringiensis* and *Staphylococcus capitis* isolated from tannery sediment. *Biomass Conversion and Biorefinery*. **11**, 383, **2021**.
 67. KHANAM R., AL ASHIK S.A., SURIEA U., MAHMUD S. Isolation of chromium resistant bacteria from tannery waste and assessment of their chromium reducing capabilities—A Bioremediation Approach. *Heliyon*. **10** (6), 2405, **2024**.
 68. CHAI L., DING C., TANG C., YANG W., YANG Z., WANG Y., LIAO Q., LI J. Discerning three novel chromate reduce and transport genes of highly efficient *Pannonibacter phragmitetus* BB: from genome to gene and protein. *Ecotoxicology and Environmental Safety*. **162**, 139, **2018**.
 69. HUANG Y., FENG H., LU H., ZENG Y. A thorough survey for Cr-resistant and/or -reducing bacteria identified comprehensive and pivotal taxa. *International Biodeterioration and Biodegradation*. **117**, 22, **2017**.
 70. KARIM M.E., SHARMIN S.A., MONIRUZZAMAN M., FARDOUS Z., DAS K.C., BANIK S., SALIMULLAH M. Biotransformation of chromium (VI) by *Bacillus* sp. isolated from chromate contaminated landfill site. *Chemistry and Ecology*. **36** (10), 922, **2020**.
 71. GU R., GAO J., DONG L., LIU Y., LI X., BAI Q., XIAO, H. Chromium metabolism characteristics of coexpression of *ChrA* and *ChrT* gene. *Ecotoxicology and Environmental Safety*. **204**, 111060, **2020**.
 72. PUSHKAR B., SEVAK P., PARAB S., NILKANTH N. Chromium pollution and its bioremediation mechanisms in bacteria: A review. *Journal of Environmental Management*. **287**, 112279, **2021**.
 73. RAMLI N.N., OTHMAN A.R., KURNIAWAN S.B., ABDULLAH S.R.S., HASAN H.A. Metabolic pathway of Cr (VI) reduction by bacteria: a review. *Microbiological Research*. **268**, 127288, **2023**.
 74. VITI C., MARCHI E., DECOROSI F., GIOVANNETTI L. Molecular mechanisms of Cr (VI) resistance in bacteria and fungi. *FEMS Microbiology Reviews*. **38** (4), 633, **2014**.
 75. BALDIRIS R., ACOSTA-TAPIA N., MONTES A., HERNÁNDEZ J., VIVAS-REYES R. Reduction of hexavalent chromium and detection of chromate reductase (*ChrR*) in *Stenotrophomonas maltophilia*. *Molecules*. **23** (2), 406, **2018**.
 76. ZHAO K., ZHANG W., LIANG Z., ZHAO H., CHAI J., YANG Y., ZHANG D. Facilitating New Chromium Reducing Microbes to Enhance Hexavalent Chromium Reduction by In Situ Sonoporation-Mediated Gene Transfer in Soils. *Environmental Science & Technology*. **57** (40), 15123, **2023**.
 77. SHI L., LIU B., ZHANG X., BU Y., SHEN Z., ZOU J., CHEN Y. Cloning of nitrate reductase and nitrite reductase

- genes and their functional analysis in regulating Cr (VI) reduction in ectomycorrhizal fungus *Pisolithus* sp. 1. *Frontiers in Microbiology*. **13**, 926748, **2022**.
78. PRADHAN S.K., SINGH N.R. RATH B.P., THATOI H. Bacterial chromate reduction: a review of important genomic, proteomic, and bioinformatic analysis. *Critical Reviews in Environmental Science and Technology*. **46**, 1659, **2016**.
 79. JUHNKE S., PEITZSCH N., HÜBENER N., GROSSE C., NIES D.H. New genes involved in chromate resistance in *Ralstonia metallidurans* strain CH34. *Archives of microbiology*. **179**, 15, **2002**.
 80. BAAZIZ H., GAMBARI C., BOYELDIEU A., ALI CHAOUCHE A., ALATOU R., MÉJEAN V., FONS M. ChrASO, the chromate efflux pump of *Shewanella oneidensis*, improves chromate survival and reduction. *PLoS One*. **12** (11), e0188516, **2017**.
 81. DAS P., SINHA S., MUKHERJEE S.K. Nickel bioremediation potential of *Bacillus thuringiensis* KUNil and some environmental factors in nickel removal. *Bioremediation Journal*. **18** (2), 169, **2014**.
 82. NAVARRO C., WU L.F., MANDRAND-BERTHELOT M.A. The nik operon of *Escherichia coli* encodes a periplasmic binding-protein-dependent transport system for nickel. *Molecular Microbiology*. **9** (6), 1181, **1993**.
 83. CHIVERS P.T., BENANTI E.L., HEIL-CHAPDELAINE V., IWIG J.S., ROWE J.L. Identification of Ni-(L-His) 2 as a substrate for NikABCDE-dependent nickel uptake in *Escherichia coli*. *Metallomics*. **4** (10), 1043, **2012**.
 84. LEBRETTE H., IANNELLO M., FONTECILLA-CAMPS J.C., CAVAZZA C. The binding mode of Ni-(L-His) 2 in NikA revealed by X-ray crystallography. *Journal of Inorganic Biochemistry*. **121**, 16, **2013**.
 85. DE PINA K., DESJARDIN V., MANDRAND-BERTHELOT M.A., GIORDANO G., WU L.F. Isolation and characterization of the nikR gene encoding a nickel-responsive regulator in *Escherichia coli*. *Journal of Bacteriology*. **181** (2), 670, **1999**.
 86. ERNST F.D., KUIPERS E.J., HEIJENS A., SARWARI R., STOOF J., PENN C.W., VAN VLIET A.H. The nickel-responsive regulator NikR controls activation and repression of gene transcription in *Helicobacter pylori*. *Infection and Immunity*. **73** (11), 7252, **2005**.
 87. BABAR Z., KHAN M., CHOTANA G.A., MURTAZA G., SHAMIM S. Evaluation of the potential role of *Bacillus altitudinis* MT422188 in nickel bioremediation from contaminated industrial effluents. *Sustainability*. **13** (13), 7353, **2021**.
 88. MAITRA S. Study of genetic determinants of nickel and cadmium resistance in bacteria-a review. *International Journal of Current Microbiology and Applied Sciences*. **5** (11), 459, **2016**.
 89. NARAYANASAMY S., THANKAPPAN S., KUMARAVEL S., RAGUPATHI S., UTHANDI S. Complete genome sequence analysis of a plant growth-promoting phyloplane *Bacillus altitudinis* FD48 offers mechanistic insights into priming drought stress tolerance in rice. *Genomics*. **115** (1), 110550, **2023**.
 90. MANDAL D., DAS S.K., ADHIKARI J., CHATTERJEE D., BANDYOPADHYAY T.K., BASU A. Genome sequencing, annotation and application of a strain of *Microbacterium paraoxydans*—a heavy metal hypertolerant and plant growth promoting bacterium. *Research Square*. **1**, **2024**.
 91. DIEP P., LEO SHEN H., WIESNER J.A., MYKYTCZUK N., PAPANGELAKIS V., YAKUNIN A.F., MAHADEVAN R. Engineered nickel bioaccumulation in *Escherichia coli* by NikABCDE transporter and metallothionein overexpression. *Engineering in Life Sciences*. **23** (7), 2200133, **2023**.
 92. LEE S., KHANAL A., CHO A.H., LEE H., KANG M.S., UNNO T., LEE J.H. Cupriavidus sp. strain Ni-2 resistant to high concentration of nickel and its genes responsible for the tolerance by genome comparison. *Archives of Microbiology*. **201**, 1323, **2019**.
 93. SHARMA P., PANDEY A.K., KIM S.H., SINGH S.P., CHATURVEDI P., VARJANI S. Critical review on microbial community during in-situ bioremediation of heavy metals from industrial wastewater. *Environmental Technology & Innovation*. **24**, 101826, **2021**.
 94. NAZARI M.T., SIMON V., MACHADO B.S., CRESTANI L., MARCHEZI G., CONCOLATO G., PICCIN J.S. *Rhodococcus*: A promising genus of actinomycetes for the bioremediation of organic and inorganic contaminants. *Journal of Environmental Management*. **323**, 116220, **2022**.
 95. HU Y.M., ZHOU J., DU B.Y., LIU H.L., ZHANG W.T., LIANG J.N. Health risks to local residents from the exposure of heavy metals around the largest copper smelter in China. *Ecotoxicology and Environmental Safety*. **171**, 329, **2019**.
 96. TIQUIA-ARASHIRO S.M. Lead absorption mechanisms in bacteria as strategies for lead bioremediation. *Applied Microbiology and Biotechnology*. **102** (13), 5437, **2018**.
 97. IGIRI B.E., OKODUWA S.I., IDOKO G.O., AKABUOGU E.P., ADEYI A.O., EJIOGU I.K. Toxicity and bioremediation of heavy metals contaminated ecosystem from tannery wastewater: a review. *Journal of Toxicology*. **1**, **2018**.
 98. MITRA A., CHATTERJEE S., KATAKI S., RASTOGI R.P., GUPTA D.K. Bacterial tolerance strategies against lead toxicity and their relevance in bioremediation application. *Environmental Science and Pollution Research*. **28**, 14271, **2021**.
 99. HUI C.Y., MA B.C., WANG Y.Q., YANG X.Q., CAI J.M. Designed bacteria based on natural pbr operons for detecting and detoxifying environmental lead: A mini-review. *Ecotoxicology and Environmental Safety*. **267**, 115662, **2023**.
 100. IBÁÑEZ M.M., CHECA S.K., SONCINI F.C. A single serine residue determines selectivity to monovalent metal ions in metalloregulators of the MerR family. *Journal of Bacteriology*. **197** (9), 1606, **2015**.
 101. SEXTON R., GILL P.R., DOWLING D.N., O'GARA F. Transcriptional regulation of the iron-responsive sigma factor gene pbrA. *Molecular and General Genetics MGG*. **250**, 50, **1996**.
 102. AMBROSI C., LEONI L., PUTIGNANI L., ORSI N., VISCA P. Pseudobactin biogenesis in the plant growth-promoting rhizobacterium *Pseudomonas* strain B10: identification and functional analysis of the L-ornithine N5-oxygenase (psbA) gene. *Journal of bacteriology*. **182** (21), 6233, **2000**.
 103. GROSSE C., BRANDT N., VAN ANTWERPEN P., WINTJENS R., MATTHIJS S. Two new siderophores produced by *Pseudomonas* sp. NCIMB 10586: The anti-oomycete non-ribosomal peptide synthetase-dependent mupirochelin and the NRPS-independent triabactin. *Frontiers in Microbiology*. **14**, 1143861, **2023**.

104. DANOUCHE M., EL GHACHTOULI N., EL BAOUCHI A., EL ARROUSSI H. Heavy metals phycoremediation using tolerant green microalgae : enzymatic and non-enzymatic antioxidant systems for the management of oxidative stress. *Journal of Environmental Chemical Engineering*. **8**, 104460, **2020**.
105. BALZANO S., SARDO A., BLASIO M., CHAHINE T.B., DELL'ANNO F., SANSONE C., BRUNET C. Microalgal metallothioneins and phytochelatins and their potential use in bioremediation. *Frontiers in Microbiology*. **11**, 527494, **2020**.
106. DEAN A.P., HARTLEY A., MCINTOSH O.A., SMITH A., FEORD H.K., HOLMBERG N.H., KING T., YARDLEY E., WHITE K.N., PITTMAN J.K. Metabolic adaptation of a *Chlamydomonas acidophila* strain isolated from acid mine drainage ponds with low eukaryotic diversity. *Science of the Total Environment*. **647**, 75, **2019**.
107. KOLAJ-ROBIN O., RUSSELL D., HAYES K.A., PEMBROKE J.T., SOULIMANE T. Cation diffusion facilitator family: structure and function. *FEBS letters*. **589** (12), 1283, **2015**.
108. VARGHESE B.R., GOH K.G., DESAI D., ACHARYA D., CHEE C., SULLIVAN M.J., ULETT G.C. Variable resistance to zinc intoxication among *Streptococcus agalactiae* reveals a novel IS 1381 insertion element within the zinc efflux transporter gene *czcD*. *Frontiers in Immunology*. **14**, 1174695, **2023**.
109. SURYAWATI B. Zinc homeostasis mechanism and its role in bacterial virulence capacity. In *AIP Conference Proceedings*, AIP Publishing. Volume **2021**, 1, **2018**.
110. CHATTERJEE S., KUMARI S., RATH S., PRIYADARSHANEE M., DAS S. Diversity, structure and regulation of microbial metallothionein: Metal resistance and possible applications in sequestration of toxic metals. *Metallomics*. **12** (11), 1637, **2020**.
111. LIU T., NAKASHIMA S., HIROSE K., UEMURA Y., SHIBASAKA M., KATSUHARA M., KASAMO K. A metallothionein and CPx-ATPase handle heavy-metal tolerance in the filamentous cyanobacterium *Oscillatoria brevis*. *FEBS letters*. **542** (1-3), 159, **2003**.
112. NATASHA SHAHID M., KHALID S., BIBI I., BUNDSCHUH J., KHAN NIAZI N., DUMAT C. A critical review of mercury speciation, bioavailability, toxicity and detoxification in soil-plant environment: ecotoxicology and health risk assessment. *Science of the Total Environment*. **711**, 134749, **2020**.
113. DURAND M.J., HUA A., JOUANNEAU S., CREGUT M., THOUAND G. Detection of metal and organometallic compounds with bioluminescent bacterial bioassays. In *Bioluminescence: Fundamentals and Applications in Biotechnology*, 1st ed.; Thouand G., Marks R., (eds) Publisher: Springer Cham, Switzerland. **3**, 77, **2016**.
114. MEYER F., PUPKES H., STEINBÜCHEL A. Development of an improved system for the generation of knockout mutants of *Amycolatopsis* sp. strain ATCC 39116. *Applied and Environmental Microbiology*. **83** (3), e02660, **2017**.
115. PRIYADARSHANEE M., CHATTERJEE S., RATH S., DASH H.R., DAS S. Cellular and genetic mechanism of bacterial mercury resistance and their role in biogeochemistry and bioremediation. *Journal of Hazardous Materials*. **423**, 126985, **2022**.
116. DASH H.R., DAS S. Bioremediation of mercury and the importance of bacterial mer genes. *International Biodeterioration & Biodegradation*. **75**, 207, **2012**.
117. NAGUIB M.M., EL-GENDY A.O., KHAIRALLA A.S. Microbial diversity of operon genes and their potential roles in mercury bioremediation and resistance. *The Open Biotechnology Journal*. **12** (1), 56, **2018**.
118. WILSON J.R., LEANG C., MORBY A.P., HOBMAN J.L., BROWN N.L. MerF is a mercury transport protein: different structures but a common mechanism for mercuric ion transporters? *FEBS Letters*. **472**, 78, **2000**.
119. CHRISTAKIS C.A., BARKAY T., BOYD E.S. Expanded diversity and phylogeny of mer genes broadens mercury resistance paradigms and reveals an origin for MerA among thermophilic Archaea. *Frontiers in Microbiology*. **12**, 682605, **2021**.
120. COOPER C.J., ZHENG K., RUSH K.W., JOHS A., SANDERS B.C., PAVLOPOULOU G.A., PARKS J.M. Structure determination of the HgcAB complex using metagenome sequence data: insights into microbial mercury methylation. *Communications Biology*. **3** (1), 320, **2020**.
121. GIONFRIDDO C.M., WYMORE A.M., JONES D.S., LYNES M.M., CHRISTENSEN G.A., PODAR M., ELIAS D.A. An improved hgcAB primer set and direct high-throughput sequencing expand Hg-methylator diversity in nature. *Frontiers in Microbiology*. **11**, 541554, **2020**.
122. VILLAR E., CABROL L., HEIMBÜRGER-BOAVIDA L.E. Widespread microbial mercury methylation genes in the global ocean. *Environmental Microbiology Reports*. **12** (3), 277, **2020**.
123. SHI W., MA X. Effects of heavy metal Cd pollution on microbial activities in soil. *Annals of Agricultural and Environmental Medicine*. **24** (4), **2017**.
124. CHEN J., WANG L., LI W., ZHENG X., LI X. Genomic insights into cadmium resistance of a newly isolated, plasmid-free *Cellulomonas* sp. strain Y8. *Frontiers in Microbiology*. **12**, 784575, **2022**.
125. CHEN J., XING C., ZHENG X., LI X. Complete genome sequence of *Cellulomonas* sp. Strain Y8, a high-GC-content plasmid-free heavy metalresistant bacterium isolated from farmland soil. *Microbiology Resource Announcements*. **8**, e01066, **2019**.
126. CHEN J., XING C., ZHENG X., LI X., CHEN J., XING C., ZHENG X., LI X. Functional genomic identification of cadmium resistance genes from a high gc clone library by coupling the sanger and PacBio sequencing strategies. *Genes (Basel)*. **11**, 7, **2019**.
127. GRØNBERG C., HU Q., MAHATO D.R., LONGHIN E., SALUSTROS N., DUELLI A., GOURDON P. Structure and ion-release mechanism of PIB-4-type ATPases. *Elife*. **10**, e73124, **2021**.
128. XIA Y., XU Y., ZHOU Y., YU Y., CHEN Y., LI C., TAO J. Comparative genome analyses uncovered the cadmium resistance mechanism of *Enterobacter cloacae*. *International Microbiology*. **26** (1), 99, **2023**.
129. YONG X., CHEN Y., LIU W., XU L., ZHOU J., WANG S., ZHENG T. Enhanced cadmium resistance and accumulation in *Pseudomonas putida* KT2440 expressing the phytochelatin synthase gene of *Schizosaccharomyces pombe*. *Letters in Applied Microbiology*. **58** (3), 255, **2014**.
130. LUKÁCS M., CSILLA PÁLINKÁS D., SZUNYOG G., VÁRNAGY K. Metal binding ability of small peptides containing cysteine residues. *ChemistryOpen*. **10** (4), 451, **2021**.
131. PATIL A., CHAKRABORTY S., YADAV Y., SHARMA B., SINGH S., ARYA M. Bioremediation strategies

- and mechanisms of bacteria for resistance against heavy metals: a review. *Bioremediation Journal*. **1**, 2024.
132. LIU H., HUANG H., LIANG K., LIN K., SHANGGUAN Y., XU H. Characterization of a cadmium-resistant functional bacteria (*Burkholderia* sp. SRB-1) and mechanism analysis at physiochemical and genetic level. *Environmental Science and Pollution Research*. **30** (32), 78408, 2023.
 133. HUANG H., WANG K., LI S., LIANG K., DAI J., JIAN J., XU H. Different survival strategies of the phosphate-mineralizing bacterium *Enterobacter* sp. PMB-5 in response to cadmium stress: Biomineralization, biosorption, and bioaccumulation. *Journal of Hazardous Materials*. **465**, 133284, 2024.
 134. JANG S. Multidrug efflux pumps in *Staphylococcus aureus* and their clinical implications. *Journal of Microbiology*. **54**, 1, 2016.
 135. ABBAS S.Z., RAFATULLAH M., HOSSAIN K., ISMAIL N., TAJARUDIN H.A., KHALIL H.P.S.A. A review on mechanism and future perspectives of cadmium-resistant bacteria. *International Journal of Environmental Science and Technology*. **15**, 243, 2018.
 136. XIA L., YIN C., CAI L., QIU G., QIN W., PENG B., LIU J. Metabolic changes of *Acidithiobacillus caldus* under Cu2 stress? *Journal of Basic Microbiology*. **50**, 591, 2010.
 137. NIES D.H. Efflux-mediated heavy metal resistance in prokaryotes. *FEMS Microbiology Reviews*. **27**(2-3), 313, 2003.
 138. CHEN M., LI Y., ZHANG L., WANG J., ZHENG C., ZHANG X. Analysis of gene expression provides insights into the mechanism of cadmium tolerance in *Acidithiobacillus ferrooxidans*. *Current Microbiology*. **70**, 290, 2015.
 139. LIU H., ZHANG Y., WANG Y., XIE X., SHI Q. The connection between Czc and Cad systems involved in cadmium resistance in *Pseudomonas putida*. *International Journal of Molecular Sciences*. **22**(18), 9697, 2021.
 140. AYANGBENRO A.S., BABALOLA O.O. Genomic analysis of *Bacillus cereus* NWUAB01 and its heavy metal removal from polluted soil. *Scientific Reports*. **10**(1), 19660, 2020.
 141. NIES D.H. CzcR and CzcD, gene products affecting regulation of resistance to cobalt, zinc, and cadmium (czc system) in *Alcaligenes eutrophus*. *Journal of Bacteriology*. **174**(24), 8102, 1992.
 142. SHIN M.N., SHIM J., YOU Y., MYUNG H., BANG K.S., CHO M., OH B.T. Characterization of lead resistant endophytic *Bacillus* sp. MN3-4 and its potential for promoting lead accumulation in metal hyperaccumulator *Alnus firma*. *Journal of Hazardous Materials*. **199**, 314, 2012.
 143. SHI L., DENG X.P., YANG Y., JIA Q.Y., WANG C.C., SHEN Z.G. A Cr(VI)-tolerant strain, *Pisolithus* sp.1, with a high accumulation capacity of Cr in mycelium and highly efficient assisting *Pinus thunbergii* for phytoremediation. *Chemosphere*. **224**, 862, 2019.
 144. RAFEEQ H., AFSHEEN N., RAFIQUE S., ARSHAD A., INTISAR M., HUSSAIN A., IQBAL H.M. Genetically engineered microorganisms for environmental remediation. *Chemosphere*. **310**, 136751, 2023.
 145. MAQBOOL Z., HUSSAIN S., AHMAD T., NADEEM H., IMRAN M., KHALID A., MARTIN-LAURENT F. Use of RSM modeling for optimizing decolorization of simulated textile wastewater by *Pseudomonas aeruginosa* strain ZM130 capable of simultaneous removal of reactive dyes and hexavalent chromium. *Environmental Science and Pollution Research*. **23** (11), 11224, 2016.
 146. LAWAL A.T. Polycyclic aromatic hydrocarbons. A review. *Cogent Environmental Science*. **3**(1), 1339841, 2017.
 147. ISMAIL N.A., KASMURI N., HAMZAH N., JAAFAR J., MOJIRI A., KINDAICHI T. Influence of pH and concentration on the growth of bacteria-fungus and benzo [a] pyrene degradation. *Environmental Technology & Innovation*. **29**, 102995, 2023.
 148. KEBEDE G., TAFESE T., ABDA E.M., KAMARAJ M., ASSEFA F. Factors influencing the bacterial bioremediation of hydrocarbon contaminants in the soil: mechanisms and impacts. *Journal of Chemistry*. **2021**(1), 9823362, 2021.
 149. BHANDARI S., POUDEL D.K., MARAHATHA R., DAWADI S., KHADAYAT K., PHUYAL S., PARAJULI N. Microbial enzymes used in bioremediation. *Journal of Chemistry*. **2021**(1), 8849512, 2021.
 150. IMAM A., SUMAN S.K., KANAUJIA P.K., RAY A. Biological machinery for polycyclic aromatic hydrocarbons degradation: A review. *Bioresource Technology*. **343**, 126121, 2022.
 151. SINGH S.K., SINGH R.K. An overview on remediation technologies for polycyclic aromatic hydrocarbons in contaminated lands: a critical approach. *Environment, Development and Sustainability*. **1**, 2023.
 152. JOUANNEAU Y., MEYER C., DURAFFOURG N. Dihydroxylation of four- and five-ring aromatic hydrocarbons by the naphthalene dioxygenase from *Sphingomonas* CHY-1. *Applied Microbiology and Biotechnology*. **100**, 1253, 2016.
 153. ZADA S., ZHOU H., XIE J., HU Z., ALI S., SAJJAD W., WANG H. Bacterial degradation of pyrene: biochemical reactions and mechanisms. *International Biodeterioration & Biodegradation*. **162**, 105233, 2021.
 154. WANG Q., HOU J., HUANG Y., LIU W., CHRISTIE P. Metagenomics reveals mechanism of pyrene degradation by an enriched bacterial consortium from a coking site contaminated with PAHs. *Science of The Total Environment*. **904**, 166759, 2023.
 155. ZHANG L., QIU X., HUANG L., XU J., WANG W., LI Z., XU, P. TANG H. Microbial degradation of multiple PAHs by a microbial consortium and its application on contaminated wastewater. *The Journal of Hazardous Materials*. **419**, 126524, 2021.
 156. ZHOU G., QIAO H., LIU Y., YU X., NIU X. High phenanthrene degrading efficiency by different microbial compositions construction. *Frontiers in Microbiology*. **15**, 1439216, 2024.
 157. KIM B.J., KIM B.R., JEONG J., LIM J.H., PARK S.H., LEE S.H., KIM B.J. A description of *Mycobacterium chelonae* subsp. *gwanakae* subsp. nov., a rapidly growing mycobacterium with a smooth colony phenotype due to glycopeptidolipids. *International Journal of Systematic and Evolutionary Microbiology*. **68** (12), 3772, 2018.
 158. WU Y., XU Y., ZHOU N. A newly defined dioxygenase system from *Mycobacterium vanbaalenii* PYR-1 endowed with an enhanced activity of dihydroxylation of high-molecular-weight polyaromatic hydrocarbons. *Frontiers of Environmental Science & Engineering*. **14**, 1, 2020.
 159. KIM S.J., SONG J., KWEON O., HOLLAND R.D., KIM D.W., KIM J., CERNIGLIA C.E. Functional robustness of a polycyclic aromatic hydrocarbon metabolic network examined in a *nidA* aromatic ring-hydroxylating oxygenase mutant of *Mycobacterium vanbaalenii* PYR-1.

- Applied and Environmental Microbiology. **78** (10), 3715, **2012**.
160. MATHEW J.J., NAIR A., SAJESHKUMAR N., JOSE P. Analysis of the phylloremediation capability of *Mangifera indica* in hydrocarbon polluted area: An outlook study. *Journal of Medicinal Plants*. **10**, 125, **2022**.
 161. MAWAD A.M., ABDEL-MAGEED W.S., HESHAM A.E.-L. (2020). Quantification of naphthalene dioxygenase (NahAC) and catechol dioxygenase (C23O) catabolic genes produced by phenanthrene-degrading *Pseudomonas fluorescens* AH-40. *Current Genomics*. **21** (2), 111, **2020**.
 162. ZADA S., ZHOU H., XIE J., HU Z., ALI S., SAJJAD W., WANG H. Bacterial degradation of pyrene: biochemical reactions and mechanisms. *International Biodeterioration & Biodegradation*. **162**, 105233, **2021**.
 163. FERNÁNDEZ M., NIQUI-ARROYO J.L., CONDE S., RAMOS J.L., DUQUE E. Enhanced tolerance to naphthalene and enhanced rhizoremediation performance for *Pseudomonas putida* KT2440 via the NAH7 catabolic plasmid. *Applied and Environmental Microbiology*. **78**(15), 5104, **2012**.
 164. SCHELL M.A. Homology between nucleotide sequences of promoter regions of nah and sal operons of NAH7 plasmid of *Pseudomonas putida*. *Proceedings of the National Academy of Sciences*. **83** (2), 369, **1986**.
 165. GHOSAL D., GHOSH S., DUTTA T.K., AHN Y. Current state of knowledge in microbial degradation of polycyclic aromatic hydrocarbons (PAHs): a review. *Frontiers in Microbiology*. **7**, 1369, **2016**.
 166. LLOYD-JONES G., LAURIE A.D., HUNTER D.W., FRASER R. Analysis of catabolic genes for naphthalene and phenanthrene degradation in contaminated New Zealand soils. *FEMS Microbiology Ecology*. **29** (1), 69, **1999**.
 167. OVERHAGE J., KRESSE A.U., PRIEFERT H., SOMMER H., KRAMMER G., RABENHORST J., STEINBÜCHEL A. (1999). Molecular characterization of the genes *pcaG* and *pcaH*, encoding protocatechuate 3, 4-dioxygenase, which are essential for vanillin catabolism in *Pseudomonas* sp. strain HR199. *Applied and Environmental Microbiology*. **65** (3), 951, **1999**.
 168. GUZIK U., HUPERT-KOCUREK K., SITNIK M., WOJCISZYŃSKA D. Protocatechuate 3,4-dioxygenase: a wide substrate specificity enzyme isolated from *Stenotrophomonas maltophilia* KB2 as a useful tool in aromatic acid biodegradation. *Journal of Molecular Microbiology and Biotechnology*. **24** (3), 150, **2014**.
 169. WOLF M.E., LALANDE A.T., NEWMAN B.L. BLEEM A.C., PALUMBO C.T., BECKHAM G.T., ELTIS L.D. The catabolism of lignin-derived p-methoxylated aromatic compounds by *Rhodococcus jostii* RHA1. *Applied and Environmental Microbiology*. **90** (3), e02155, **2024**.
 170. QUTOB M., RAFATULLAH M., MUHAMMAD S.A., ALOSAIMI A.M., ALORFI H.S., HUSSEIN M.A. A Review of Pyrene Bioremediation Using *Mycobacterium* Strains in a Different Matrix. *Fermentation*. **8** (6), 260, **2022**.
 171. KWEON O., KIM S.J., JONES R.C., FREEMAN J.P., ADJEI M.D., EDMONDSON R.D., CERNIGLIA C.E. A polyomic approach to elucidate the fluoranthene-degradative pathway in *Mycobacterium vanbaalenii* PYR-1. *Journal of Bacteriology*. **189** (13), 4635, **2007**.
 172. DONG C., BAI X., LAI Q., XIE Y., CHEN X., SHAO Z. Draft Genome sequence of *Sphingobium* sp. strain C100, a polycyclic aromatic hydrocarbon-degrading bacterium from the deep-sea sediment of the Arctic Ocean. *Genome Announc.* **2** (1), e01210, **2014**.
 173. ZHAO Q., HU H., WANG W., PENG H., ZHANG X. Genome sequence of *Sphingobium yanoikuyae* B1, a polycyclic aromatic hydrocarbon-degrading strain. *Genome Announc.* **3** (1):e01522, **2015**.
 174. ZHAO Q., YUE S., BILAL M., HU H., WANG W., ZHANG X. Comparative genomic analysis of 26 *Sphingomonas* and *Sphingobium* strains: dissemination of bioremediation capabilities, biodegradation potential and horizontal gene transfer. *Science of the Total Environment*. **609**, 1238, **2017**.
 175. MACCHI M., MARTINEZ M., TAUIL R.N., VALACCO M.P., MORELLI I.S., COPPOTELLI B.M. Insights into the genome and proteome of *Sphingomonas paucimobilis* strain 20006FA involved in the regulation of polycyclic aromatic hydrocarbon degradation. *World Journal of Microbiology and Biotechnology*. **34**, 1, **2018**.
 176. ASHRAF M.A. Persistent organic pollutants (POPs): a global issue, a global challenge. *Environmental Science and Pollution Research*. **24**, 4223, **2017**.
 177. LAURIE A.D., LLOYD-JONES G. The *phn* genes of *Burkholderia* sp. strain RP007 constitute a divergent gene cluster for polycyclic aromatic hydrocarbon catabolism. *Journal of Bacteriology*. **181** (2), 531, **1999**.
 178. OKOLAFOR F.I., EKHAISE F.O. Molecular characterisation of bacteria monooxygenase and dehydrogenase genes involved in PAHs remediation. *Open Journal of Environmental Research*. **3** (2), 25, **2022**.
 179. CARPENTER D.O. Polychlorinated biphenyls (PCBs): routes of exposure and effects on human health. *Reviews on Environmental Health*. **21**(1), 1, **2006**.
 180. SHEBL S., GHAREEB D.A., ALI S.M., GHANEM N.B.E.D., OLAMA Z.A. Aerobic phenol degradation using native bacterial consortium via ortho- and meta-cleavage pathways. *Frontiers in Microbiology*. **15**, 1400033, **2024**.
 181. MARDANI G., MAHVI A.H., HASHEMZADEH-CHALESHTORI M., NASERI S., DEGHANI M.H., GHASEMI-DEHKORDI P. Application of genetically engineered dioxygenase producing *Pseudomonas putida* on decomposition of oil from spiked soil. *Jundishapur Journal of Natural Pharmaceutical Products*. **12** (3), e64313, **2017**.
 182. JIANG Y., YANG X., LIU B., ZHAO H., CHENG Q., CAI B. Catechol 2, 3-dioxygenase from *Pseudomonas* sp. strain ND6: gene sequence and enzyme characterization. *Bioscience, Biotechnology, and Biochemistry*. **68** (8), 1798, **2004**.
 183. LI S., QIN K., LI H., GUO J., LI D., LIU F., ZHAO H. Cloning and characterisation of four *catA* genes located on the chromosome and large plasmid of *Pseudomonas putida* ND6. *Electronic Journal of Biotechnology*. **34**, 83, **2018**.
 184. ABBAS N.H., ELSAYED A., HASSAN H.A., EL-SABBAGH S., ELBAZ A.F., KHALIL H. Characterization and Expression Analysis of Extradiol and Intradiol Dioxygenase of Phenol-Degrading Haloalkaliphilic Bacterial Isolates. *Current Microbiology*. **79** (10), 294, **2022**.
 185. JIANG J., TIAN W., LU Z., CHU M., CAO H., ZHANG, D. Cometabolic degradation of pyrene with phenanthrene as substrate: assisted by halophilic *Pseudomonas stutzeri* DJP1. *Biodegradation*. **34** (6), 519, **2023**.

186. ZEYAUULLAH M., AHMAD RA. Z.I., NASEEM A.S.M.A., ISLAM B.A.D.R.U.L., HASAN H.M., ABDELKAFE A.S., ALI A.R.I.F. Catechol biodegradation by *Pseudomonas* strain: a critical analysis. *International Journal of Chemical Sciences*. **7** (3), 2211, **2009**.
187. FUJIHARA H., HIROSE J., SUENAGA H. Evolution of genetic architecture and gene regulation in biphenyl/PCB-degrading bacteria. *Frontiers in Microbiology*. **14**, 1168246, **2023**.
188. SUENAGA H., YAMAZOE A., HOSOYAMA A., KIMURA N., HIROSE J., WATANABE T. Draft genome sequence of the polychlorinated biphenyl-degrading bacterium *Cupriavidus basilensis* KF708 (NBRC 110671) isolated from biphenyl-contaminated soil. *Genome Announcements*. **3**, 143, **2015**.
189. HIROSE J., FUJIHARA H., WATANABE T., KIMURA N., SUENAGA H., FUTAGAMI T., FURUKAWA K. Biphenyl/PCB degrading bph genes of ten bacterial strains isolated from biphenyl-contaminated soil in Kitakyushu, Japan: comparative and dynamic features as integrative conjugative elements (ICEs). *Genes*. **10** (5), 404, **2019**.
190. SUENAGA H., FUJIHARA H., KIMURA N., HIROSE J., WATANABE T., FUTAGAMI T. Insights into the genomic plasticity of *Pseudomonas putida* KF715, a strain with unique biphenyl-utilizing activity and genome instability properties. *Environmental Microbiology Reports*. **9**, 589, **2017**.
191. SAMIN, AFZAL B., REHMANJI M., NAAZ H., YASIN D., JUTUR P.P., FATMA T. Insight into the mechanism of estrone biodegradation by *Spirulina* CPCC-695. *Environment, Development and Sustainability*. **1**, **2023**.
192. SUENAGA H., MATSUZAWA T., SAHARA T. Discovery by metagenomics of a functional tandem repeat sequence that controls gene expression in bacteria. *FEMS Microbiology Ecology*. **98** (4), fiac037, **2022**.
193. FOLAKE O. Molecular identification of crude oil-degrading bacteria and screening for catechol 2, 3 dioxygenase (C23O) gene. *Biotechnology Journal International*. **23** (4), 1, **2020**.
194. YEN K.M., GUNSALUS I.C. Regulation of naphthalene catabolic genes of plasmid NAH7. *Journal of Bacteriology*. **162** (3), 1008, **1985**.
195. EIP-AGRI. Protecting agricultural soils from contamination. MINI PAPER 3: Biological remediation of contaminated agricultural soils. Available online: <https://ec.europa.eu/eip/agriculture/en/focus-groups/protecting-agricultural-soils-contamination.html> (accessed on 14.09.2024). **2024**.
196. ADETUNJI C.O., ANANI O.A. Recent advances in the application of genetically engineered microorganisms for microbial rejuvenation of contaminated environment. *Microbial Rejuvenation of Polluted Environment*. **3**, 303, **2021**.
197. WU C., LI F., YI S., GE F. Genetically engineered microbial remediation of soils co-contaminated by heavy metals and polycyclic aromatic hydrocarbons: advances and ecological risk assessment. *Journal of Environmental Management*. **296**, 113185, **2021**.
198. JAISWAL S., SHUKLA P. Alternative strategies for microbial remediation of pollutants via synthetic biology. *Frontiers in Microbiology*. **11**, 808, **2020**.
199. CORDIS-EU. Clean, fast and effective – new approach transforms contaminated soil. 2020. Available online: <https://cordis.europa.eu/article/id/415943-clean-fast-and-effective-new-approach-transforms-contaminated-soil> (accessed on 3.01.2025). **2019**.
200. FERRARESE E., ANDREOTTOLA G., OPREA I.A. Remediation of PAH-contaminated sediments by chemical oxidation. *Journal of Hazardous Materials*. **152** (1), 128, **2008**.
201. CHEN B., XU J., LU H., ZHU L. Remediation of benzo [a] pyrene contaminated soils by moderate chemical oxidation coupled with microbial degradation. *Science of The Total Environment*. **871**, 161801, **2023**.
202. LEMAIRE J., MORA V., FAURE P., HANNA K., BUÈS M., SIMONNOT M.O. Chemical oxidation efficiency for aged, PAH-contaminated sites: An investigation of limiting factors. *Journal of Environmental Chemical Engineering*. **7** (3), 103061, **2019**.
203. REDDY V.A., SOLANKI C.H., KUMAR S., REDDY K.R., DU Y.J. Stabilization/solidification of zinc-and lead-contaminated soil using limestone calcined clay cement (LC3): An environmentally friendly alternative. *Sustainability*. **12** (9), 3725, **2020**.
204. SAEED K., AL-KHYAT S., ABD HACHEEM Z., FARTOSY S.H. Evaluating the Efficiency of Alkaline Activator with Silica-Rich Wastes in Stabilizing Cadmium-Contaminated Soil. *Civil Engineering Journal*. **10** (7), 2123, **2024**.
205. KUMARI P., KUMAR A. Advanced Oxidation Process: A remediation technique for organic and non-biodegradable pollutant. *Results in Surfaces and Interfaces*. **11**, 100122, **2023**.
206. AZIZ S., REHMAN N.U., SOONMIN H., AHMAD N. Novel Ni/ZnO Nanocomposites for the Effective Photocatalytic Degradation of Malachite Green Dye. *Civil Engineering Journal*. **10** (8), **2024**.
207. KHAN S., NAUSHAD M., AL-GHEETHI A., IQBAL J. Engineered nanoparticles for removal of pollutants from wastewater: Current status and future prospects of nanotechnology for remediation strategies. *Journal of Environmental Chemical Engineering*. **9** (5), 106160, **2021**.
208. ROY A., SHARMA A., YADAV S., JULE L.T., KRISHNARAJ R. Nanomaterials for remediation of environmental pollutants. *Bioinorganic Chemistry and Applications*. **2021** (1), 1764647, **2021**.
209. CEBALLOS-ESCALERA A., POUS N., BALAGUER M.D., PUIG S. Nitrate electro-bioremediation and water disinfection for rural areas. *Chemosphere*. **352**, 141370, **2024**.
210. ANAND U., DEY S., PARIAL D., FEDERICI S., DUCOLI S., BOLAN N.S., BONTEMPI E. Algae and bacteria consortia for wastewater decontamination and transformation into biodiesel, bioethanol, biohydrogen, biofertilizers and animal feed: a review. *Environmental Chemistry Letters*. **21** (3), 1585, **2023**.
211. CEBALLOS-ESCALERA A., POUS N., BAÑERAS L., BALAGUER M.D., PUIG S. Advancing towards electro-bioremediation scaling-up: On-site pilot plant for successful nitrate-contaminated groundwater treatment. *Water Research*. **256**, 121618, **2024**.
212. ADHIKARY S., SAHA J., DUTTA P., PAL A. Bacterial Homeostasis and Tolerance to Potentially Toxic Metals and Metalloids through Diverse Transporters: Metal-Specific Insights. *Geomicrobiology Journal*. **41** (5), 496, **2024**.

213. KUPPAN N., PADMAN M., MAHADEVA M., SRINIVASAN S., DEVARAJAN R. A comprehensive review of sustainable bioremediation techniques: Eco friendly solutions for waste and pollution management. *Waste Management Bulletin*. **2** (3), 154, **2024**.
214. MADISON A.S., SORSBY S.J., WANG Y., KEY T.A. Increasing in situ bioremediation effectiveness through field-scale application of molecular biological tools. *Frontiers in Microbiology*. **13** (2), 1, **2022**.
215. KOOLIVAND A., COULON F., BALL A.S., ISMAIL N.I., KHUDUR L.S., PARSIMEHR M., GODINI K. Challenges with bioaugmentation and field-scale application of bioremediation processes for petroleum-contaminated sites: A review. *Indian Journal of Microbiology*. **1**, **2024**.
216. RAKLAMI A., MEDDICH A., OUFDOU K., BASLAM M. Plants – Microorganisms-based bioremediation for heavy metal cleanup: Recent developments, phytoremediation techniques, regulation mechanisms, and molecular responses. *International Journal of Molecular Sciences*. **23** (9), 5031, **2022**.
217. ABO-ALKASEM M.I., HASSAN N.M.H., ABO ELSOUD M.M. Microbial bioremediation as a tool for the removal of heavy metals. *Bulletin of the National Research Centre*. **47** (1), 31, **2023**.
218. BOLAN S., PADHYE L.P., JASEMIZAD T., GOVARTHANAN M., KARMEGAM N., WIJESEKARA H., BOLAN N. Impacts of climate change on the fate of contaminants through extreme weather events. *Science of The Total Environment*. **909**, 168388, **2024**.
219. SARAVANAN A., KUMAR P.S., VO D.V.N., JEEVANANTHAM S., KARISHMA S., YAASHIKAA P.R. A review on catalytic-enzyme degradation of toxic environmental pollutants: Microbial enzymes. *Journal of Hazardous Materials*. **419**, 126451, **2021**.