

Original Research

Biogas Slurry Application to Enhance Spinach (*Spinacia oleracea* L.) Growth and Improve Polluted Soil Properties

**Hossam S. El-Beltagi^{10*}, Mohamed Fathi El-Nady², Mohamed M. El-Mogy³,
Metwaly Mahfouz Salem Metwaly², Ahmed Mahmoud Ismail^{3,4},
Mahmoud A. Shabana⁵, Hosny Hamed Kesba³, Saffaa S. M. Abd El-Al⁵,
Wael Elmenofy³, Mohamed A. Abd El-Aziz⁵, Lamy Hamed⁶,
Kholoud Ahmed El-Naqma⁵**

¹Agricultural Biotechnology Department, College of Agriculture and Food Sciences, King Faisal University, Al-Ahsa 31982, Saudi Arabia

²Agricultural Botany Department, Faculty of Agriculture, Kafrelsheikh University, 33516, Kafr El-Sheikh, Egypt

³Department of Arid Land Agriculture, College of Agriculture and Food Sciences, King Faisal University, Al-Ahsa 31982, Saudi Arabia

⁴Pests and Plant Diseases Unit, College of Agriculture and Food Sciences, King Faisal University, Al-Ahsa 31982, Saudi Arabia

⁵Soil Water and Environment Research Institute, Agricultural Research Center, Giza, 12619, Egypt

⁶Department of Environment and Agricultural Natural Resources, College of Agricultural and Food Sciences, King Faisal University, Al-Ahsa 31982, Saudi Arabia

Received: 24 February 2025

Accepted: 19 April 2025

Abstract

Biogas slurry contains high levels of organic matter and essential nutrients, which enhance soil fertility and structure. Although it can reduce the uptake of heavy metals by plants, improper application may result in soil contamination. This research evaluated the effects of applying semi-solid biogas slurry (SSBS) at 72 or 96 m³/ha, either on its own or combined with a foliar spray of liquid biogas slurry (LBS) at 48 L/ha or a recommended NPK fertilizer on spinach growth. Results showed that all SSBS treatments, whether individually or combined with NPK or LBS, improved growth traits such as plant height and total leaf area. They also increased chlorophyll content (chlorophyll a, b, and total), vitamin C, total phenolic compounds in leaves, and concentrations of nitrogen, phosphorus, potassium, iron, and zinc, along with enhanced nitrate reductase activity. Furthermore, these treatments boosted leafy yield and lowered levels of lead (Pb), cadmium (Cd), nitrate, and oxalate content in spinach leaves. The most pronounced positive effects on growth and physiological-biochemical parameters occurred when SSBS

*e-mail: helbeltagi@kfu.edu.sa

Tel.: +966541775875

°ORCID iD: 0000-0003-4433-2034

was applied at 96 m³/ha along with either the recommended NPK dose or LBS. These findings indicate that biogas slurry could serve as a sustainable alternative to chemical fertilizers in agriculture.

Keywords: chlorophyll, lead, cadmium, oxalate, vitamin C

Introduction

Spinach (*Spinacia oleracea* L.), a member of the Chenopodiaceae family, is a widely cultivated leafy vegetable with an annual global production of 26.2 million tons (FAO 2021). Its leaves are highly nutritious and are an excellent source of various vitamins (A, B2, B6, B9, C, E, K, or folate), minerals (including potassium, calcium, magnesium, iron, manganese, selenium), and dietary fiber [1].

Soil contamination by heavy metals occurs due to natural processes like weathering and erosion of parent rock material or ore deposits, as well as artificial sources such as wastewater irrigation and mining activities. Heavy metal contamination in soil is a significant environmental issue that poses risks to ecosystems, agricultural productivity, and human health. The presence of one contaminant can influence the effects of others. Most research to date has focused on the impact of individual metals on plants [2]. Heavy metals typically exist in an insoluble form in soil, making them inaccessible to plants. However, plants may enhance their bioavailability by releasing root exudates that modify the pH of the rhizosphere, thereby increasing heavy metal uptake [3]. Common heavy metals that contaminate soils include lead (Pb), cadmium (Cd), mercury, arsenic, chromium, and nickel. Heavy metals may disrupt soil microbial communities, affecting nutrient cycling and soil structure [4].

Heavy metals like cadmium (Cd) and lead (Pb) can hinder plant growth by disrupting root and shoot development, resulting in stunted growth, reduced biomass, and decreased crop yields [5, 6]. They also trigger the production of reactive oxygen species (ROS) in plants, leading to oxidative stress that damages cellular components such as lipids, proteins, and DNA, affecting overall cell function or viability [5, 6]. Additionally, heavy metals impair photosynthesis by harming chloroplasts and reducing chlorophyll levels, which lowers photosynthetic efficiency and energy production necessary for plant growth [6, 7]. These metals can also interfere with enzymes that are essential for metabolic processes, such as those involved in nitrogen assimilation and carbohydrate metabolism, ultimately reducing plant productivity [6, 7]. On a cellular level, heavy metals can damage structures like cell membranes, mitochondria, and chloroplasts, leading to cell death and necrosis, further weakening plant growth [6, 7]. Moreover, heavy metals can cause mutations and alter gene expression, disrupting growth regulation and stress response pathways, potentially leading to long-term changes in plant physiology [6, 7].

Accumulation of heavy metals in plants can have significant impacts on animals and humans who consume these contaminated plants. These metals can build up in plant tissues and enter the food chain. As animals and humans eat these plants, the metals can accumulate in their bodies, leading to bioaccumulation. This may result in biomagnification, where the concentration of heavy metals increases with each level of the food chain [8]. Certain heavy metals, like cadmium (Cd), are known carcinogens and may elevate cancer risk. Lead (Pb) exposure can cause neurological harm, leading to cognitive and developmental problems, particularly in children. Prolonged exposure to heavy metals is also linked to cardiovascular diseases and kidney damage [8].

Biogas slurry can significantly decrease plants' uptake of heavy metals. The organic matter in the slurry contains functional groups capable of binding with heavy metals, lowering their bioavailability in the soil [9, 10]. This binding process restricts plants' absorption of heavy metals, mitigating their toxicity and improving soil properties [9]. Additionally, biogas slurry enhances soil fertility by providing essential nutrients such as nitrogen, phosphorus, or potassium, promoting microbial activity, and improving soil structure by increasing water retention and aeration [10]. The bioactive compounds in the slurry can also help suppress soilborne diseases, fostering healthier plant growth [11]. However, it's important to conduct a thorough risk assessment before using biogas slurry in agricultural fields to avoid introducing harmful levels of heavy metals [12].

Building on the previous information, a hypothesis was proposed that using biogas slurry as an organic fertilizer may significantly enhance plant growth and improve soil properties. By providing a natural source of nutrients, biogas slurry may reduce the reliance on chemical fertilizers, supporting more sustainable farming practices. To explore this, the study examined the effects of biogas slurry application on the growth, physio-biochemical responses of spinach plants, and soil properties over two growing seasons.

Materials and Methods

Experimental Location

Two field experiments were carried out in Al-Hamul district (latitude 31.24°N, longitude 31.04°E, and elevation of 14 m above sea level) in Kafr El-Sheikh Governorate, Egypt, during two consecutive growing seasons (2022 and 2023). These studies aimed

Table 1. Fundamental soil characteristics before cultivation during the seasons of 2022 and 2023.

Characteristics	Seasons values		Characteristics 2022	Seasons values	
	2022	2023		2023	
Chemical traits			Total porosity, %	46.04	45.66
OM, %	1.13	1.15	CEC	40.13	40.22
EC, dSm ⁻¹	3.32	3.27	F.C, %	45.28	45.54
pH	8.24	8.27	Bulk density	1.43	1.44
ESP%	13.21	12.91	Particle size distribution		
CaCo3	2.46	2.41	Clay, %	52.8	52.1
			Texture class	Clayey	Clayey
Availability of macro-nutrients (mg kg ⁻¹)					
Available	2022 season			2023 Season	
N	32.59			34.72	
P	5.86			6.27	
K	218			234	
Availability of micro-nutrients (mg kg ⁻¹)					
Available	2022 Season	2023 Season	Available	2022 Season	2023 Season
Fe	3.52	3.83	Pb	117	114
Zn	0.25	0.24	Cd	8	7.4
Cu	0.12	0.15			

to improve the effects of different forms of organic fertilizers, including semi-solid biogas slurry and liquid biogas, on the growth, physio-biochemical responses of spinach plants, and soil properties under conditions of soil contamination with lead (Pb) and cadmium (Cd).

Initial Soil Sample Analyses

Before the sowing process, initial soil samples were collected at a depth of 0-25 cm for analysis using standard methods based on Sparks et al. [13] and Dane and Topp [14]. Table 1 presents the fundamental soil characteristics before cultivation for the 2022 and 2023 seasons.

Experimental Design and Treatments

A completely randomized design in three replicates was utilized as follows:

1. Recommended dose of mineral fertilizers (Control, RD).
2. Semi-solid biogas slurry (SSBS) as 72 m³ha⁻¹ (Soil application).
3. SSBS 72 m³ with 48 Lha⁻¹ of liquid biogas slurry (LBS) as a foliar application.
4. SSBS 72 m³ha⁻¹ with RD as a foliar application.
5. SSBS 96 m³ha⁻¹ (Soil application).
6. SSBS 96 m³ha⁻¹ with 48 L (LBS-foliar application).
7. SSBS 96 m³ha⁻¹ with RD as a foliar application.

Studied Forms of Biogas Slurry

Semi-solid biogas slurry (SSBS) or liquid biogas slurry (LBS) were procured from the Egyptian commercial market, Corn Company for Sustainable Development, and their characteristics are shown in Table 2.

Cultivation and Harvest

Spinach (*Spinacia oleracea*) seeds of the Balady variety were sown in rows at a rate of 28.8 kg per hectare in the second week of September for both seasons studied. The cultivation area measured 12 m² (3.0×4.0 m). A recommended dose of 144 nitrogen units per hectare was applied for spinach production, adjusted for the soil's available nitrogen content (30 mg/kg). Mineral nitrogen was supplied as ammonium sulfate (21% nitrogen). In addition, all plots received 28.8 phosphorus units per hectare through calcium superphosphate (6.7% phosphorus) during soil preparation and 60 potassium units per hectare as potassium sulfate (39.8% potassium) at the second irrigation. Irrigation was carried out every 10 days. Semi-solid biogas slurry treatments were incorporated into the soil before planting. Liquid biogas slurry was applied in two doses: the first with the initial irrigation at planting and the second as a foliar spray 20 days after the first irrigation.

Table 2. Characteristics of semi-solid biogas slurry (SSBS) and liquid biogas slurry (LBS).

Characteristics	SSBS	LBS
pH	7.52	8.1
Ec, dSm ⁻¹	8.69	1.73
Organic matter, %	55.98	92.93
C/N ratio	20.10	1.13
N, %	1.3	1
P, %	1.1	1.91
K, %	3.68	1.21
Fe, mg kg ⁻¹	1.12	0.95
Zn, mg kg ⁻¹	25.1	0.003
Cu, mg kg ⁻¹	33.7	21.3

Sampling, Measurements, and Determination

Spinach was harvested at the pre-flowering stage, specifically when it reached the 5-6 leaf growth stage, occurring around 60 days after sowing in both experimental seasons. The following characteristics were then analyzed.

Vegetative Growth Characters

The parameters assessed included plant height (cm), dry weight (determined by drying plants in electric oven (BINDER Drying and Heating Chambers, China) at 70°C for 72 hours, measured in grams per plant), and leaf area (cm² per plant) using a portable laser leaf area meter (Ci-02) from CiD Bio-Science, USA. Additionally, at harvest, measurements were taken for plant height (cm), number of leaves per plant, and yield (Mg per hectare).

Chlorophyll Pigments Content

Thirty days after sowing, leaflets from the third leaf counted from the plant tip were collected and submerged in 5 mL of N,N-dimethylformamide for extraction. This extraction was conducted overnight at room temperature in the dark. Chlorophyll pigment concentrations (chlorophyll a, b, and total) were measured in µg/g of fresh weight following the procedure described by Moran [15].

Proline Content

At the thirty-day post-sowing, fresh leaves were utilized to determine proline content, measured in micromoles per gram of fresh weight (µmol g⁻¹ FW). This analysis was performed using a spectrophotometer (Shimadzu UV 1601) at a wavelength of 520 nm, following the protocol established by Bates et al. [16].

Total Phenols Content

The total phenolic compounds were quantified following the method outlined by Bessada et al. [17], with careful adherence to each step to ensure accuracy and reliability of measurements.

Mineral Element Nutrients

The chemical composition of leaves (N, P, K%, and Fe, Zn, Cu, Pb, Cd ppm) was determined. Plant sample digestion was done using HClO₄ + H₂SO₄ as mixture [18].

NPK Percentages

Then N, P, K percentages were determined utilizing MicroKjeldahl (Model: 054120-01 Japan), spectrophotometric and flame photometer (FP910 Brochure, Russian), respectively [19].

Fe, Mn, Zn, Cu, Pb, and Cd Content

The concentrations of iron (Fe), manganese (Mn), zinc (Zn), copper (Cu) in spinach plants were determined in parts per million (ppm) using Atomic Absorption Spectroscopy (SP-IAA320, InfiteK, China), following the Soil Science Research Institute by Chen et al. [20]. Lead (Pb) and cadmium (Cd) levels were measured in ppm using inductively coupled plasma-optical emission spectrometry (ICP-OES, ICAPTM 7000 Plus Series, Thermo ScientificTM, USA) after acid digestion with HNO₃ (69%) and H₂O₂ (30%) in a microwave digestion apparatus (Milestone MLS 1200 Mega), as described by Bettinelli et al. [21].

Nitrate and Nitrite (mg kg⁻¹) Contents

Nitrate or nitrite (mg kg⁻¹) contents in spinach leaves at the harvest stage were determined as described by Ozdestan and Uren [22].

Oxalate Content

Leaves were dried at 60°C, homogenized in deionized water, and oxalate concentration was measured using an oxalate kit following Palaniswamy et al. [23].

Nitrate Reductase Assay

The activity of nitrate reductase (NR) was assessed by measuring the production of NO²⁻. Leaf material extraction and NR activity measurements followed the method of Kaiser and Brendle-Behnisch [24], with some modifications. At specific times, leaves were flash-frozen in liquid nitrogen, ground into fine powder, and stored in liquid nitrogen until analyzed. For activity assay, 1 mL of a buffer containing 50 mM Hepes-KOH (pH 7.5), 2 mM DTT, 0.2% (v/v) Triton X-100, or MgCl₂

(at specified concentrations) was added to a frozen leaf sample (around 0.2 mg chlorophyll). The sample was then homogenized, and a 100 μL portion was added to 450 μL of reaction medium containing 40 mM Hepes-KOH, 5 mM KNO_3 , 0.2 mM NADH, and MgCl_2 . The interval between adding the medium to the leaf material and starting the assay was 2 min. Enzyme assay was conducted at 30°C for 2 min and halted by adding 100 μL of 1 M zinc acetate solution. Any remaining NADH was oxidized by adding 50 μL of 0.15 mM phenazine methosulfate. Nitrite formation was then measured according to the method of Hageman and Reed [25].

Soil Physical Analysis

In addition, soil properties were assessed at harvest, including bulk density (Mg m^{-3}), total porosity (%), pH, electrical conductivity (EC, dS m^{-1}), organic matter content (%), and available nitrogen (mg kg^{-1}), following methods outlined by Sparks et al. [13] and Dane and Topp [14]. Total microbial count (CFU mL^{-1}) in soil was determined by estimating the total number of microorganisms in the plant rhizosphere, as described by Allen [26].

Soil samples were air-dried and analyzed for some chemical characteristics, i.e., soil pH according to McLean [27]. Total soluble salts (EC) were determined utilizing electrical conductivity meters at 25°C in soil paste extract as dS/m [28]. Cation exchange capacity (CEC) expressed in (C moles kg^{-1}) was determined by the sodium acetate (NaOAC) method by Kim et al. [29]. Organic matter content (O.M) was measured according to Bhattacharyya et al [30].

The undisturbed soil samples were used to evaluate some of the physical properties, i.e., Soil bulk density (g cm^{-3}) and Total porosity (%) were determined according to Campbell [31]. Field capacity (FC) and permanent wilting point (PWP) were obtained by the pressure plate method, using pressure of 0.033 and 1.5 MPa, respectively, according to Klute and Dirksen [32].

Soil Chemical Analysis

The available soil nutrients were determined using various extraction methods following formal protocols. Specifically, the following extraction methods were used to assess the availability of different nutrients in the soil. Potassium sulfate (K_2SO_4) at a concentration of 1% was employed to extract available soil nitrogen (Kjeldahl method). The method was used to extract available soil phosphorus (spectrophotometric method) [33]. Ammonium acetate was used to extract available soil potassium (flame photometer method). Diethylene triamine penta acetic acid (DTPA) was utilized to extract available soil Fe, Zn, Cu, Pb, and Cd (BY AAS) [33].

Statistical Analyses

Statistical analysis was done as described by Gomez and Gomez [34] via CoStat (Version 6.303, CoHort, USA, 1998-2004) and DMRT (Duncan's)

Results and Discussion

Plant Height, Leaf Number/Plant, and Plant Leaf Area

The data in Fig. 1 indicate that treatments using semi-solid biogas slurry, whether alone or combined with liquid biogas slurry or the recommended NPK dose, improved spinach plant height and number of leaves per plant in comparison to control plants during the 2022 and 2023 seasons. The highest plant heights or leaf number per plant were observed for BS1 and BS2 combined with the recommended NPK dose, as well as for BS2 combined with 48L/ha of liquid biogas slurry, across both seasons. Fig. 1 illustrates that using both forms of biogas slurry, either combined with each other or with the recommended NPK dose, significantly increased spinach leaf area compared to the control across both growing seasons. The largest leaf area was observed with 96 m^3/ha of semi-solid biogas slurry (SSBS) combined with either the recommended NPK dose or 48 L/ha of liquid biogas slurry foliar application in both seasons.

The improvement in spinach plant height, leaf count, and total leaf area following the application of biogas slurry could be attributed to its high nutrient content, which boosts photosynthesis and promotes the development of plant tissues and organs [35]. Additionally, biogas slurry improves the soil's physical or chemical properties, raising total nitrogen and available phosphorus levels [36], creating a nutrient-rich environment that fosters plant growth. Biogas slurry also accelerates nutrient cycling and influences soil microbial community structure, further promoting plant growth [35]. Combining biogas slurry with NPK fertilizers has been shown to boost plant height. For instance, in okra, a mix of 50% nitrogen from biogas slurry and 50% from NPK fertilizers resulted in a 15% height increase compared to using only NPK [37]. This combination enhances soil nitrogen, phosphorus, and potassium availability, supporting root and overall plant growth [38]. Study suggests that substituting a portion of NPK with biogas slurry can be beneficial; for example, replacing 25% of nitrogen fertilizer with biogas slurry may increase soil nutrients or promote plant growth [38].

Chlorophyll Pigments Content

According to Fig. 2, applying semi-solid biogas slurry (SSBS) at 72 m^3/ha (BS1) significantly decreased chlorophyll pigments (chl. a, b, and total chl.) compared to the recommended NPK dose (control) in both




Fig. 1. Effect of semi-solid biogas slurry at 72 m³/ha (BS1) and 96 m³/ha (BS2), BS1 in combination with recommended dose of NPK (BS1 RD), BS1 in combination with liquid biogas slurry foliar application at 48 L/ha (BS 1 BL), BS2 in combination with recommended dose (BS2 RD), BS2 in combination with liquid biogas slurry foliar application at 48 L/ha (BS2 BL) on plant height a), leaves number b) and leaf area c) in comparing with spinach plants treated with recommended NPK dose (Control: Cont.) during 2022 and 2023 seasons. Various letters on each bar indicate significant difference among the treatments ($P < 0.05$).

seasons. However, no notable differences in chl. a and total chlorophyll were observed with the 96 m³/ha (BS2) application when compared to the control. Combined application of SSBS with either recommended NPK dose or liquid biogas slurry (LBS) foliar application significantly increased chlorophyll pigment levels across both seasons, with the highest levels observed when BS1 and BS2 were combined with the recommended NPK dose. The increase in chlorophyll pigments resulting from SSBS application, combined with the recommended NPK dose, can be attributed to its rich nutrient and organic matter content. When applied alongside NPK fertilizers, SSBS significantly enhances chlorophyll content, essential for photosynthesis and overall plant health [39]. This synergy stems from

a balanced nutrient supply and boosted soil microbial activity, which together improve nutrient absorption and utilization by plants [39].

Both semi-solid and liquid biogas slurry combined applications positively affect chlorophyll content. Liquid biogas slurry provides nutrients in a readily absorbable form, giving plants an immediate nutrient boost [36], while semi-solid slurry offers a slow-release nitrogen source, improving nitrogen uptake over time. This form also enhances soil structure and microbial activity, improving nitrogen availability and plant absorption. The organic matter in semi-solid slurry raises soil cation exchange capacity (CEC), aiding magnesium retention and absorption by plants [40]. Liquid slurry supplies immediately available nitrogen and magnesium, vital




Fig. 2. Effect of semi-solid biogas slurry at 72 m³/ha (BS1) and 96 m³/ha (BS2), BS1 in combination with recommended dose of NPK (BS1 RD), BS1 in combination with liquid biogas slurry foliar application at 48 L/ha (BS1 BL), BS2 in combination with NPK recommended dose (BS2 RD), BS2 in combination with liquid biogas slurry foliar application at 48 L/ha (BS2 BL) on chlorophyll pigments content (chl. a, b and total) in comparing with spinach plants treated with recommended NPK dose (Control: Cont.) during 2022 and 2023 seasons. Various letters on each bar indicate significant difference among the treatments ($P < 0.05$).

for chlorophyll synthesis, supporting rapid nitrogen uptake during critical growth stages. Its liquid form promotes even nutrient distribution in the soil [40]. Nitrogen and magnesium are crucial in chlorophyll molecules; nitrogen aids in amino acid, protein, and enzyme synthesis, all vital for chlorophyll production [41]. Magnesium, as the core atom in chlorophyll, is indispensable for chlorophyll synthesis. It activates enzymes that drive chlorophyll biosynthesis, including those inserting Mg into protoporphyrin IX, a chlorophyll precursor [42].

Vitamin C Content

Fig. 3a) shows that applying both forms of biogas slurry, either individually or in combination with or without the recommended NPK dose, significantly

increased vitamin C content (mg/100 g fresh weight) compared to the control across both seasons. The highest vitamin C content was achieved with semi-solid biogas slurry at 96 m³/ha combined with either the recommended NPK dose (BS2+RD) or liquid biogas slurry (BS2+BL) in both seasons.

The increase in vitamin C content from biogas slurry application may result from its high concentration of essential nutrients and organic matter, which enhance soil fertility and improve crop nutritional quality [43]. Studies have shown that biogas slurry may increase vitamin C levels in fruits; for instance, concentrated biogas slurry significantly boosts vitamin C in tomatoes [44]. Nitrogen, phosphorus, and potassium fertilizers also supply critical nutrients supporting overall plant growth and development, which enhances the synthesis of various metabolites, including vitamin C [45].




Fig. 3. Effect of semi-solid biogas slurry at 72 m³/ha (BS1) and 96 m³/ha (BS2), BS1 in combination with recommended dose of NPK (BS1 RD), BS1 in combination with liquid biogas slurry foliar application at 48 L/ha (BS1 BL), BS2 in combination with NPK recommended dose (BS2 RD), BS2 in combination with liquid biogas slurry foliar application at 48 L/ha (BS2 BL) on vitamin C content a), proline content b) and total phenol c) in comparing with spinach plants treated with recommended NPK dose (Control: Cont.) during 2022 and 2023 seasons. Various letters on each bar indicate significant difference among the treatments ($P<0.05$).

A balanced supply of nitrogen, phosphorus, and potassium is crucial for plant health and vitamin C biosynthesis. Nitrogen, in particular, is essential for synthesizing amino acids and proteins, precursors to vitamin C production [46].

Proline and Leaf Total Phenols Content

The data in Fig. 3b) indicated that all biogas slurry applications, whether applied alone or in combination with the recommended NPK dose, resulted in a reduction of proline content compared to the control in both seasons. The lowest proline levels were observed with the application of semi-solid biogas slurry at 96 m³/ha, either combined with the recommended NPK dose or

with foliar application of liquid biogas slurry, during both seasons. Fig. 3c) shows that applying both forms of biogas slurry, either individually or in combination with each other or with the recommended NPK dose, significantly increased total phenols content in spinach leaves compared to the control treatment across each season. The highest leaf total phenols content was observed with the application of semi-solid biogas slurry at 96 m³/ha combined with the recommended NPK dose in both seasons.

The increase in total phenol content in spinach leaves from applying biogas slurry, whether individually or in combination with each other or the recommended NPK dose, may be attributed to the nutrients in biogas slurry that facilitate the synthesis of these compounds




Fig. 4. Effect of semi-solid biogas slurry at 72 m³/ha (BS1) and 96 m³/ha (BS2), BS1 in combination with recommended dose of NPK (BS1 RD), BS1 in combination with liquid biogas slurry foliar application at 48 L/ha (BS1 BL), BS2 in combination with NPK recommended dose (BS2 RD), BS2 in combination with liquid biogas slurry foliar application at 48 L/ha (BS2 BL) on nitrogen, phosphorous and potassium% in spinach plant in comparing with control during 2022 and 2023 seasons. Various letters on each bar indicate significant difference among the treatments ($P < 0.05$).

[10]. Research shows that NPK fertilizers can also boost total phenolic content in plants; for example, a study on purslane (*Portulaca grandiflora*) found that NPK fertilizer significantly increased total phenolic content in the leaves [47]. A balanced supply of nitrogen, phosphorus, or potassium is crucial for maintaining optimal plant health, which is essential for biosynthesis of phenolic compounds. Nitrogen, in particular, plays a vital role in synthesizing amino acids and proteins, which are precursors to phenolic production [48]. The combination of NPK and biogas slurry creates a more favorable environment for plant growth, promoting increased production of secondary metabolites like phenolic compounds, which are important for plant defense and antioxidant activity [49].

Percentage of Nitrogen, Phosphorus, and Potassium

The data in Fig. 4 shows that applying semi-solid biogas slurry (SSBS) at 72 m³/ha had no significant effect on the nitrogen, phosphorus, and potassium (N, P, K) percentages in spinach plants compared to the control in both seasons. However, a significant increase in N, P, K% was observed when SSBS at 72 m³/ha was combined with the recommended NPK dose or liquid biogas slurry (LBS) foliar application at 48 L/ha in each season. Likewise, SSBS at 96 m³/ha, either alone or combined with the recommended NPK dose or LBS, significantly raised N, P, and K percentages in both seasons. The highest N, P, K% values were achieved with SSBS at 72 m³/ha combined with the recommended NPK dose,




Fig. 5. Effect of semi-solid biogas slurry at 72 m³/ha (BS1) and 96 m³/ha (BS2), BS1 in combination with recommended dose of NPK (BS1 RD), BS1 in combination with liquid biogas slurry foliar application at 48 L/ha (BS1 BL), BS2 in combination with NPK recommended dose (BS2 RD), BS2 in combination with liquid biogas slurry foliar application at 48 L/ha (BS2 BL) on Fe, Cu and Zn content in spinach plant in comparing with control during 2022 and 2023 seasons. Various letters on each bar indicate significant difference among the treatments ($P < 0.05$).

as well as with SSBS at 96 m³/ha combined with either the recommended NPK dose or LBS foliar application in both seasons. The increase in essential nutrients like nitrogen (N), phosphorus (P), and potassium (K) is vital for plant growth. Through anaerobic digestion in biogas production, these nutrients are mineralized, making them more accessible to plants [50]. Applying biogas slurry (BGS) can enhance soil properties by raising pH, cation exchange capacity (CEC), and levels of bioactive substances like humic acids, leading to improved nutrient retention or availability in soil [51, 52]. Additionally, BGS can partially replace synthetic fertilizers, thus lowering both environmental impact and

the costs associated with chemical fertilizers, making it a sustainable choice for nutrient management in agriculture [53].

Iron (Fe), Copper (Cu), Zinc (Zn), Lead (Pb), and Cadmium (Cd) Content

Fig. 5 demonstrates that applying semi-solid biogas slurry (SSBS) at 72 and 96 m³/ha, combined with the recommended NPK dose, significantly increased Fe content compared to the control across each season. The highest Fe content was achieved with SSBS at 96 m³/ha combined with NPK during both seasons.




Fig. 6. Effect of semi-solid biogas slurry at 72 m³/ha (BS1) and 96 m³/ha (BS2), BS1 in combination with recommended dose of NPK (BS1 RD), BS1 in combination with liquid biogas slurry foliar application at 48 L/ha (BS1 BL), BS2 in combination with NPK recommended dose (BS2 RD), BS2 in combination with liquid biogas slurry foliar application at 48 L/ha (BS2 BL) on Pb and Cd content in spinach plant in comparing with control during 2022 and 2023 seasons. Various letters on each bar indicate significant difference among the treatments ($P < 0.05$).

Conversely, all treatments led to a reduction in Cu or Zn content compared to the control. The lowest Cu and Zn levels were observed with SSBS at 96 m³/ha combined with LGS at 48 L/ha as a foliar application and with SSBS at 72 m³/ha, respectively, across both seasons. Fig. 6 shows that all treatments reduced Pb and Cd content compared to the control across both seasons. The lowest Pb and Cd levels were achieved with the application of SSBS at 96 m³/ha combined with 48 L/ha as a foliar application in both seasons. Biogas slurry is nutrient-rich, especially in iron, which plants readily absorb. This application not only raises Fe content but also enhances soil health and fertility overall [54]. Biogas slurry increases iron bioavailability in the soil, facilitating plant uptake of this essential nutrient [54]. However, while biogas slurry typically enhances nutrient availability, it may sometimes reduce Cu and Zn absorption by plants. Biogas slurry can alter soil pH, potentially decreasing the bioavailability of Cu and Zn. Higher pH levels can cause these metals to precipitate, reducing plant absorption [55]. Additionally, excessive biogas slurry can disrupt soil nutrient balance, potentially hindering the uptake of certain micronutrients like Cu and Zn [10].

Nitrate, Nitrite Content, and Nitrate Reductase Activity

Fig. 7a) data showed that applying biogas slurry (BS), in various forms – whether individually, in combinations, or with the recommended NPK dose decreased nitrate content compared to using only the NPK recommended dose across both seasons. The lowest nitrate content was observed with the foliar application of SSBS at 96 m³/ha, either alone or combined with the application of liquid biogas slurry in both seasons.

Data in Fig. 7b) showed that all biogas slurry treatments, whether used alone, in combination with each other, or with the recommended NPK dose, increased nitrite content compared to the control, except for the combination of SSBS at 96 m³/ha with foliar application of liquid biogas slurry in both seasons. The highest nitrite content was recorded with the SSBS application at 72 m³/ha across both seasons. The data in Fig. 7c) demonstrated that all treatments involving biogas slurry, whether applied alone or with the recommended NPK dose, significantly enhanced nitrate reductase activity in both seasons. The highest nitrate reductase activity in both seasons was observed




Fig. 7. Effect of semi-solid biogas slurry at 72 m³/ha (BS1) and 96 m³/ha (BS2), BS1 in combination with recommended dose of NPK (BS1 RD), BS1 in combination with liquid biogas slurry foliar application at 48 L/ha (BS1 BL), BS2 in combination with NPK recommended dose (BS2 RD), BS2 in combination with liquid biogas slurry foliar application at 48 L/ha (BS2 BL) on nitrate a) and nitrite b) content as well as nitrate reductase activity c) in spinach plant in comparing with control during 2022 and 2023 seasons. Various letters on each bar indicate significant difference among the treatments ($P < 0.05$).

with the application of semi-solid biogas slurry at 96 m³/ha, either on its own or in combination with foliar application of liquid biogas slurry.

Oxalate Content and Yield

The data in Fig. 8a) indicate that applying SSBS at 72 or 96 m³/ha, either alone or in combination with the recommended NPK dose or LBS at 48 L/ha as a foliar spray, reduced oxalate content in spinach leaves compared to the control in both growing seasons. The lowest oxalate levels were observed with the individual application of SSBS at 72 m³/ha during both seasons. Regarding spinach yield (mg. ha⁻¹), SSBS at 72 and 96 m³/ha⁻¹ as a soil application in combination

with NPK recommended dose and SSBS at 96 m³/ha⁻¹ resulted in the highest total yield per ha during both seasons (Fig. 8b)). Application of biogas slurry (BGS) may have various effects on plant growth or soil health, although its specific impact on oxalate content in plants is less commonly explored. Biogas slurry is rich in nutrients such as nitrogen, phosphorus, and potassium, which can promote plant growth or enhance nutrient uptake [10]. This increased nutrient availability may influence metabolic processes in plants, potentially affecting oxalate synthesis. The reduction in oxalate content following BGS application could be attributed to several factors. First, BGS provides a balanced supply of essential nutrients, which can optimize plant metabolism and reduce oxalate production, as oxalates are often




Fig. 8. Effect of semi-solid biogas slurry at 72 m³/ha (BS1) and 96 m³/ha (BS2), BS1 in combination with recommended dose of NPK (BS1 RD), BS1 in combination with liquid biogas slurry foliar application at 48 L/ha (BS1 BL), BS2 in combination with NPK recommended dose (BS2 RD), BS2 in combination with liquid biogas slurry foliar application at 48 L/ha (BS2 BL) oxalate content a) and total yield b) in spinach plant in comparing with control during 2022 and 2023 seasons. Various letters on each bar indicate significant difference among the treatments ($P < 0.05$).

synthesized in response to nutrient deficiencies. Second, the organic matter in BGS enhances soil microbial activity, improving nutrient availability and uptake by plants. This, in turn, can result in healthier plants with lower oxalate levels.

The application of BS to soil can significantly affect the nitrate and nitrite levels in plants. Rich in nitrogen, biogas slurry boosts nitrate content in soil, which in turn enhances the plant's ability to absorb nitrate [56]. When semi-solid biogas slurry (SSBS) is combined with foliar application of liquid biogas slurry, it notably reduces nitrite content, as biogas slurry often contains beneficial microbes that facilitate the conversion of nitrite into nitrate through a process known as nitrification [56]. Furthermore, biogas slurry typically increases nitrate reductase activity in plants. This enzyme is essential for converting nitrate to nitrite during nitrogen assimilation, allowing plants to utilize nitrogen more efficiently, thereby improving growth and boosting yields [10, 35]. The application of biogas slurry (BS) can influence proline content in plants. By reducing stress levels, plants may produce lower amounts of proline, which is typically synthesized in response to stress [10, 35].

Soil Bulk Density, Total Porosity, and Organic Matter Content

The data in Fig. 9 indicate that the soil application of SSBS at 72 and 96 m³/ha, either alone or in combination with the recommended NPK dose or foliar application of liquid biogas slurry, resulted in a reduction in soil bulk density compared to the control treatment in both seasons. Conversely, these treatments enhanced total porosity or organic matter content of the soil during both seasons.

Application of biogas slurry has been shown to significantly affect soil bulk density, improving soil structure by making it less compact and more porous [49]. This reduction in bulk density also enhances the soil's water-holding capacity, increases organic matter content, and boosts nutrient availability [11]. Additionally, biogas slurry promotes the formation of soil aggregates, which is vital for sustaining soil fertility, creating macro-pores that facilitate water movement, root penetration, and gas exchange [49]. The slurry's organic matter content plays a key role in improving soil total porosity by creating more pore spaces, influencing both macro- and micro-porosity within the soil matrix, which in turn enhances soil aeration and water




Fig. 9. Soil bulk density a), total porosity b), and organic matter content c) as affected by application of semi-solid biogas slurry at 72 m³/ha (BS1) and 96 m³/ha (BS2), BS1 in combination with recommended dose of NPK (BS1 RD), BS1 in combination with liquid biogas slurry foliar application at 48 L/ha (BS1 BL), BS2 in combination with NPK recommended dose (BS2 RD), BS2 in combination with liquid biogas slurry foliar application at 48 L/ha (BS2 BL) at harvest for the 2022 and 2023 growing seasons. Various letters on each bar indicate significant difference among the treatments ($P < 0.05$).

infiltration [57]. While biogas slurry generally increases organic matter content in soil, the extent of this increase can vary. Some studies suggest that the organic matter from biogas slurry may be rapidly mineralized, which could limit long-term increases in soil organic matter [58, 59]. The utilization of organic manure, such as biogas slurry, significantly affects soil properties, including cation exchange capacity (CEC), pH, organic carbon content (OC%), and nutrient availability.

Soil pH and Electrical Conductivity (EC)

The data in Fig. 10a) show that the application of SSBS, either alone or in combination with the recommended NPK dose or LBS foliar application, significantly reduced soil pH in both seasons. Regarding soil electrical conductivity (EC), all treatments resulted

in a decrease in EC compared to the control in both seasons (Fig. 10b)). The changes in organic matter and pH caused by biogas slurry may lead to the formation of complexes with heavy metals, reducing their solubility and availability for plant absorption [60]. Increased microbial activity stimulated by biogas slurry may also promote sequestration of heavy metals by soil microbes, further limiting their uptake by plants [60]. Additionally, biogas slurry can raise the soil's overall nutrient content, potentially diluting Pb and Cd concentrations and making them less accessible for plant absorption [10].

Soil Mineral Nutrient Content

Macronutrients such as nitrogen (N), phosphorus (P), potassium (K) (Fig. 11), as well as micronutrients




Fig. 10. Soil pH a) and electrical conductivity EC b) as affected by application of semi-solid biogas slurry at 72 m³/ha (BS1) and 96 m³/ha (BS2), BS1 in combination with recommended dose of NPK (BS1 RD), BS1 in combination with liquid biogas slurry foliar application at 48 L/ha (BS1 BL), BS2 in combination with NPK recommended dose (BS2 RD), BS2 in combination with liquid biogas slurry foliar application at 48 L/ha (BS2 BL) at harvest for the 2022 and 2023 growing seasons. Various letters on each bar indicate significant difference among the treatments ($P < 0.05$).

like iron (Fe), copper (Cu), or zinc (Zn) (Fig. 12), were significantly elevated in the soil following the application of SSBS, either alone or in combination with the recommended NPK dose or LBS as a foliar spray, at the harvest date during both seasons. Organic manure (OM) is a valuable source of essential nutrients like nitrogen, phosphorus, potassium, micronutrients, and organic matter, as noted [61]. During decomposition and mineralization, these nutrients are gradually released into the soil [62]. This slow release provides a steady supply of plant-available nutrients throughout the growing season, supporting crop growth and development [63]. Additionally, OM enhances microbial activity and improves soil structure, which promotes nutrient cycling and uptake by plants [64]. Thus, organic manure improves soil fertility, increases nutrient availability, and reduces dependence on synthetic fertilizers, contributing to sustainable agriculture.

Organic manure can also affect soil pH, though its impact is less significant compared to liming or acidic amendments. The pH of organic manure varies based on its composition and stage of decomposition, typically ranging from acidic to neutral or slightly alkaline. Organic manure acts as a buffer, helping to moderate pH changes and stabilize soil acidity [61]. Generally, less decomposed organic matter may have a slightly acidic

pH due to organic acids, while well-decomposed matter tends to be neutral or slightly alkaline [64].

Biological characteristics of soil are significantly impacted by organic manure, which improves microbial diversity, abundance, activity, and general soil health. Organic manure supports nutrient cycling, organic matter turnover, and soil ecosystem functioning by giving soil microorganisms a rich source of organic matter, nutrients, and energy. This encourages the growth of beneficial microbial communities. Improved soil structure, nutrient availability, and disease suppression are all facilitated by increased microbial diversity and activity, which also increases soil fertility, health, and resistance to environmental pressures. Maintaining soil biological qualities, bolstering soil health, and advancing agricultural sustainability in the face of persistent environmental concerns worldwide all depend on sustainable soil management techniques that incorporate the application of organic manure [65]. Biogas slurry can influence both soil pH and electrical conductivity (EC). When applied to soil, it often leads to a decrease in pH, making the soil more acidic. However, biogas slurry can also help neutralize acidic soils, resulting in an increase in soil pH, which is beneficial for crops that thrive in neutral to slightly alkaline conditions [66]. In some cases, though,




Fig. 11. Nitrogen a), phosphorous b) and potassium c) content in soil as affected by application of semi-solid biogas slurry at 72 m³/ha (BS1) and 96 m³/ha (BS2), BS1 in combination with recommended dose of NPK (BS1 RD), BS1 in combination with liquid biogas slurry foliar application at 48 L/ha (BS1 BL), BS2 in combination with NPK recommended dose (BS2 RD), BS2 in combination with liquid biogas slurry foliar application at 48 L/ha (BS2 BL) at harvest for the 2022 and 2023 growing seasons. Various letters on each bar indicate significant difference among the treatments ($P < 0.05$).

application of biogas slurry has not significantly altered soil pH, suggesting that its effect can depend on initial soil conditions and the quantity of slurry applied [58]. The reduction in EC indicates a decrease in soil salinity, which can improve soil health and promote better plant growth. In certain studies, biogas slurry did not lead to a significant increase in EC, suggesting that its application can be managed without causing soil salinization [58]. This is important for maintaining soil health and preventing negative impacts on crop growth.

The application of biogas slurry may significantly enhance the mineral nutrient content of soil. Its application can increase the availability of these nutrients in soil, which are crucial for plant growth [52]. Organic matter in biogas slurry can improve soil structure or increase the soil's capacity to retain water and nutrients [67]. Biogas slurry may boost microbial activity in the soil, which helps in the decomposition of

organic matter and the release of nutrients [66]. This can lead to better nutrient cycling or availability for plants.

Soil Cadmium and Lead Content

Data in Fig. 13 show that applying SSBS at 72 m³/ha, either alone or combined with the recommended NPK dose, significantly increased lead content in soil during both growing seasons. Additionally, the application of SSBS at 96 m³/ha combined with the NPK recommended dose significantly raised lead soil content during the second season. Regarding cadmium (Cd) soil levels, SSBS at 96 m³/ha combined with the NPK recommended dose, or LBS applied as a foliar treatment at 48 L/ha, significantly increased Cd content in the soil during both seasons.

The use of biogas slurry (BGS) on soil can influence its properties, including levels of heavy metals like lead




Fig. 12. Iron a), copper b) and zinc c) content in soil as affected by application of semi-solid biogas slurry at 72 m³/ha (BS1) and 96 m³/ha (BS2), BS1 in combination with recommended dose of NPK (BS1 RD), BS1 in combination with liquid biogas slurry foliar application at 48 L/ha (BS1 BL), BS2 in combination with NPK recommended dose (BS2 RD), BS2 in combination with liquid biogas slurry foliar application at 48 L/ha (BS2 BL) at harvest for the 2022 and 2023 growing seasons. Various letters on each bar indicate significant difference among the treatments ($P<0.05$).

(Pb) or cadmium (Cd). Depending on the feedstock used in biogas production, biogas slurry may contain these heavy metals, which can accumulate in the soil when applied. This accumulation can potentially raise the concentrations of Pb or Cd [59]. However, despite an increase in soil's heavy metal content, both before (Table 1) and after planting (Fig. 13), the application of biogas slurry resulted in a decrease in buildup of these metals in tissues of spinach plants (Fig. 6).

The best organic matter for stabilizing heavy metals, lowering heavy metal uptake, and encouraging plant growth in contaminated soil was found to have a favorable impact on plant and soil characteristics, indicating its potential for safe crop production. With the benefits of being inexpensive and readily available, organic manure (OM), a common organic fertilizer, may improve soil quality and increase crop yields. According

to Huang et al. [68], organic fertilizer can also lessen the toxicity of heavy metals by complexing and adsorbing them. Functional elements like hydroxyl and carboxyl groups found in organic components in manure mix with heavy metals to produce organic-metal complexes, which change the availability and movement of heavy metals [69-72]. *Bidens tripartite* biomass rose, and plant Cd content fell in Cd-contaminated soil treated with organic manure [73, 74]. Guo et al [75] found that manure enhanced phytoextraction of heavy metals by *Streptomyces pactum* in a Cd-contaminated soil. Heavy metal ions are strongly adsorbed and chelated by organic matter and beneficial microorganisms found in organic fertilizer. Because there were fewer water-soluble and exchangeable heavy metals in the soil, the bioavailability decreased [76].




Fig. 13. Lead a) and cadmium b) in soil as affected by application of semi-solid biogas slurry at 72 m³/ha (BS1) and 96 m³/ha (BS2), BS1 in combination with recommended dose of NPK (BS1 RD), BS1 in combination with liquid biogas slurry foliar application at 48 L/ha (BS1 BL), BS2 in combination with NPK recommended dose (BS2 RD), BS2 in combination with liquid biogas slurry foliar application at 48 L/ha (BS2 BL) at harvest for the 2022 and 2023 growing seasons. Various letters on each bar indicate significant difference among the treatments ($P < 0.05$).




Fig. 14. Diagram illustrating the impact of semi-solid biogas slurry (SSBS) soil application at rates of 72 and 96 m³/ha, either alone or combined with the recommended NPK dose, or liquid biogas slurry (LBS) applied as a foliar spray at 48 L/ha. The treatments positively influenced parameters such as plant height, dry weight, leaf area, and leaf number per plant, as well as chlorophyll pigment content, vitamin C, total phenols in leaves, and nitrate reductase activity. Additionally, reductions were observed in cadmium (Cd), lead (Pb), proline, nitrate, and nitrite contents, alongside improved yield per hectare.

Ultimately, because organic fertilizers boost soil fertility, sustainability, and crop yields in a synergistic way, they are frequently used in China's intensive crop farming systems in place of chemical fertilizers [77]. However, total substitution of organic fertilizers for chemical fertilizers can frequently result in immediate lack of nutrient availability, lowering crop yields, because of the sluggish release of nutrients and inefficiency of organic fertilizers. According to a meta-analysis research, where the replacement ratio was 43% or less, using manure instead of chemical fertilizer did not lower wheat yields [78].

Graphical Summary

The diagram (Fig. 14) highlights the beneficial effects of applying semi-solid biogas slurry soil (SSBS) at rates of 72 and 96 m³/ha, either alone or in combination with the recommended NPK dose or liquid biogas slurry (LBS), on spinach plants. These treatments significantly improved growth metrics, including plant height, dry weight, leaf area, and the number of leaves per plant. They also enhanced chlorophyll pigment content (chlorophyll a, b, and total), vitamin C levels, total phenols in leaves, and nitrate reductase activity. Furthermore, these treatments reduced concentrations of cadmium (Cd), lead (Pb), proline, nitrate, or nitrite in spinach plants across both growing seasons.

Conclusions

The results demonstrated that applying semi-solid biogas slurry, whether used alone, alongside recommended rates of nitrogen, phosphorus, and potassium, or in combination with a foliar spray of liquid biogas slurry, significantly enhanced spinach growth, yield, and nutritional quality. Additionally, the use of biogas slurry individually or combined with other treatments reduced concentrations of cadmium and lead in spinach leaves and improved soil chemical characteristics, including nutrient availability. These findings indicate that biogas slurry has the potential to lessen reliance on organic fertilizers, thereby contributing to more sustainable agricultural practices.

Acknowledgements

We thank the Deanship of Scientific Research, Vice Presidency for Graduate Studies and Scientific Research, King Faisal University, Saudi Arabia (KFU242525), for supporting this research work.

Conflict of Interest

The authors declare that they have no conflicts of interest.

References

1. RABIE K.A.E., ASHOUR H.M., MANAF H.H. Influence of nitrogen forms on nitrate and nitrite accumulation in the edible parts of spinach (*Spinacia oleracea*, L.) plant with maintenance for yield production. *Journal of Horticultural Science & Ornamental Plants*. **6** (3), 126, **2014**.
2. CAO H., JIANG Y., CHEN J., ZHANG H., HUANG W., LI L., ZHANG W. Arsenic accumulation in *Scutellaria baicalensis* Georgi and its effects on plant growth and pharmaceutical components. *Journal of Hazardous Materials*. **171**, 508, **2009**.
3. DALVI A.A., BHALERAO S.A. Response of plants towards heavy metal toxicity: an overview of avoidance, tolerance and uptake mechanism. *Annals of Plant Sciences*. **2** 362, **2013**.
4. POPOOLA O.J., OGUNDELE O.D., LADAPPO E.A., SENBORE S. The impact of heavy metal contamination in soils on soil microbial communities and its potential health risks for humans. In: *Soil Microbiome in Green Technology Sustainability*. Aransiola S.A., Atta H.I., Maddela N.R., Eds., Cham: Springer Nature Switzerland. **15**, 351, **2024**.
5. GOYAL D., YADAV A., PRASAD M., SINGH T.B., SHRIVASTAV P., ALI A., DANTU P.K., MISHRA S. Effect of heavy metals on plant growth: An Overview. In: *Contaminants in Agriculture*. Naeem, M., Ansari, A., Gill, S. (eds). Cham: Springer Nature Switzerland. **4**, 78, **2020**.
6. GILL R.A., KANWAR M.K., RODRIGUES R.A., ALI B. Heavy metal toxicity in plants: recent insights on physiological and molecular aspects. *Frontiers in Plant Science*. **12**, 830682, **2022**.
7. PANDEY A.K., ZORIĆ L., SUN T., KARANOVIC D., FANG P., BORISEV M., WU X., LUKOVIC J., XU P. The anatomical basis of heavy metal responses in legumes and their impact on plant–rhizosphere interactions. *Plants*. **11** (19), 2554, **2022**.
8. KHAN A., KHAN S., KHAN M.A., QAMAR Z., WAQAS M. The uptake and bioaccumulation of heavy metals by food plants, their effects on plants nutrients, associated health risk: a review. *Environmental Science and Pollution Research*. **22**, 13772, **2015**.
9. TIAN Y., WU Y., PENG Y., GUO X., LI Y., DAI B., HUANG T. Study on the complexation of heavy metals onto biogas slurry DOM using two-dimensional correlation spectroscopy combined with the log-transformed synchronous fluorescence spectroscopy. *Environmental Science and Pollution Research*. **28**, 22878, **2021**.
10. KUMAR A., VERMA L.M., SHARMA S., SINGH N. Overview on agricultural potentials of biogas slurry (BGS): applications, challenges, solutions. *Biomass Conversion and Biorefinery*. **13**, 13729, **2023**.
11. TANG J., DAVY A.J., WANG W., ZHANG X., WU D., HU L., YIN J. Effects of biogas slurry on crop yield, physicochemical properties and aggregation characteristics of lime concretion soil in wheat–maize rotation in the North China Plain. *Journal of Soil Science and Plant Nutrition*. **22** (2), 2406, **2022**.
12. LIU Q., ZHAO Z., XUE Z., LI D., WEN Z., RAN Y., MEI X., HE L. Comprehensive risk assessment of applying biogas slurry in peanut cultivation. *Frontiers in Nutrition*. **8**, 702096, **2021**.
13. SPARKS D.L., PAGE A.L., HELMKE P.A., LOEPPERT R.H. *Methods of soil analysis, part 3: Chemical methods*. John Wiley & Sons, pp. 417-435, **2020**.

14. DANE J.H., TOPP C.G. Methods of soil analysis. Part 4: Physical methods, John Wiley & Sons, pp. 1-200, **2020**.
15. MORAN R. Formulae for determination of chlorophyllous pigments extracted with N, N-dimethylformamide. *Plant Physiology*. **69** (6), 1376, **1982**.
16. BATES L.S., WALDREN R.P., TEARE I.D. Rapid determination of free proline for water stress studies. *Plant and Soil*. **39** (1), 205, **1973**.
17. BESSADA S.M., BARREIRA J.C., BARROS L., FERREIRA J.C., OLIVEIRA M.B.P. Phenolic profile and antioxidant activity of *Coleostephus myconis* (L.) rchb. F.: an underexploited and highly disseminated species. *Industrial Crops and Products*. **89**, 45, **2016**.
18. PETERBURGSKI A.V. Handbook of Agronomic Chemistry. Kolos Publishing House, Moscow, pp. 29-86, **1968** [in Russian].
19. WALINGA I., VAN D.E., RLEE J.J., HOUBA V.J., VAN VARK W., NOVOZAMSKY I. Plant analysis manual. Springer Dordrecht, Netherlands. **2012**.
20. Chen Y., JIANG Z., ZHU R. A study of the surface phase change of zinc electrode under anodic passivation in KOH solutions. *Acta Chimica Sinica*. **41**(3), 193, **1983**.
21. BETTINELLI M., BEONE G.M., SPEZIA S., BAFFI C. Determination of heavy metals in soils and sediments by microwave-assisted digestion and inductively coupled plasma optical emission spectrometry analysis. *Analytica Chimica Acta*. **424**, 289, **2000**.
22. OZDESTAN O., UREN A. Development of a costeffective method for nitrate and nitrite determination in leafy plants and nitrate and nitrite contents of some green leafy vegetables grown in the Aegean region of Turkey. *Journal of Agricultural and Food Chemistry*. **58** (9), 5235, **2010**.
23. PALANISWAMY U.R., BIBLE B.B., MCAVOY R.J. Oxalic acid concentrations in purslane (*Portulaca oleraceae* L.) is altered by the stage of harvest and the nitrate to ammonium ratios in hydroponics. *Scientia Horticulturae*. **102**, 267, **2004**.
24. KAISER W.M., BRENDLE-BEHNISCH E. Rapid modulation of spinach leaf nitrate reductase activity by photosynthesis. I. Modulation in vivo by CO₂ availability. *Plant Physiology*. **96**, 363, **1991**.
25. HAGEMAN R.H., REED A.J. Nitrate reductase from higher plants. *Methods Enzymology*, **69**, 270, **1980**.
26. ALLEN O.N. Experiments in soil Bacteriology. University of Minnesota, Minneapolis: Burgess Pub. Co., 3rd edition, **1959**.
27. MCLEAN E.O. Soil pH and lime requirement. In: Methods of soil analysis, part 2, chemical and microbiological properties, 2nd ed.; A. L. Page, (eds.) American Soc. Agron. Inc, Madison, w1, USA. **12**, 199, **1982**.
28. PAGE A.L., MILLER R.H., KEENEY D.R. Methods of Soil Analysis. Part II: Chemical and microbiological properties, 2nd ed. American Society of Agronomy. In Soil Science Society of America, Madison, USA, **1982**.
29. KIM Y.W., PETROV I., GREENE J.E., ROSSNAGEL S.M. Development of 111 textures in Al films grown on SiO₂/Si (001) by ultrahigh-vacuum primary-ion deposition. *Journal of Vacuum Science and Technology*. **14** (2), 346, **1996**.
30. BHATTACHARYA T., CHANDRAN P., RAY S.K., MANDAL C., TIWARY P., PAL D.K., MAURYA U.K., NIMKAR A.M., KUCHANKAR H., SHEIKH S., TELPANDE B.A., KOLHE A. Walkley-Black recovery factor to reassess soil organic matter: Indogangetic plains and black soil region of India case studies. *Communications in Soil Science and Plant Analysis*. **46** (20), 2628, **2015**.
31. CAMPBELL D.J. Determination and use of soil bulk density in relation to soil compaction. *Developments in Agricultural Engineering*. **11**, 113, **1994**.
32. KLUTE A., DIRKSON C. Hydraulic conductivity and diffusivity: Laboratory methods. In: Methods of soil analysis: part 1-physical and mineralogical methods. American Society of Agronomy, Inc. **1986**.
33. JACKSON M.L. Soil chemical analysis: advanced course: a manual of methods useful for instruction and research in soil chemistry, physical chemistry of soils, soil fertility, soil genesis. 2nd ed, Madison, Wis.: Parallel Press, University of Wisconsin-Madison Libraries, **2005**.
34. GOMEZ K.A., GOMEZ A.A. Statistical procedures for agricultural research. 2nd ed, John Wiley and Sons, Inc, New York, p. 680, **1984**.
35. CAI J., RAN Q., ZHOU J., BI M., LIU Y., YANG S., FAN Y., NIE G., HE W. Effects of arbuscular mycorrhizal fungi and biogas slurry application on plant growth, soil composition, microbial communities of hybrid Pennisetum. *Sustainability*. **16** (19), 8416, **2024**.
36. YIN D., CHEN D., QIN J., LIU W., WANG Y., LAN W., ZHONG N., MA P. Effect of drip irrigation with biogas slurry on the yield and quality of Chinese Cabbage. *Bio Resources*. **19** (3), 5717, **2024**.
37. MAQBOOL S., UL HASSAN A., JAVEDAKHTAR M., TAHIR M. Integrated use of biogas slurry and chemical fertilizer to improve growth and yield of okra. *Science Letters*. **2** (1), 56, **2014**.
38. LU Y., XIAO Q., WU S., YUAN H., GAO T., CAI T., WU X., MA Y., LIAO X. Partial substitution of nitrogen fertilizer with biogas slurry increases rice yield and fertilizer utilization efficiency, enhancing soil fertility in the Chaohu lake basin. *Plants*. **13** (15), **2024**].
39. ZHANG H., LI S., ZHENG X., ZHANG J., BAI N., ZHANG H., LV W. Effects of biogas slurry combined with chemical fertilizer on soil bacterial and fungal community composition in a paddy field. *Frontiers Microbiology*. **12**, 65515, **2021**.
40. LI M., LIU Y., LUO L., YING S., JIANG P. Effects of different biogas slurry application patterns on nitrogen and phosphorus losses in a paddy field. *Paddy and Water Environment*. **22**, 521, **2024**.
41. ISHFAQ M., WANG Y., YAN M., WANG Z., WU L., LI C., LI X. Physiological essence of magnesium in plants and its widespread deficiency in the farming system of China. *Frontiers of Plant Science*. **13**, 802274, **2022**.
42. TRIPATHY B.C., PATTANAYAK G.K. Chlorophyll biosynthesis in higher plants. In: Photosynthesis: plastid biology, energy conversion and carbon assimilation, Springer, pp. 63-94, **2012**.
43. EL-BELTAGI H.S., AHMAD I., BASIT A., SHEHATA W.F., HASSAN U., SHAH S.T., HALEEMA B., JALAL A., AMIN R., KHALID M.A., NOOR F., MOHAMED H.I. Ascorbic acid enhances growth and yield of sweet peppers (*Capsicum annum*) by mitigating salinity stress. *Gesunde Pflanzen*. **74**, 423, **2022**.
44. YU F.B., LUO X.P., SONG C.F., ZHANG M.X., SHAN S.D. Concentrated biogas slurry enhanced soil fertility and tomato quality. *Acta Agriculturae Scandinavica Section B*. **60** (3), 262, **2010**.
45. SUN J., LUO H., JIANG Y., WANG L., XIAO C., WENG L. Influence of nutrient (NPK) factors on growth, pharmacodynamic component biosynthesis of *Atractylodes chinensis*: An insight on acetyl-CoA carboxylase (ACC), 3-hydroxy-3-methylglutaryl-CoA reductase (HMGR),

- farnesyl pyrophosphate synthase (FPPS) signaling responses. *Frontiers in Plant Science*. **13**, 799201, **2022**.
46. GHIFARI A.F., KOESRIHARTI M.R. Influence of NPK fertilizer doses on yield and vitamin C content of two varieties of amaranth (*Amaranthus tricolor* L.). *Jurnal Produksi Tanaman*. **7**, 1780, **2019**.
 47. PUTRA R.P., MEDIARTHA I.K., SETIAWAN M.A.D., SUJAI P.A.N., ARISTA R.A., KANDI R.P., AISYAH S.L., NURCHOLIS W. Effects of NPK fertilizer on growth, phytochemical content and antioxidant activity of Purslane (*Portulaca grandiflora*). *Current Applied Science and Technology*. **24**, e0257237, **2024**.
 48. SIDDIQUI Y., MUNUSAMY U., NAIDU Y., AHMAD K. Integrated effect of plant growth-promoting compost and NPK fertilizer on nutrient uptake, phenolic content, antioxidant properties of *Orthosiphon stamineus* and *Cosmos caudatus*. *Horticulture, Environment, and Biotechnology*. **61**, 1051, **2020**.
 49. ZHENG X., FAN J., XU L., ZHOU J. Effects of combined application of biogas slurry and chemical fertilizer on soil aggregation and C/N distribution in an Ultisol. *PLoS One*. **12** (1), e0170491, **2017**.
 50. XU W., ZHU Y., WANG X., JI L., WANG H., YAO L., LIN C. The effect of biogas slurry application on biomass production and forage quality of *Lolium multiflorum*. *Sustainability*. **13** (7), 3605, **2021**.
 51. XU M., XIAN Y., WU J., GU Y., YANG G., ZHANG X., PENG H., YU X., XIAO Y., LI L. Effect of biogas slurry addition on soil properties, yields, bacterial composition in the rice-rape rotation ecosystem over 3 years. *Journal of Soils and Sediments*. **19**, 2534, **2019**.
 52. NIYUNGEKO C., LIANG X., LIU C., ZHOU J., CHEN L., LU Y., TIIMUB B.M., LI F. Effect of biogas slurry application on soil nutrients, phosphomonoesterase activities, phosphorus species distribution. *Journal of Soils and Sediments*. **20**, 900, **2020**.
 53. YADAV R., SUDHISHRI S., KHANNA M., LAL K., DASS A., KUSHWAHA H.L., BANDYOPADHYAY K., DEY A., KUSHWAH A., NAG R.H. Temporal characterization of biogas slurry: a pre-requisite for sustainable nitrification in crop production. *Frontiers in Sustainable Food Systems*. **7**, 1234472, **2023**.
 54. REN T., LIAO J., JIN L., DELGADO-BAQUERIZO M., RUAN H. Application of biogas-slurry and biochar improves soil multifunctionality in a poplar plantation during afforestation processes. *Plant and Soil*. **501** (1), 107, **2024**.
 55. WIN A.T., TOYOTA K., WIN K.T., MOTOBAYASHI T., OOKAWA T., HIRASAWA T., CHEN D., LU J. Effect of biogas slurry application on CH₄ and N₂O emissions, Cu and Zn uptakes by whole crop rice in a paddy field in Japan. *Journal of Soil Science and Plant Nutrition*. **60** (3), 411, **2014**.
 56. LEI J., SUN Y., YIN J., LIU R., CHEN Q. Influence of biogas slurry and a nitrification inhibitor application in nitrous oxide emissions by soil. *Agronomy*. **14** (8), 1844, **2024**.
 57. DU Z., CHEN X., QI X., LI Z., NAN J., DENG J. The effects of biochar and hogger biogas slurry on fluvo-aquic soil physical and hydraulic properties: a field study of four consecutive wheat–maize rotations. *Journal of Soils and Sediments*. **16**, 2050, **2016**.
 58. TANG Y., WEN G., LI P., DAI C., HAN J. Effects of biogas slurry application on crop production and soil properties in a rice–wheat rotation on coastal reclaimed farmland. *Water, Air, & Soil Pollution*. **230**, 51, **2019**.
 59. WANG Z., SANUSI I.A., WANG J., YE X., KANA E.B.G., OLANIRAN A.O., SHAO H. Developments and prospects of farmland application of biogas slurry in China – a review. *Microorganisms*. **11** (11), 2675, **2023**.
 60. WANG L., LIU S., LI J., LI S. Effects of several organic fertilizers on heavy metal passivation in Cd-contaminated gray-purple soil. *Frontiers in Environmental Science*. **10**, 895646, **2022**.
 61. CHEN Y., LIU Y., WANG Z. Influence of organic manure on soil chemical properties. *Journal of Agricultural Chemistry and Environment*. **47** (2) 89, **2020**.
 62. MAGDOFF F.R., VAN ES H.M. Building soils for better crops: Sustainable soil management. 3rd ed. Sustainable Agriculture Research and Education Program (SARE), **2009**.
 63. LEHMANN J., RILLIG M.C., THIES J., MASIELLO C.A., HOCKADAY W.C., CROWLEY D. Biochar effects on soil biota—a review. *Soil Biology and Biochemistry*. **43** (9), 1812, **2008**.
 64. KHAITOV B., YUN H.J., LEE Y., RUZIEV F., LE T.H., UMURZOKOV M., BO BO A., CHO K.M., PARK K.W. Impact of organic manure on growth, nutrient content and yield of Chilli Pepper under various temperature environments. *International Journal of Environmental Research and Public Health*. **16**, 3031, **2019**.
 65. VERMA S., PRADHAN S.S., SINGH A., KUSHUWAHA M. Effect of organic manure on different soil properties: A Review. *International Journal of Plant and Soil Science*. **36** (5), 182, **2024**.
 66. FENG G., HAO F., HE W., RAN Q., NIE G., HUANG L., WANG X., YUAN S., XU W., ZHANG X. Effect of biogas slurry on the soil properties and microbial composition in an annual ryegrass-silage maize rotation system over a five-year period. *Microorganisms*. **12** (4), 716, **2024**.
 67. RAMADAN K.M.A., EL-BELTAGI H.S., EL-MAGEED T.A.A., SAUDY H.S., AL-OTAIBI H.H., MAHMOUD M.A.A. The Changes in various physio-biochemical parameters and yield traits of faba bean due to humic acid plus 6-benzylaminopurine application under deficit irrigation. *Agronomy*. **13**, 1227, **2023**.
 68. HUANG M., ZHU Y., LI Z., HUANG B., LUO N., LIU C., ZENG G. Compost as a soil amendment to remediate heavy metal-contaminated agricultural soil: Mechanisms, efficacy, problems, strategies. *Water, Air and Soil Pollution*. **227**, 359, **2016**.
 69. MAYANS B., PÉREZ-ESTEBAN J., ESCOLÁSTICO C., EYMAR E., MASAGUER A. Evaluation of commercial humic substances and other organic amendments for the immobilization of copper through ¹³C CPMAS NMR, FT-IR, DSC analyses. *Agronomy*. **9**, 762, **2019**.
 70. MOHAMED AA, EL-BELTAGI HS, RASHED MM. Cadmium stress induced change in some hydrolytic enzymes, free radical formation and ultrastructural disorders in radish plant. *Electronic Journal of Environmental, Agricultural and Food Chemistry*. **8**, 969, **2009**.
 71. EL-BELTAGI, H.S., MOHAMED, A.A. Changes in non protein thiols, some antioxidant enzymes activity and ultrastructural alteration in radish plant (*Raphanus sativus* L.) Grown under lead toxicity. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*. **38**, 76, **2010**.
 72. AFIFY A., EL-BELTAGI H.S. Effect of insecticide cyanophos on liver function in adult male rats. *Fresenius Environmental Bulletin*. **20** (4a), 1084, **2011**.
 73. EL-BELTAGI H.S., HALEMA A.A., ALMUTAIRI Z.M., ALMUTAIRI H.H., ELARABI N.I.,

- ABDELHADI A.A., HENAWY A.R., ABDELHALEEM H.A.R. Draft genome analysis for *Enterobacter kobei*, a promising lead bioremediation bacterium. *Frontiers in Bioengineering and Biotechnology*. **11**,1335854, **2024**.
74. HALEMA A.A., EL-BELTAGI H.S., AL-DOSSARY O., ALSUBAIE B., HENAWY A.R., REZK A.A., ELARABI N.I., ABDELHADI A.A. Omics technology draws a comprehensive heavy metal resistance strategy in bacteria. *World Journal of Microbiology and Biotechnology*. **40**, 193, **2024**.
75. GUO D., REN C., ALI A., LI R., DU J., LIU X., GUAN W., ZHANG Z. *Streptomyces pactum* combined with manure compost alters soil fertility and enzymatic activities, enhancing phytoextraction of potentially toxic metals (PTMs) in a smelter-contaminated soil. *Ecotoxicology and Environmental Safety*. **181**, 312, **2019**.
76. WANG Q., CHEN Z., ZHAO J., MA J, YU Q., ZOU P., LIN H., MA J. Fate of heavy metals and bacterial community composition following biogas slurry application in a single rice cropping system. *Journal of Soils and Sediments*. **22**, 968, **2022**.
77. YANG Q., LIU P., DONG S., ZHANG J., ZHAO B. Combined application of organic and inorganic fertilizers mitigates ammonia and nitrous oxide emissions in a maize field. *Nutrient Cycling in Agroecosystems*. **117**, 13, **2020**.
78. LI Y., WU X.P., HE G., WANG Z.H. Benefits of yield, environment and economy from substituting fertilizer by manure for wheat production of China. *Scientia Agricultura Sinica*. **53**, 4879, **2020**.