

Original Research

Meteorological Parameters versus PM₁₀: Statistical Analysis of the City of Skopje, Republic of North Macedonia Using Multiple Linear Regression

Emilija Manevska^{o*}, Olgica Dimitrovska, Ivan Radevski, Svemir Gorin, Arse Kuzmanoski, Blagoja Markoski

Institute for Geography, Faculty of Natural Science and Mathematics, Ss. Cyril and Methodius University;
Arhimedova 3, 1000, Skopje

Received: 15 October 2024

Accepted: 19 April 2025

Abstract

This scientific paper aims to analyze the multilinear regression between the meteorological parameters air temperature (T_{air}), relative humidity (RH), wind speed (WS), and precipitation (P) to statistically determine the impact on the concentration of PM₁₀ pollutant. Since the city of Skopje has the most polluted air in the region, particularly in winter, this scientific research focuses on exploring the correlations between the meteorological parameters and the air pollution from PM₁₀. The results proved the moderate negative correlation ($r = -0.76$, $p = 0.0001$) between air temperature and PM₁₀ versus the moderate positive correlation ($r = 0.42$, $p = 0.24$) between relative humidity and PM₁₀ and the weak negative correlation between wind speed and PM₁₀ ($r = -0.32$, $p = 0.06$) and ($r = -0.19$, $p = 0.001$) for precipitation. The model was tested with a goodness-of-fit test and ANOVA, and it was proved that the model was adequate. It can be concluded that there can be different results in the correlation because of different climate, geographical position, topography, and socioeconomic factors in the region.

Keywords: Skopje, air pollution, meteorological parameters, correlation, multilinear regression

Introduction

Air pollution is a great environmental challenge in today's society. Air pollutants are a complex combination of gases and particulate matter [1]. Particulate air pollution is a mixture of solid, liquid, or solid and liquid particles suspended in the air [2]. PM is emitted directly into the atmosphere (primary) or formed in the atmosphere through gas-to-particle

conversion (secondary) [1]. PM particles often contain heavy metals such as Mn, Co, Ni, Cu, Cd, and Pb, with their concentrations significantly higher in industrial areas within urban settings [3]. Air pollution in cities is a serious environmental problem where the air quality is the result of a complex interaction between natural and anthropogenic environmental conditions, especially in developing countries [4, 5]. Today, Skopje is one of the most polluted cities in Europe [5], where numerous studies show that PM₁₀ concentrations exceeded the annual limit value, peaking during the winter [5, 6]. Often, these pollutants cause severe conditions in human health, particularly respiratory diseases [7, 8].

*e-mail: emilijamanevska@ukim.pmf.mk

^oORCID iD: 0009-0000-0605-8340

The consequences of air pollution with particulate matter suggest that in selected North Macedonian cities 1,903 human lives (excess deaths) are lost annually due to PM exposures [9], while the estimated economic cost associated with mortality from exposure to air pollution in North Macedonia is in the range of US\$ 500-900 million annually, equivalent to 5.2-8.5% of gross domestic product (GDP) in 2016 [10]. The air in Skopje is one of the most polluted in the world because of the city's natural position and people's reliance on fossil fuels [11]. The geographical position of the city plays a key role in the air pollution. Topography and meteorological conditions contribute to poor air quality in Skopje [12]. Skopje is geographically located in a basin surrounded by mountains, which entraps the pollution; there is insufficient aerial circulation that can otherwise help in diluting the toxic air [13, 14]. Contemporary studies have noted that the climate change impact is more influential in the mountainous areas [15], which is a main geographical characteristic in North Macedonia as a country with around 80% hilly mountain surface. The main anthropogenic sources of ambient PM air pollution in North Macedonia in 2014 were identified as: residential heating (36% of total primary emissions), industrial processes (33%), and energy production (20%). Traffic contributed 2% of the total emissions of PM particles [9]. The low traffic effect was proven elsewhere in the neighboring countries [16]. The conducted studies for the air pollution in Skopje provide indicative data that domestic households' biomass combustion has the largest impact on the disturbed air quality in Skopje, especially for the emission of suspended particles PM_{10} . This also corresponds with the fact that households in Skopje are using stoves with inefficient fuel combustion and low-quality fuels of suspicious origin [17].

Many scientific studies confirm the impact of meteorological factors on air pollution [18-20].

Although previously mentioned anthropogenic factors contribute to the natural conditions in the city of Skopje, the meteorological parameters exert a substantial influence on the prevalence of air pollution in Skopje. This is confirmed by the fact that the highest concentrations of air pollution and PM_{10} pollutants are measured during the winter period. The high concentrations of air pollutants are observed during stable atmospheric conditions, with emitted substances accumulating in the basin. During these periods, there is reduced circulation in the atmosphere due to prolonged periods with weak winds, foggy days, and very little rain, as well as the occurrence of temperature inversion [21].

In general, the studies showed a negative correlation of PM_{10} with the air temperature and the atmospheric pressure, and a positive correlation with the relative humidity [22-26].

Wider European study analyzed the effect of geographical parameters on PM_{10} in European landscapes. The results showed that in the cooling season, air temperature, precipitation, and wind speed

are the most important factors for PM_{10} concentrations in suburban regions. Although all of the previously mentioned climate parameters are important, the results showed that precipitation has the highest impact on the pollutant. In the urban areas, the effect of these three climate variables is significantly lower [27].

Despite these findings, the specific relationship between meteorological factors and PM_{10} concentrations in Skopje remains underexplored. Given that climatic factors influence each other as well as particulate matter concentrations – for instance, the speed and direction of the wind directly influence environmental factors such as air temperature, humidity, and dust transport [28], all of which play a crucial role in air pollution dynamics – this study aims to fill this gap by analyzing how meteorological parameters impact PM_{10} levels in Skopje's unique geographical and climatic context.

This study enhances the understanding of PM_{10} variability by applying Multiple Linear Regression (MLR) to quantify the impact of meteorological factors while controlling for multiple variables simultaneously. Unlike generalized assessments, it ensures statistical validity through data transformation techniques and addresses autocorrelation issues, improving the reliability of the model. By tailoring MLR to Skopje's unique geographical and climatic conditions, this research provides a more precise and data-driven approach to analyzing air pollution dynamics.

Understanding this relationship is vital for developing targeted strategies to manage air pollution in the city. The results will provide critical insights for authorities to mitigate air pollution through a better understanding of these correlations.

Study Area

The City of Skopje is situated in the Skopje Basin, located in the northern region of the Republic of North Macedonia, between 41°42'15"N and 42°16'20"N, 21°09'40"E and 21°49'15"E., along Greenwich (Fig. 1 and 2). The geographical boundaries of this area are defined by the mountain massif Karadzica and Goleshnica from the south, Mountain Gradeshka from east to the Ovche Pole (plain), the Kumanovo Valley and the Skopska Crna Gora massif from north, and the Zheden Mountain from west separates it from the Polog Valley. Within this geographical frame, the Skopje basin covers an area of 1924.2 km², of which 343.9 km² are flatland located in a hypsometric belt of 150-300 m.a.s.l. [29].

From an administrative perspective, Skopje is a key component of the Skopje region, one of the eight statistical regions in the Republic of North Macedonia. Within its boundaries, Skopje encompasses ten municipalities, while the entire region covers an expanse of 1812 km²; the city extends over an area of 571.97 km² [30].

As of the most recent census conducted in 2021, Skopje is home to a total population of 526,502 residents,




Fig. 1. Geographical position of the city of Skopje within the Republic of North Macedonia. DEM Source: USGS 2015, Data Source: OSM.




Fig. 2. Geographical position of the city of Skopje within the Skopje Basin. DEM Source: USGS 2015, Data Source: OSM.

which is 27.7% of the entire population of the Republic of North Macedonia [30]. However, it is estimated that on a daily basis, the number is higher due to the fact that a large number of residents from other regions often migrate and stay in the city due to work obligations, study, etc. Both the increasing population and migration from rural to urban areas lead to the inevitable opening of new areas for settlement in urban centers [31]. Constant internal migrations to the capital are also the main reason for the need to build new housing facilities.

With the increase of the urbanization in the city, the air purification is additionally interrupted, and with the construction of high buildings in a narrow space in the central area of the city, the natural and necessary circulation of the winds, which would be used to purify the air in a natural way, becomes more difficult [32].

The majority of industrial, commercial, and service facilities are concentrated within the city. Functioning as the capital of the Republic of North Macedonia, Skopje serves as the nation's economic, administrative, and cultural hub.

Due to the constant expansion of the city, rapid urbanization, and the adoption of a fast-paced, modern lifestyle, Skopje has become a subject of research regarding its longstanding issues with air pollution, mainly focusing on analyzing the causes, consequences, and potential solutions to address the present environmental challenges in the capital.

The geographical position and orographic peculiarities are the main modifiers of the climate in the city of Skopje.

The Skopje basin is influenced by the continental climate from the north, the Mediterranean climate from the south, and their modifications. The average annual air temperature is 12.7°C [33]. Air temperature inversions in the Skopje basin occur in all months, but they are strongly expressed in the winter months, most often during anticyclone weather situations.

The Skopje Basin is a specific area with frequent occurrences of fog and temperature inversions. In such weather conditions, air pollution increases. On the days with temperature inversions, it is coldest in the low parts of the basin, while the temperature increases with altitude towards the higher regions of Vodno and Skopska Crna Gora mountains [33].

Precipitation is unevenly distributed throughout the year (monthly and seasonally). The lowest values were registered in the months of February and July. The average annual amount of precipitation (1951-2010) ranges from 497.5 mm in the plains to about 700 mm in the highest parts of Vodno and the low slopes of Skopska Crna Gora Mountain [33].

In the Skopje Basin in the average multi-year wind time series, the most frequent wind directions are from the north and south, and a factor that contributes to the higher air pollution is the fact that the urban part of the Skopje basin has a much higher frequency of silences - conditions without wind [33].

The relative humidity in the Skopje basin decreases from January to August and increases from August to December. The highest monthly values of relative humidity are in November, December, and January at 82-84%, and the lowest average monthly relative humidity values are in July and August, at 56-57% [33].

The Skopje basin is characterized by an increased frequency of days with fog. Fog occurs throughout the day, but with the highest frequency in the morning.

The highest frequency of fog occurs in the winter period, in December, with an average of 15 days with fog, and in November, December, and January, with 61% of the total annual average number of days with fog occurring. On average, there are 63 days with fog in the Skopje basin [33].

Materials and Methods

In this scientific paper, the focus is placed on analyzing the data predominantly sourced from state institutions of the Republic of North Macedonia. The data analysis for the meteorological parameters for the period of nine years (2012-2020) was taken from the main meteorological gauge "Zajchev Rid" – Hydro meteorological Service of the Republic of North Macedonia and the data for the concentrations of the air pollutant PM_{10} was gathered from the gauge for air quality "Centar" – Ministry of Environment and Spatial Planning of North Macedonia for the same time period as the meteorological parameters. The problem of the relatively short analyzed period of the time series is a consequence of the short-term organized air pollution measurement in North Macedonia, where continuous measurement started in 2012.

The whole statistical analysis (descriptive statistics, testing normality and multiple linear regression, ANOVA testing) was run on the software "XLSTAT".

Since all of the cities in the Republic of North Macedonia, including the city of Skopje, have serious issues with air pollution, especially with the concentration of PM_{10} in the winter period, the standards for permitted concentrations are shown in Table 1.

Descriptive statistics were used for summarizing, organizing, and presenting data meaningfully and concisely. It can be presented as measures of central tendency (mean, median, mode, and range). In order to complete the multiple linear regression on the meteorological variables and PM_{10} , a normality analysis was conducted on the raw data using the Shapiro-Wilk test.

The Shapiro-Wilk test has two hypotheses:

- Null Hypothesis (H_0): The data follows a normal distribution.
- Alternative Hypothesis (H_a): The data does not follow a normal distribution

The Shapiro-Wilk test is particularly suitable for small to medium-sized datasets [37].

Table 1. Standards for permitted concentrations of PM_{10} .

Community/ Variable	Air Pollutant	Period	Limit Value ($\mu\text{g}/\text{m}^3$)
MKD	PM_{10}	24 h	50
		Annual	40
EU	PM_{10}	24 h	50
		Annual	40
WHO	PM_{10}	24 h	45
		Annual	15

Source: [34, 35, 36].

The main statistic W is estimated as follows:

$$W = \frac{(\sum_{i=1}^n a_i x_{(i)})^2}{\sum_{i=1}^n (x_i - \bar{x})^2} \quad (1)$$

Where x_i are ordered sample values, a_i are constants derived from the covariance matrix, and \bar{x} is the sample mean.

For the Shapiro-Wilk test, the significance level is 0.05, which means that if the p value is higher than the significance level, the data has a normal distribution [37].

Since some of the parameters didn't have a normal distribution, we used the Box-Cox transformation on the chosen data, and the results can be seen in Table 2. The Box-Cox transformation is applied to stabilize the variance and make the dataset closely approximate a normal distribution. It was necessary to transform this variable before attempting a multilinear regression [38].

$$y_i^{(\lambda)} = \begin{cases} y_i^\lambda - 1, & \text{if } \lambda \neq 0, \\ \ln(y), & \text{if } \lambda = 0, \end{cases} \quad (2)$$

λ determines the type of transformation (e.g., logarithmic when $\lambda = 0$). It is commonly used in regression and time series analysis to meet normality and homoscedasticity assumptions.

After the data had a normal distribution (tested again with the normality test Shapiro-Wilk), the transformed data of four meteorological variables and PM_{10} particles, similar time series data, were used in the multiple linear regression analysis.

Table 2. Shapiro-Wilk normality test results.

Distribution of Parameters	p -value
T_{air}	0.051
RH	0.143
WS	0.1
P	0.6
PM_{10}	0.06

Source: authors.

Multiple regression analysis (MR) is a highly flexible system for examining the relationship of a collection of independent variables (or predictors) to a single dependent variable (or criterion) [39, 40]. The equation for multiple linear regression has the same form as that for simple linear regression, but has more terms:

$$y_i = \beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i} + \dots + \beta_p x_{pi} + e_i \quad (2)$$

As for the simple case, β_0 is the constant, which will be the predicted value of y when all explanatory variables are 0. In a model with p explanatory variables, each explanatory variable has its own β -coefficient. Again, the analysis does not allow us to make causal inferences, but it does allow us to investigate how a set of explanatory variables is associated with a response variable of interest [41].

The degree of association is measured by a correlation coefficient, denoted by r . It is sometimes called Pearson's correlation coefficient after its originator and is a measure of linear association (Table 3). If a curved line is needed to express the relationship, other and more complicated measures of the correlation must be used [42].

The goodness of fit for the multiple linear regression (MLR) model was evaluated using several statistical metrics. The R^2 value was analyzed to determine the proportion of variance in the dependent variable explained by the independent variables. It contains various variables, the most important of which are the F-statistic and p -value [37].

The F-statistic tests the significance of the MLR model as follows:

$$F = \frac{\text{Explained variance per predictor}}{\text{Unexplained variance per residual degree of freedom}} \quad (3)$$

High F-value with a low p -value (<0.05) suggests that the model is significant and at least one independent variable is useful for predicting the dependent variable. Additionally, the Adjusted R^2 , AIC, and BIC values were examined to account for the number of predictors and ensure model accuracy and parsimony [41].

Following the MLR analysis, we conducted an Analysis of Variance (ANOVA) to test the significance of the overall model and to identify which meteorological parameters significantly influence PM_{10} levels. The ANOVA decomposes the total variability in PM_{10} concentrations into components attributed to the model and residual error, providing F-statistics to assess the explanatory power of the predictors [43]:

$$F = \frac{MS_{\text{model}}}{MS_{\text{error}}} \quad (4)$$

The MS model is usually used when testing the MLR model [44], which is the mean square of the model, and MS error is the mean square of the error. This two-step

Table 3. Detailed classification of Pearson correlation coefficient (r) values.

r value	Interpretation
$r = 1$	Perfect positive linear correlation
$1 > r \geq 0.8$	Strong positive linear correlation
$0.8 > r > 0.4$	Moderate positive linear correlation
$0.4 > r > 0$	Weak positive linear correlation
$r = 0$	No correlation
$0 > r \geq -0.4$	Weak negative linear correlation
$-0.4 > r \geq -0.8$	Moderate negative linear correlation
$-0.8 > r > -1$	Strong negative linear correlation
$R = -1$	Perfect negative linear correlation

Source: [45].

approach allowed us to evaluate the relationship between meteorological conditions and PM_{10} pollution rigorously.

Results

The descriptive statistics of the data collected from January 2012 to December 2021 of the data collected from the gauges Zajchev Rid and Centar are given in Table 4.

The mean air temperature (T_{air}) measured at the gauge Zajchev Rid was 13.8°C while the standard deviation was 8.3°C . There is a huge range between the minimum, which is -4.0°C , and the maximum, which is 27.7°C . The mean RH was 65.3% while the standard deviation was 10.9%. The minimal range for the measured period was 39.0%, and the maximum was 88.0%. The mean WS has a mean value of 2.5 m/s, and the standard deviation is 0.4 m/s. The minimum was 1.8 m/s while the maximum was 3.5 m/s. The mean precipitation measured for the given period was 44.8 mm/m^2 , and the standard deviation was 34.0 mm/m^2 . For the PM_{10} , the mean was $61.0 \mu\text{g/m}^3$ while the standard deviation was $40.0 \mu\text{g/m}^3$. The minimal range was $23.9 \mu\text{g/m}^3$ and the maximum

was $238.3 \mu\text{g/m}^3$, which is a difference of $214.4 \mu\text{g/m}^3$ (Table 4 and Fig. 3).

Since the descriptive analysis was made on raw data, the next step was the normality test, which was applied to the data. All of the parameters, except the data for wind speed, didn't follow a normal distribution, so the Box-Cox transformation was applied to the chosen data.

After the data had a normal distribution, the Shapiro-Wilk normality test was applied, which is shown in Table 2.

After the normality test, the multiple linear regression was conducted in order to test the significance of the independent variables, i.e., the meteorological parameters, in a linear regression where PM_{10} is chosen as the dependent variable.

From Table 5. It can be seen that air temperature has the strongest negative correlation in comparison with the other observed meteorological parameters, with PM_{10} , which is $r = -0.76$, ($p = 0.0001$), where, according to the classification given in Table 3, is classified as a moderate negative linear correlation. The negative correlation between air temperature and PM_{10} concentration indicates that higher average temperatures are associated with lower PM_{10} levels, while lower temperatures correspond to higher PM_{10} concentrations. The second strongest correlation of all climate parameters is the positive correlation between relative humidity and PM_{10} , which according to the classification signifies a moderate positive correlation with correlation coefficient $r = 0.42$ ($p = 0.24$). Contrary to the negative correlation between air temperature and PM_{10} , the positive correlation between relative humidity and PM_{10} means that when the relative humidity percentage is higher, the concentrations of PM_{10} in the air are also higher, and when the relative humidity percentage is lower, the concentrations of PM_{10} in the air are also lower.

The other climate parameters – wind speed (WS) and precipitation (P) have weak negative correlation with PM_{10} where the correlation coefficient between wind speed and PM_{10} is $r = -0.32$ ($p = 0.06$), while the correlation coefficient between precipitation and PM_{10} is the lowest of all previously analyzed climate parameters, which is $r = -0.19$ ($p = 0.01$).

From the previously analyzed results, it can be

Table 4. Descriptive statistic of the collected data.

Gauge Zajchev Rid / Centar	Unit	M \pm SD	Range [Min; Max]
T_{air}	$^{\circ}\text{C}$	13.78 ± 8.27	[-4.00; 27.70]
RH	%	65.28 ± 10.92	[39.00; 88.00]
WS	m/s	2.48 ± 0.36	[1.80; 3.50]
P	mm/m^2	44.81 ± 34.02	[0.00; 167.50]
PM_{10}	$\mu\text{g/m}^3$	61.01 ± 40.04	[23.90; 238.30]

Source: authors.




Fig. 3. Box plots 1. The median (denoted by a horizontal bar in the box), the 25th percentile (denoted by the bottom edge of the box), the 75th percentile (denoted by the top edge of the box), the 5th percentiles (denoted by the bottom edge of the whisker), the 95th percentiles (denoted by the top edge of the whisker). Source: authors.

Table 5. Pearson correlation (r) between meteorological parameters and PM_{10} concentrations.

	RH	P	WS	T_{air}	PM_{10}
RH	1	0.010	0.097	-0.659	0.415
P	0.010	1	-0.071	-0.007	-0.192
WS	0.097	-0.071	1	0.246	-0.316
T_{air}	-0.659	-0.007	0.246	1	-0.763
PM_{10}	0.415	-0.192	-0.316	-0.763	1

Source: authors.

noticed that in the investigated period for the given gauges, the meteorological parameters: air temperature, wind speed, and precipitation have a negative correlation with PM_{10} , while relative humidity has a positive correlation with PM_{10} . The obtained MLR equation is as follows:

$$PM_{10} = 0.16 - 1.13 \cdot RH + 4.05 \cdot P - 3.27 \cdot WS - 2.87 \cdot T_{air} \quad (5)$$

This Equation represents the relationship between PM_{10} concentration and the meteorological factors (Relative Humidity, Precipitation, Wind Speed, and Air Temperature) based on the coefficients obtained from the regression analysis.

Where:

- The constant term (0.164) represents the intercept of the regression equation, indicating the value of PM_{10} when all independent variables are zero;
- The beta values, or regression coefficients, indicate the expected change in the dependent variable for a one-unit change in the corresponding independent variable, while holding other variables constant.

The goodness of fit procedure was performed (see Table 6).

The sum of weights indicates that each observation is equally weighted (weight of 1 per observation). Degrees of freedom for the error term, which is the total

number of observations minus the number of parameters estimated (including the intercept).

R^2 (Coefficient of Determination): 0.648 indicates that 64.8% of the variability in PM_{10} levels is explained by the model.

Adjusted for the number of predictors in the model, providing a more accurate measure than R^2 when multiple predictors are involved. This value indicates that 63.5% of the variability is explained by the model after adjusting for the number of predictors. The average of the squared differences between observed and predicted values. This value, being very close to zero, indicates a good fit. The square root of the MSE provides a measure of the average magnitude of the error in the same units as the response variable (PM_{10}). A small RMSE indicates a good fit of the multiple linear regression, the average absolute percentage error between observed and predicted values. A MAPE of 3.476% indicates that, on average, the predictions are within 3.476% of the actual values. The Durbin-Watson statistic is a measure of autocorrelation in the residuals. Values close to 2 indicate no autocorrelation, while values closer to 0 or 4 suggest positive or negative autocorrelation, respectively. A DW of 0.587 indicates potential positive autocorrelation. Cp is a measure used to assess the fit of regression models, ideally close to the number of predictors plus the intercept. Here, Cp being 5 suggests a model with potentially 4 predictors is well-specified. AIC (Akaike Information Criterion)

Table 6. Goodness-of-fit of the data.

N	Observations	108.000
1	Sum of weights	108.000
2	DF (degrees of freedom)	103.000
3	R ² (ratio between the sum of explained variation by the model and the actual variation)	0.648
4	Adjusted R ²	0.635
5	MSE (mean squared error)	0.000
6	RMSE (root mean squared error)	0.006
7	MAPE (Mean Absolute Percentage Error)	3.476
8	DW (Durbin Watson statistics)	0.587
9	Cp (Mallows' Cp depends on the MSE)	5.000
10	AIC (Akaike Information Criterion)	-1109.333
11	SBC (Schwarz Bayesian Criterion)	-1095.923
12	PC (Amemiya Prediction Criterion)	0.386

is a measure of model quality, with lower values indicating a better fit. Negative values suggest a very good model fit.

SBC (Schwarz Bayesian Criterion/BIC) is similar to AIC but includes a penalty for the number of parameters. Lower values indicate a better model. PC (Prediction Criterion) is another measure of model fit, with lower values indicating better predictive accuracy.

The p-value (<0.0001) in ANOVA indicates the probability that the observed F-value would occur if the null hypothesis were true. A very small p-value (less than 0.05) suggests strong evidence against the null hypothesis, indicating that the model is statistically significant. The ANOVA results (Table 7) indicate that the model is highly significant ($p<0.0001$), meaning that the variations explained by the model are much greater than would be expected by chance. The model explains a significant portion of the variability in PM₁₀ levels. A high F-value (47.449) suggests that the model provides a good fit to the data. The degrees of freedom and the sum of squares give insight into how much variation is being explained by the model versus the error.

The results section offers a detailed analysis of meteorological factors and their relationship with PM₁₀ concentrations. Air temperature showed a moderate negative correlation with PM₁₀ ($r = -0.76$), indicating

reduced concentrations during warmer conditions. In contrast, relative humidity displayed a moderate positive correlation ($r = 0.42$), suggesting that higher moisture levels contribute to increased PM₁₀. Wind speed and precipitation had weaker negative correlations, indicating limited effects on PM₁₀ dispersion or removal. The multiple linear regression model explained 64.8% of the variability in PM₁₀ levels, with goodness-of-fit measures supporting its reliability.

The findings indicate a statistically significant relationship between meteorological parameters and PM₁₀ levels, yet the model's explanatory power ($R^2 = 64.8\%$) suggests that nearly 35% of the variation remains unexplained, potentially influenced by unaccounted emission sources or secondary aerosol formation. The strong negative correlation between air temperature and PM₁₀ may reflect seasonal heating and emission patterns rather than a purely meteorological effect, warranting further investigation into emission inventories. Additionally, the positive autocorrelation ($DW = 0.587$) suggests potential temporal dependencies in PM₁₀ levels, indicating the need for time-series modeling or non-linear approaches for improved predictive accuracy.

Discussion

Several scientific studies proved that the meteorological conditions can affect air pollution. More precisely, some meteorological parameters affect the atmospheric pollution [46-51]. Multiple linear regression has been widely applied in various cities to investigate the relationship between meteorological factors and air pollution, as urban areas often face challenges with poor air quality. Some other scientific research that analyzed the correlation between the meteorological parameters (T_{air} , WS, RH, and P) on PM₁₀ proved similar results to this research (Table 8).

Table 8 contains the correlation data between PM₁₀ levels and key meteorological parameters - air temperature (T_{air}), precipitation (P), relative humidity (RH), and wind speed (WS) - across various locations, including cities and regions. For instance, in Tirana and Skopje, a strong negative correlation with temperature indicates that higher temperatures correspond to lower PM₁₀ levels, likely due to enhanced atmospheric dispersion. In contrast, regions like Northern Thailand and cities such as Athens show positive correlations between air temperature and PM₁₀,

Table 7. Analysis of variance, ANOVA (PM₁₀).

Source	DF	Sum of squares	Mean squares	F	Pr > F
Model	4	0.006	0.002	47.449	< 0.0001
Error	103	0.003	0.000		
Corrected Total	107	0.010			

Table 8. Correlation coefficient (r) between PM_{10} and meteorological parameters in the researched cities.

Gauge/ Parameters	T_{air}	P	RH	WS	Citation
Tirana	-0.88	/	0.21	/	[49]
Sosnowiec	0.08	0.03	/	0.25	[50]
Izmir	0.11	/	0.70	- 0.41	[51]
Budapest	-0.38	/	0.25	-0.44	[52]
Pécs	-0.37	/	0.24	-0.33	[52]
Miskolc	-0.48	/	0.30	-0.26	[52]
Athens	0.44	0.17	0.01	-0.59	[53]
Istanbul	-0.15	/	0.01	-0.12	[54]
Transylvania	-0.68	0.34	0.38	-0.47	[55]
Northern Thailand	0.53	/	0.60	0.04	[56]
Moldova region	-0.34	-0.06	0.15	-0.14	[57]
China	-0.36	-0.07	-0.18	-0.04	[58]
Pune	-0.25	-0.17	-0.23	-0.37	[59]
Skopje	-0.76	-0.19	0.42	-0.32	Own research

Source: authors.

suggesting that warmer conditions may contribute to increased particulate matter concentrations. Wind speed generally has a negative correlation with PM_{10} in most areas, including Athens and Transylvania, indicating that stronger winds help disperse pollutants. These correlations highlight the varied influence of meteorological factors on PM_{10} levels, shaped by local geographical and climatic conditions.

In the Table 8, samples of research in different cities are presented, using the same methodology.

From the given table, it can be noticed that the correlation between air temperature (T_{air}) and PM_{10} varies from strong negative correlation ($r = -0.88$) in Tirana to moderate positive correlation in Northern Thailand ($r = 0.53$) and in Athens ($r = 0.44$). In the city of Skopje, the correlation between the two variables is ($r = -0.76$), which is a moderate negative correlation. The similarity between the results from Tirana and Skopje can be explained by the geographical closeness between the two cities, although the cities have different climates, they have similar population rates, and both cities have bad air quality due to some anthropogenic factors such as urbanization and residential heating. In all of the researched areas, the correlation between PM_{10} and air temperature was negative.

Skopje indicated a weak negative correlation between PM_{10} and precipitation ($r = -0.19$), suggesting that rainfall marginally helps reduce PM_{10} levels by washing particles out of the air. In comparison, regions like Pune ($r = -0.17$) and Moldova ($r = -0.06$) show similar weak negative correlations. In contrast, Transylvania ($r = 0.34$)

shows a moderate positive correlation, which is unusual since rainfall typically reduces particulate matter. This discrepancy may be related to local factors such as soil composition or post-rain re-suspension of particles. Other cities, such as Sosnowiec ($r = 0.03$) and Athens ($r = 0.17$), also show weak positive correlations.

Considering the fact that relative humidity has seasonal variability, which shapes the dynamics of the local ecosystems, which then affects the urban life quality [60], Skopje has a moderate positive correlation between PM_{10} and relative humidity ($r = 0.42$). As mentioned before, the Skopje basin has a lot of days with fog, which suggests that in the winter period, the fog could keep the PM_{10} pollutant longer in the air, which could explain the positive correlation between relative humidity and PM_{10} .

This trend is mirrored in almost all of the researched areas, for instance, regions like Izmir ($r = 0.70$), where it was concluded that the RH affects the natural deposition process of PM_{10} , and Northern Thailand ($r = 0.60$), where higher humidity is linked to higher PM_{10} levels. This could be due to moisture facilitating the aggregation of particles, increasing their size and concentration in the air. However, in some cities such as Athens ($r = 0.01$) and Istanbul ($r = 0.01$), the correlation is almost negligible. In some studies, relative humidity has been found to exhibit a negative correlation with PM_{10} levels [61-63]. However, in locations such as Pune ($r = -0.23$) and China ($r = -0.18$), the relationship is more varied, displaying a weak inverse correlation.

The Pearson correlation between wind speed and

PM₁₀ is generally negative, ranging from a moderate negative correlation ($r = -0.59$) in Athens to a weak positive correlation ($r = 0.04$) in Northern Thailand. In most of the studied areas, the correlation remains negative, as observed in Skopje, where the correlation coefficient is ($r = -0.32$).

We took into consideration the fact that the relationship between PM₁₀ concentrations and meteorological parameters differs between seasons, as well as between years, i.e., have seasonal and annual variation [64] and also the geographical position of the region has a key role in the impact of the meteorological parameters on PM₁₀ as previously mentioned.

Comparing the seasonal differences, research made in Shenyang, China, showed results where PM₁₀ had a negative correlation with wind speed and air temperature in autumn and winter, but no significant correlation in spring and summer [48].

The very low number of $MSE < 0.0009$ confirmed that the model of MLR in this research is very well fitted, which is complementary with a previous study about the impact of meteorological factors on PM₁₀ in Kocaeli, Turkey, where MSE varies from 0.00019 to 0.00086 [61].

The results of RMSE show a much lower value of 0.0006 and an excellently fitted MLR compared with the results from Pune, India, where the $RMSE = 20.5$ [59].

Although, in the city of Skopje, air pollution is a consequence of the geographical position, anthropogenic factors such as high population density, urbanization and the frequent usage of fossil fuels, in this paper was also proved the contribution of the meteorological parameters (air temperature, precipitation, wind speed and relative humidity) on the concentration of PM₁₀ with the correlation between all observed meteorological parameters.

In future studies, these results could be combined with other complementary methodologies like canonical correlation analysis, which determines air temperature as the best-correlated factor to PM₁₀ concentrations [65].

Conclusions

Air pollution is a significant problem in today's city of Skopje. A lot of factors cause the bad air quality, which also has great consequences for the residents of the city.

The Skopje basin is surrounded by mountains, which entrap the pollution that is affected by insufficient aerial circulation. The climate conditions, which are characterized by frequent occurrences of fogs and temperature inversions, worsen the air pollution.

To the mentioned geographical conditions, huge contributors to the bad air quality are the anthropogenic factors, which play a key role in the air pollution that occurs during the winter, such as residential heating, which is overall the main cause of the air pollution in the city.

The results of the multilinear regression between the

meteorological parameters and PM₁₀ showed that, mostly in the researched areas, the meteorological parameters were in negative correlation with PM₁₀, except relative humidity, which in the researched areas mostly had positive correlation with PM₁₀. The results showed that in the city of Skopje, as the air temperature lowers, the PM₁₀ values go higher, i.e., the air quality worsens as a consequence of frequent use of non-environmentally friendly ways of residential heating. Precipitation and wind speed also have a negative correlation with PM₁₀. Although the correlation between precipitation and wind speed is not significant it also showed that precipitation and wind speed can have an impact on the concentrations of PM₁₀, which can be explained with the fact that wind speed and direction significantly influence local microclimates, thermal comfort and air quality [28] where stronger winds help disperse airborne pollutants, thus lowering PM₁₀ levels, while rainfall cleans the air by washing away particulate matter. Humidity is positively correlated with PM₁₀ as a result of the frequent days with fog, which can be explained by the fact that the fog entraps the PM₁₀.

The methodology in this paper is complementary to many studies that explored the correlation between meteorology and PM₁₀ in many urban cities. It can be concluded that variations in correlation may arise due to differences in climate, geographical location, topography, and socio-economic factors in the region. The strong negative correlation between air temperature and PM₁₀ concentrations may be influenced not only by meteorological conditions but also by seasonal variations in emissions, such as increased heating-related pollution in colder months. This highlights the need for a more comprehensive analysis incorporating emission inventories to distinguish meteorological effects from anthropogenic influences. Additionally, the positive autocorrelation ($DW = 0.587$) suggests that PM₁₀ levels exhibit temporal dependencies, meaning past concentrations may influence future values. This indicates that traditional regression models may not fully capture the underlying patterns, and more advanced time-series or non-linear modeling approaches could improve predictive accuracy.

This study could be extrapolated to all Balkan capitals in order to investigate the main causes for air pollution, since the Balkan Peninsula, as a geographical region, has the most polluted air in Europe.

Conflict of Interest

The authors declare no conflict of interest.

References

1. ZHANG R., WANG G., GUO S., ZAMORA M.L., YING Q., LIN Y., WANG W., HU M., WANG Y. Formation of Urban Fine Particulate Matter. *Chemical Reviews*. **115**

- (10), 3803, **2015**.
2. BRUNEKREEF B., HOLGATE S.T. Air pollution and health. *The Lancet*. **360** (9341), 1233, **2002**.
3. YU Y., JIAYING W., JIASHU Y., HONGQUAN C., MIN L. Spatial and Temporal Distribution Characteristics of PM_{2.5} and PM₁₀ in the Urban Agglomeration of China's Yangtze River Delta, China. *Polish Journal of Environmental Studies*. **28** (1), 445, **2019**.
4. MAYER H. Air pollution in cities. *Atmospheric Environment*. **33** (24-25), 4029, **1999**.
5. UNEP. Air pollution in the Western Balkans. Available online: <https://zoynet.org/wp-content/uploads/2022/02/Pollution-Balkans-EN2.pdf> (accessed on 10.07.2024).
6. MANEVSKA E., DIMITROVSKA O., RADEVSKI I. Analysis of the concentrations of PM₁₀ and PM_{2.5} particles in the air in the city of Skopje and their impact on its quality. *Geographical Reviews*. **53**, 65, **2020**.
7. JIN X., JINGSEN F., HONGYA N., PEI L., QIANQIAN Y. Analysis of Sources and Concentrations of Heavy Metal Contents in PM₁₀ over a Four-Season Cycle in a Heavily Industrialised City in China. *Polish Journal of Environmental Studies*. **28** (5), 3227, **2019**.
8. SUN R., NA T., ZHIJUN C., XIAOWEI G., LIANHONG Z., JUNHUA Z. Effects of Meteorological Factors and Air Pollutants on Tuberculosis Incidence: A Distribution Lag Non-Linear Analysis. *Polish Journal of Environmental Studies*. **34** (3), 3269, **2025**.
9. AVIS R. Causes and Consequences of Air Pollution in North Macedonia. Available online: https://mvdsl.seeu.edu.mk/papers/K4D_NM2.pdf (accessed on 27.08.2024), **2022**.
10. WORLD BANK. Air Pollution Management in North Macedonia. Available online: <https://documents1.worldbank.org/curated/en/116521576516981237/pdf/Air-Quality-Management-in-North-Macedonia.pdf> (accessed on 01.06.2024).
11. EUROPEAN COMMISSION, CORDIS. Air pollution in Skopje: how citizens spurred policymakers towards the change. Available online: <https://cordis.europa.eu/article/id/413275-air-pollution-in-skopje-how-citizens-spurred-policymakers-towards-the-change> (accessed on 08.06.2024).
12. MARTINEZ G., SPADARO J., CHAPIZANIS D., KENDROVSKI V., KOCHUBOVSKI M., MUDU P. Health Impacts and Economic Costs of Air Pollution in the Metropolitan Area of Skopje. *International Journal of Environmental Research and Public Health*. **15** (4), 626, **2018**.
13. CEE INSTITUTE. Macedonia Social briefing: Air Pollution Crisis in Macedonia: Sources, Public Reaction and Policy Responses. Available online: <https://china-cee.eu/2018/02/19/macedonia-social-briefing-air-pollution-crisis-in-macedoniasourcespublic-reaction-and-policy-responses/> (accessed on 04.06.2024).
14. ECOHUBMAP. Air pollution in Skopje, North Macedonia, Available online: <https://www.ecohubmap.com/hot-spot/air-pollution-in-skopje-north-macedonia/9mwr0mml9f94w25> (accessed on 11.02.2024).
15. GUR E., PALTA S., OZEL B.H., VAROL T., SEVIK H., CETIN M., KOCAN N. Assessment of Climate Change Impact on Highland Areas in Kastamonu, Turkey. *Anthropocene*. **46** (1), 100432, **2024**.
16. BEŽANOVIĆ V., MLADENKA N., DUŠAN M., ZORAN Č., DRAGAN A. Effects of the COVID-19 Quarantine Measures on PM_{2.5}, PM₁₀ and NO₂ Concentration Levels in Serbia. *Polish Journal of Environmental Studies*. **31** (4), 3509, **2022**.
17. TRPEVSKA-TRAJKOVSKA M., STEFANOVA E. The influence of domestic fireplaces on air quality. *Techolab DOO Skopje, Proceedings - Macedonian Academy of Sciences and Arts*, **2019** [In Macedonian].
18. PÉREZ I.A., GARCIA A., SANCHEZ L., PARDO N., FERNANDEZ-DUKUE F. Key Points in Air Pollution Meteorology. *International Journal of Environmental Research and Public Health*. **17** (22), 8349, **2020**.
19. GUTTIKUNDA S.K., GURJAR B.R. Role of meteorology in seasonality of air pollution in megacity Delhi, India. *Environ Monit Assess*. **184**, 3199, **2011**.
20. LIU Y., ZHOU Y., LU J. Exploring the relationship between air pollution and meteorological conditions in China under environmental governance. *Scientific Reports*. **10**, 14518, **2020**.
21. FINNISH METEOROLOGICAL INSTITUTE, MINISTRY OF ENVIRONMENT AND PHYSICAL PLANNING. Air quality improvement plan for Skopje agglomeration. Available online: https://www.eeas.europa.eu/sites/default/files/activity_3.2_benchmark_3.2.1_skopje_air_quality_improvement_plan.pdf (accessed on 14.07.2024).
22. KIREŠOVÁ S., GUZAN M. Determining the correlation between particulate matter PM10 and meteorological factors. *Eng*. **3** (3), 343, **2022**.
23. JIANG B., SUN C., MU S., ZHAO Z., CHEN Y., LIN Y., QIU L., GAO T. Differences in Airborne Particulate Matter Concentration in Urban Green Spaces with Different Spatial Structures in Xi'an, China. *Forests*. **13** (1), 14, **2022**.
24. YANG H., PENG Q., ZHOU J., SONG G., GONG X. The Unidirectional Causality Influence of Factors on PM_{2.5} in Shenyang City of China. *Scientific Reports*. **10**, 8403, **2020**.
25. KIM M.J. Changes in the Relationship between Particulate Matter and Surface Temperature in Seoul from 2002–2017. *Atmosphere*. **10** (5), 238, **2019**.
26. LUO H., ZHOU W., JISKANI I.M., WANG Z. Analyzing Characteristics of Particulate Matter Pollution in Open-Pit Coal Mines: Implications for Green Mining. *Energies*. **14** (9), 2680, **2021**.
27. SOHRAB S., CSIKÓS N., SZILASSI P. Effect of geographical parameters on PM10 pollution in European landscapes: a machine learning algorithm-based analysis. *Environmental Sciences Europe*. **36** (1), 152, **2024**.
28. CETIN Z.I. Wind Speeds of Bartın. *Impacts of Climate Change and Settlements. Menba Journal of Fisheries Faculty*. **10** (3), 223, **2024**.
29. LAZARESKA M., MARKOSKI B. Chronology of the urban expansion of Skopje, *Acta Geobalcánica*. **5-2**, 57, **2019**.
30. STATE STATISTICS OFFICE OF THE REPUBLIC OF MACEDONIA. Census of the population, households and apartments in the Republic of North Macedonia. Available online: <https://www.stat.gov.mk/pdf/2022/2.1.22.10-mk-en.pdf> (accessed on 17.11.2023) [In Macedonian].
31. KILICOGLU C., CETIN M., ARICAK B., SEVIK H. Site selection by using the multi-criteria technique – a case study of Bafra, Turkey. *Environmental Monitoring and Assessment*. **192** (9), 608, **2020**.
32. LEAP OF THE CITY OF SKOPJE (2020-2026), Skopje. Available online: <https://skopje.gov.mk/media/6058/leap-2020-2026.pdf> (accessed on 18.11.2023) [In Macedonian].
33. LAZAREVSKI A. Climate in Macedonia. 1st ed.;

- Publisher: Kultura Skopje, North Macedonia, pp. 253, **1993** [In Macedonian].
34. MINISTRY OF ENVIRONMENT AND PHYSICAL PLANNING. Available online: https://air.moepp.gov.mk/?page_id=118 (accessed on 23.11.2023), **2023**.
 35. EUROPEAN COMISSION. Available online: https://environment.ec.europa.eu/topics/air/air-quality/eu-air-quality-standards_en (accessed on 25.04.2024).
 36. WHO. What are the WHO Air quality guidelines? Available online: <https://www.who.int/news-room/feature-stories/detail/what-are-the-who-air-quality-guidelines> (accessed on 27.06.2024).
 37. SHAPIRO S.S., WILK M.B. An Analysis of Variance Test for Normality (Complete Samples). *Biometrika*, **52** (3/4), 591, **1965**.
 38. BOX G.E.P., COX D.R. An Analysis of Transformations. *Journal of the Royal Statistical Society. Series B (Methodological)*. **26** (2), 211, **1964**.
 39. TRANMER M., MURPHY J., ELLIOT M., PAMPAKA M. Multiple Linear Regression, 2nd ed.; Publisher: Cathie Marsh Institute, Manchester, UK, pp. 59, **2020**.
 40. BOUNAKHLA Y., ABDELFAH B., FRANCESCA C., MOUNIA T., BASSMA E.G., EL K.E.H., FATIHA Z., MOUSSA B. Overview of PM₁₀, PM_{2.5} and BC and their dependent relationships with meteorological variables in an urban area in Northwestern Morocco. *Atmosphere*. **14** (1), 162, **2023**.
 41. CAMPBELL M.J., SWINSCOW T.D.V. Statistics at square one, 11th ed.; Publisher: John Wiley and Sons Ltd Publication, Chichester, UK, pp. 193, **2009**.
 42. FROST J. Regression Analysis: An Intuitive Guide for Using and Interpreting Linear Models, 1st ed.; Publisher: Jim Publishing, Costa Mesa, CA, USA, pp. 340, **2019**.
 43. CUEVAS A., FEBRERO M., FRAIMAN R. An anova test for functional data. *Computational Statistics and Data Analysis*. **47** (1), 111, **2004**.
 44. JIE L., HOU K., WANG X., YANG P., Temporal-Spatial Variations of Concentrations of PM₁₀ and PM_{2.5} in Ambient Air. *Polish Journal of Environmental Studies*. **25** (6), 2435, **2016**.
 45. MOORE D.S. The basic practice of statistics, 9th ed.; Publisher: W. H. Freeman, New York, USA, pp. 800, **2004**.
 46. LEE S., HO C.H., CHOI Y.S. High-PM₁₀ concentration episodes in Seoul, Korea: Background sources and related meteorological conditions. *Atmospheric Environment*. **45** (39), 7240, **2021**.
 47. KAYES I., SHAHRIAR S.A., HASAN K., AKHTER M., KABIR M.M., SALAM M.A. The relationships between meteorological parameters and air pollutants in an urban environment. *Global Journal of Environmental Science and Management*. **5** (3), 265, **2019**.
 48. LI X., MA Y., WANG Y., LIU N., HONG Y. Temporal and spatial analyses of particulate matter (PM₁₀ and PM_{2.5}) and its relationship with meteorological parameters over an urban city in northeast China. *Atmospheric Research*. **198**, 185, **2017**.
 49. HYSENAJ M., DURAJ S. Spatiotemporal variations of particulate matter in Tirana. *Journal of Ecological Engineering*. **22** (11), 209, **2021**.
 50. ADAMEK A., ZIERNICKA-WOJTASZEK A. Comparison of regression models of PM10 particulate concentration in relation to selected meteorological elements based on the example of Sosnowiec, Poland. *Időjárás/Quarterly Journal of the Hungarian Meteorological Service*. **123** (1), 127, **2019**.
 51. BIRIM N.G., TURHAN C., ATALAY A.S., AKKURT G.G. The Influence of Meteorological Parameters on PM₁₀: A Statistical Analysis of an Urban and Rural Environment in Izmir/Türkiye. *Atmosphere*. **14** (3), 421, **2023**.
 52. FERENCZI Z., IMRE K., LAKATOS M., MOLNÁR Á., BOZÓ L., HOMOLYA E., GELENCSEER A. Long-term Characterization of Urban PM₁₀ in Hungary. *Aerosol and Air Quality Research*. **21**, 210048, **2021**.
 53. VARDOULAKISA S., PAVLOS K. Sources and factors affecting PM₁₀ levels in two European cities: Implications for local air quality management. *Atmospheric Environment*. **42** (17), 3949, **2008**.
 54. UNAL Y.S., TOROS H., DENIZ A., INCECIK S. Influence of meteorological factors and emission sources on spatial and temporal variations of PM₁₀ concentrations in Istanbul metropolitan area. *Atmospheric Environment*. **45** (31), 5504, **2021**.
 55. BODOR Z., BODOR K., KERESZTESI Á. SZÉP R. Major air pollutants seasonal variation analysis and long-range transport of PM₁₀ in an urban environment with specific climate condition in Transylvania (Romania). *Environmental Science and Pollution Research*. **27**, 38181, **2020**.
 56. SIRITHIAN D., THANATRAKOLSRI P. Relationships between Meteorological and Particulate Matter Concentrations (PM_{2.5} and PM₁₀) during the Haze Period in Urban and Rural Areas, Northern Thailand. *Air, Soil and Water Research*. **15**, **2022**.
 57. NISTOR A., NISTOR B., MIHAILĂ D., BISTRICEAN P-A. Atmospheric levels of PM10 in relation to meteorological elements. Case study: Moldova Region, Romania. *Georeview*. **33** (2), 95, **2023**.
 58. LIU Y., ZHOU Y., LU J. Exploring the relationship between air pollution and meteorological conditions in China under environmental governance. *Scientific Reports*. **10**, 14518, **2020**.
 59. YADAV S., PRAVEEN O.D., SATSANGI P.G. The effect of climate and meteorological changes on particulate matter in Pune, India. *Environmental Monitoring and Assessment*. **187**, 1, **2015**.
 60. CETIN Z.I. The effects of Sinop province relative humidity values on bioclimatic comfort and urban and landscape planning. *Journal of Design for Resilience in Architecture and Planning*. **5** (3), 401, **2024**.
 61. ÖZDEMİR U., TANER S. Impacts of meteorological factors on PM₁₀: Artificial neural networks (ANN) and multiple linear regression (MLR) approaches. *Environmental Forensics*. **15** (4), 329, **2014**.
 62. KLIENGCHUAY W., WORAKHUNPISET S., LIMPANONT Y., MEEYAI A.C., TANTRAKARNAPA K. Influence of the meteorological conditions and some pollutants on PM₁₀ concentrations in Lamphun, Thailand. *Journal of Environmental Health Science and Engineering*. **19**, 237, **2021**.
 63. COSKUNER G., SASSIM M.G., MUNIR S. Characterizing temporal variability of PM_{2.5}/PM₁₀ ratio and its relationship with meteorological parameters in Bahrain, *Environmental Forensics*. **19** (4), 315, **2018**.
 64. MALTARE N.N., VAHORA S., ANI K. Seasonal analysis of meteorological parameters and air pollutant concentrations in Kolkata: An evaluation of their relationship. *Journal of Cleaner Production*. **436**, 140514, **2024**.
 65. SHIHAB A.S. Investigating the relationship between air pollutants and meteorology: A canonical correlation analysis. *Polish Journal of Environmental Studies*. **31** (6), 5841, **2022**.