

Original Research

The Appropriate Reduction of Nitrogen Fertilization Enhances Soil Quality without Compromising Fruit Yield and Quality in a Bayberry Orchard

Yichao Chen^{1, 2, 3#}, Lu Xiang^{4#}, Fei Li^{1, 2, 3}, Yaojun Chang^{1, 2, 3}, Hongao Yu^{1, 2, 3},
Jing Zhang^{1, 2, 3}, Zhiliang Xie^{1, 2, 3*}

¹Wenzhou Vocational College of Science and Technology, Wenzhou, 325006, China

²Wencheng Institute of Modern Agriculture and Healthcare Industry, Wenzhou, 325006, China

³Southern Zhejiang Key Laboratory of Crop Breeding, Wenzhou, 325006, China

⁴College of Environment and Resources, Zhejiang A&F University, Hangzhou 311300, China

Received: 13 February 2025

Accepted: 22 April 2025

Abstract

The reduction of chemical fertilizers is a management practice gradually being adopted in long-term fertilized soils. However, the effects of chemical fertilizer reduction on soil health have remained underexplored in orchard systems. Through a two-year field experiment, we investigated how the reduction of chemical fertilizers affects soil chemical properties, the soil quality index (SQI), and fruit yield and quality. The results indicated that chemical fertilizer reduction significantly increased soil pH and organic matter, while it decreased the available nitrogen and phosphorus contents in 2022 and 2023. The reduction of chemical P fertilizer enhanced the availability of Si, S, Mn, and Cu, while it had no impact on the availability of Mo, Fe, and B. Chemical fertilization reduction did not decrease fruit yield and quality, except for the 1/3N treatment. Moreover, the SQI for the N and 2/3N treatments was higher. The random forest model demonstrated that SQI and soil pH were the most important driving factors regulating fruit yield in response to chemical fertilizer reduction practices. Our study suggests that the appropriate reduction of chemical fertilizers can enhance SQI and the availability of micronutrients without decreasing fruit yield and quality, which may have direct implications for soil health.

Keywords: long-term fertilization, management practice, chemical fertilizer reduction, soil health, fruit quality

[#]These two authors contributed equally to this paper

*e-mail: xiezhiliang@wzvcst.edu.cn

Tel.: +86 0577 88412441

Introduction

Agrosystems are frequently subjected to various fertilizer inputs aimed at enhancing crop yield and quality, predominantly nitrogen (N) and phosphorus (P) [1]. Long-term application of chemical fertilizers results in the accumulation of available nutrients in the soil [2], potentially hindering plant productivity [3]. This practice increases farmers' costs and heightens the risk of surface runoff contamination by N and P [4]. Furthermore, soil acidification from chemical fertilization adversely affects plant growth and may reduce crop yields. Reducing chemical fertilizer usage is crucial for improving soil properties [3]. Previous studies have indicated that decreasing fertilizer application does not negatively impact the yield and quality of wheat [5]. In contrast to conventional crop production systems, limited research has examined the effects of reduced chemical fertilizer use on soil quality, fruit yield, and quality within orchard forest ecosystems [6].

Soil quality is a multifactorial indicator that reflects the potential of soil to sustain its ecological services and productivity [7]. The Soil Quality Index (SQI) is a comprehensive measure of soil degradation, quality, and health [8, 9]. Most studies have primarily focused on assessing soil quality for only one year or one growing season [7, 10, 11], resulting in a limited understanding of the dynamic changes in soil quality through continuous evaluation. However, insufficient attention has been given to how chemical fertilizer reduction affects soil quality and its linkages to fruit yield and quality within orchard agroecosystems.

Bayberry (*Myrica rubra*) is widely distributed across tropical and subtropical regions globally, including countries such as China, Japan, the United States, and Brazil [12, 13]. In China, bayberry orchards span over 334,000 hectares and provide significant economic benefits to farmers [12]. Agroforestry management practices, including chemical fertilization [14] and the removal of understory vegetation, are commonly employed in bayberry orchards to enhance the growth of bayberry plants and fruit yield. However, these practices, particularly chemical fertilization, have resulted in several adverse effects on soil health and quality, such as soil acidification [2] and soil organic carbon loss [15], which undermine the sustainable development of the bayberry industry. Consequently, reducing fertilizer usage presents a viable and effective strategy to mitigate these negative impacts [4]. Nonetheless, it remains unclear how much fertilizer should be reduced and what the appropriate proportions of the essential nutrients involved are without compromising fruit yield and quality.

This study examines soil properties and evaluates soil quality after reducing chemical fertilization (N and P) in a bayberry orchard. Additionally, we assessed fruit yield and quality to explore their relationships with soil quality and health, ultimately aiming to identify the most

appropriate ratio for chemical fertilization reduction. We hypothesized the following: (1) a suitable decrease in chemical fertilization does not negatively impact fruit yield and quality while enhancing soil quality, and (2) variations in soil quality are associated with changes in fruit yield and quality resulting from the reduction of chemical fertilization.

Materials and Methods

Study Site

The experiment was conducted in Huangtan Township, Wenzhou (27°73'N, 119°99'E), located in the southeastern region of China. The study site experiences a monsoonal subtropical climate characterized by an annual mean temperature of 17.1°C and total precipitation of 1,680 mm. The site is situated at an elevation of approximately 200 m and has an average of 245 frost-free days and 2,000 hours of sunlight annually. The soil at the site has a clay loam texture derived from gneiss and is classified as Ferralsol [16].

In July 2021, we designated a section of a bayberry forest orchard for this field study. Traditional fertilization practices for these orchards involve the application of urea (450 kg ha⁻¹), superphosphate (100 kg ha⁻¹), and potassium sulphate (450 kg ha⁻¹). The planting density of fruit trees is 300 trees per hectare. When fertilizing, dig a ring-shaped ditch around each fruit tree and apply the corresponding fertilizer. Typically, 20% of N fertilizer and 60% of K fertilizer were used in late May, while the remaining 80% of N fertilizer and 40% of K fertilizer were applied in June. All P fertilizers were applied in November. Furthermore, the orchard owner routinely clears the understory vegetation annually. During our study period, no additional fertilizers were applied; however, the understory vegetation was removed each year.

Experimental Design, Management Practices, and Sample Collection

The study used a randomized block design comprising 9 treatments (3 gradient levels of N fertilizer reduction × 3 gradient levels of P fertilizer reduction) with 3 replicates for 27 plots. Each plot measures 100 m² (10 m × 10 m). Three gradient levels of N fertilizer reduction were applied with 450 kg ha⁻¹ (N), 300 kg ha⁻¹ (2/3N), and 150 kg ha⁻¹ (1/3N) urea. Three gradient levels of P fertilizer reduction were applied with 100 kg ha⁻¹ (N), 67 kg ha⁻¹ (2/3P), and 33 kg ha⁻¹ (1/3P) superphosphate (Table 1). All plots were applied 450 kg ha⁻¹ potassium sulphate as K fertilizer.

In July 2021, March and July 2022, and March 2023, the chemical N and P fertilizers were uniformly applied to their respective treatment plots within 0–20 cm of the soil layer, with 50% of the application amount calculated based on plot area (Table 1). In each plot,

Table 1. Detailed fertilization practices of each treatment.

| Treatment | Urea (kg ha ⁻¹) | Superphosphate (kg ha ⁻¹) |
|-----------|-----------------------------|---------------------------------------|
| N-P | 450 | 100 |
| N-2/3P | 450 | 67 |
| N-1/3P | 450 | 33 |
| 2/3N-P | 300 | 100 |
| 2/3N-2/3P | 300 | 67 |
| 2/3N-1/3P | 300 | 33 |
| 1/3-NP | 150 | 100 |
| 1/3N-2/3P | 150 | 67 |
| 1/3N-1/3P | 150 | 33 |

five randomly-located soil cores (0-20 cm) separated by at least 2 m were collected and mixed thoroughly to form a composite sample. The mixed samples were air-dried at room temperature and were passed through a 2-mm sieve for soil nutrients and other properties analysis.

Analysis of Soil Chemical Properties

The soil's chemical properties were measured following the method described by Lu [17]. In detail, soil pH was determined by a glass electrode (Mettler-Toledo, Switzerland) with a soil-to-water ratio of 1:2.5. Soil organic C (SOC) and total nitrogen (TN) were determined by wet digestion using K₂CrO₇ oxidation and the Kjeldahl procedure, respectively. Soil available phosphorus (P) was determined according to the procedure described by Bray et al. [18]. Soil available N was determined by the hot alkaline permanganate method.

Soil available microelement (including Si, S, Mo, B, Fe, Mn, Cu, and Zn) contents were extracted by 0.025 M DTPA solution, and the concentrations of the elements in the extractants were determined using ICP-OES (Optima 8300, PerkinElmer, Waltham, MA, USA) [19]. Exchangeable Ca and Mg concentrations were extracted by NH₄C₂H₃O₂ (pH = 7.0) and determined using ICP-OES.

Analysis of Fruit Yield and Quality

The fruit yield of bayberry in each plot was weighted and recorded between May and June every year. At the same time, the responses of chemical fertilizer reduction on bayberry fruit quality were evaluated by measuring soluble solids and titratable acidity. The content of soluble solids was determined using a portable reflectometer [20]. Titratable acidity was determined by the indicator-based titration method using 0.1 M NaOH-methylene blue indicator [21].

Calculation of Soil Quality Index

Soil quality index (SQI) was calculated using minimum datasets (MDS) and principal component analysis (PCA) methods [22]. In detail, PCA was employed to identify the indicators of MDS while minimizing the information loss [23]. The principal components having eigenvalues >1 and explaining >5% were considered [11]. Select the maximum Norm value from each group to enter the MDS [11], where the Norm value is the vector magnitude in the multi-dimensional space composed of principal components. The higher Norm value represents a more significant comprehensive load of the indicator on all principal components, indicating a richer reflection of soil quality information [7, 10]. Considering the important role of soil organic matter and N, P, and K elements in soil quality and ecological function, available N, P, and K, and soil organic matter were included in MDS [24, 25].

To eliminate the magnitudes of different soil indicators, they were transformed by the linear scoring method to score the values ranging from 0 to 1 [26].

$$StPi = \frac{Pi - \min(Pi)}{\max(Pi) - \min(Pi)}$$

Where StPi, Pi, min(Pi), and max(Pi) indicate the standardized index, actual observed value, maximum value observed in all samples, and minimum value observed in all samples, respectively.

Finally, the radar maps of soil quality were established using the standardized indexes [27, 28], and the coverage area reflects soil quality. SQI was calculated as follows:

$$SQI = 0.5 \sum_i^n StPi^2 \sin \frac{2\pi}{n}$$

Where SQI and n indicate the soil quality index and the number of indicators in MDS.

Statistical Analysis

Significant analysis was performed by SPSS 26.0 (IBM Inc., Chicago, IL, USA). Two-way analysis of variance (ANOVA) was explored to determine the significant changes in the soil's physical and chemical properties, fruit yield and quality, and SQI. One-way ANOVA was employed to assess the differences under the same N or P fertilizer reduction level. The correlation coefficients involving the correlative relationships between SQI and fruit yield and quality were obtained using the Spearman correlation. Random forest analysis was performed to estimate the importance of soil properties for explaining SQI and fruit yield by using "rfPermute" and "A3" packages in R [29]. Structural equation modeling (SEM) was further constructed

to identify how chemical fertilizer reduction affected SQI and fruit yield based on the linkages among soil available nutrient content. An adequate fit was indicated by the chi-square test ($df>5$; $P>0.05$), a low root-mean-square error of approximation (RMSEA) (<0.05), and a high goodness-of-fit index (GFI) [30].

Results

Changes in Soil Chemical Properties

Chemical N and P fertilizer reduction significantly affected soil chemical properties, including common nutrient and micro-element contents. The reduction of N fertilizer significantly increased soil pH and organic matter and decreased available N, available P, and total N content ($P<0.05$, Table 2). The reduction of P fertilizer did not affect soil pH and TN, but significantly

decreased soil available P and available N and increased soil organic matter ($P<0.05$). Additionally, the reduction of N and P fertilizers did not consistently affect soil exchange Ca and Mg in 2022 and 2023 (Fig. 1).

Compared to the N treatment, the 2/3N treatment had higher available S, Mo, Fe, Mn, Zn, and B contents (Table 3), while the 1/3N treatment had lower available Si, S, Fe, Cu, and Zn contents. Chemical P fertilizer reduction significantly increased available Si, S, Mn, Cu, and Zn contents, and there was no significant difference in available Fe and B contents (Table 3).

Changes in Fruit Yield and Quality

In 2022, the chemical N fertilizer reduction had significant effects on fruit yield and had no significant changes on fruit diameter, edible rate (Fig. 2), soluble solid and titratable acidity (Fig. 3), while the chemical P fertilizer significantly changed fruit yield and soluble

Table 2. Changes in soil nutrient content affected by the reduction of N and P fertilization.

| Treatment | | Soil pH | Organic matter (g kg ⁻¹) | Total N (g kg ⁻¹) | Available N (mg kg ⁻¹) | Available P (mg kg ⁻¹) |
|----------------|-----------|--------------|--------------------------------------|-------------------------------|------------------------------------|------------------------------------|
| 2022 | N-P | 4.68±0.07aB | 25.60±4.71bA | 2.64±0.32aA | 194.93±6.12aA | 139.17±28.29aAB |
| | N-2/3P | 4.75±0.02aB | 28.29±4.79abB | 2.78±0.26aA | 153.20±14.63bA | 116.39±8.34abA |
| | N-1/3P | 4.75±0.09aC | 34.52±0.44aA | 2.51±0.36aA | 128.77±4.43cA | 97.71±14.89bA |
| | 2/3N-P | 5.00±0.02bA | 30.20±4.59aA | 2.22±0.30aA | 174.28±6.39aB | 160.47±14.91aA |
| | 2/3N-2/3P | 5.01±0.08bA | 35.27±0.50aA | 2.62±0.22aA | 115.89±14.04bB | 112.29±6.96bA |
| | 2/3N-1/3P | 5.14±0.08aA | 32.47±2.71aA | 2.33±0.48aA | 122.07±6.50bA | 77.39±6.76cAB |
| | 1/3N-P | 4.88±0.07abA | 28.85±2.35abA | 1.43±0.32aB | 100.10±11.55aC | 132.38±9.47aB |
| | 1/3N-2/3P | 4.78±0.06bB | 26.58±3.81bB | 1.63±0.20aB | 96.64±12.20aB | 87.51±3.26bB |
| | 1/3N-1/3P | 4.94±0.12aB | 34.34±4.12aA | 1.51±0.11aB | 111.93±11.79aA | 62.20±2.31cB |
| 2023 | N-P | 4.60±0.05aA | 22.20±0.72bB | 2.06±0.06bA | 171.67±5.69bA | 136.07±5.89aA |
| | N-2/3P | 4.57±0.05aB | 25.43±1.17bB | 2.43±0.04aA | 199.00±8.89aA | 113.23±6.83bA |
| | N-1/3P | 4.67±0.05aB | 30.40±3.44aA | 1.97±0.06bA | 147.00±7.94cA | 94.40±8.32cA |
| | 2/3N-P | 4.66±0.08aA | 26.33±1.72bA | 1.59±0.10aB | 135.00±10.82aB | 143.43±11.73aA |
| | 2/3N-2/3P | 4.66±0.04aB | 30.17±2.74aA | 1.55±0.07abB | 136.67±5.13aB | 97.60±13.66bB |
| | 2/3N-1/3P | 4.71±0.07aB | 28.93±3.17abAB | 1.47±0.04bB | 131.67±5.13aA | 90.93±4.21bA |
| | 1/3N-P | 4.72±0.09bA | 26.37±0.60aA | 1.32±0.03bC | 105.67±11.02aC | 130.50±7.41aA |
| | 1/3N-2/3P | 4.93±0.14aA | 22.67±0.67aB | 1.37±0.06abC | 103.67±11.50aC | 93.00±7.07bB |
| | 1/3N-1/3P | 4.89±0.08aA | 26.10±1.25aB | 1.44±0.06aB | 108.33±6.81aB | 75.73±5.92cB |
| N reduction | | ** | * | ** | ** | ** |
| P reduction | | ns | ** | ns | ** | ** |
| N *P reduction | | ns | ** | ns | ** | ns |

*and ** indicated significance based on two-way ANOVA at 0.05 and 0.01 levels, respectively. Ns, no significance. Different letters in the same column indicate significant differences at the $P<0.05$ level. Lowercase letters denote statistically significant differences ($P<0.05$) in P fertilizer reduction effects on soil nutrient contents under each N fertilizer reduction levels, whereas uppercase letters represent N fertilizer reduction impacts under each P fertilizer reduction levels.




Fig. 1. Soil exchange Ca (a and c) and Mg (b and d) contents and their response to chemical fertilizer reduction in 2022 and 2023. Lowercase letters denote statistically significant differences ($P < 0.05$) in P fertilizer reduction effects on soil nutrient contents under each N fertilizer reduction levels, whereas uppercase letters represent N fertilizer reduction impacts under each P fertilizer reduction levels.

solid and not impacted fruit diameter, edible rate, and titratable acidity. In 2023, compared to 1/3N treatment, N and 2/3N treatment had a higher fruit yield and titratable acidity content (Fig. 3). N fertilizer reduction did not affect fruit diameter, edible rate, or soluble solid contents. Furthermore, the chemical P fertilizer reduction did not affect fruit yield and titratable acidity contents, while only significantly increasing soluble solid contents in 1/3P treatment.

Changes in Soil Quality Indexes

In this study, the eigenvalue of PC1-PC6 was >1 , and cumulatively explained the variance of 82.442% in the whole data using PCA analysis (Table 4). The soil pH, soil organic matter, bulk density, moisture, exchange Mg, and available N, K, P, Mn, and B were selected as important soil quality evaluation indicators by using MDS and PCA analysis (Fig. 4). Compared to the N treatment, the SQI significantly improved under the 2/3N treatment in 2022 ($P < 0.05$), while there is no significant difference between the N treatment and the 2/3N treatment in 2023 (Fig. 4g) and h)). In both 2022 and 2023, compared to the N treatment, the SQI largely decreased under the 1/3N treatment. The reduction of chemical P fertilizer had no significant impacts on SQI in 2022 and 2023 ($P > 0.05$).

The Relationships between Soil Chemical Properties, Soil Quality, and Fruit Quality

The SQI was strongly and positively correlated with fruit yield in both 2022 and 2023 (Fig. 5a)). The SQI had no significant correlations with the fruit quality, including soluble solids and titratable acidity in 2022 and 2023, as expected for SQI and fruit titratable acidity exerting a close correlation (Fig. 5b) and c)). Random forest analysis showed that soil chemical properties (including exchange Mg, moisture, available N, soil organic carbon, available K, pH, available Mn, available B, and available P) were significantly important for SQI variation. All the indicators explain 73.4% of the SQI variation. Furthermore, soil pH and SQI showed the highest importance for fruit yield variation, followed by available N, exchange Mg, and available K. All the indicators explain 49.8% of the fruit yield variation. The SEM proved a good fit to the data (Chi-square = 15.328, df = 10, GFI = 0.924, RMSEA = 0.04) and accounted for 73.7% of the variation in SQI, and 49.5% of the variation in fruit yield (Fig. 6). According to the model, the chemical N fertilizer reduction significantly and directly influenced soil pH, available N, and available B. SQI was affected by soil pH, available N, available B, and exchange Mg. Moreover, fruit yield is impacted by soil pH, exchange Mg and SQI.

Table 3. Changes in micro-elements (including Si, B, Mo, Fe, Mn, Cu, Zn) content in the bayberry orchard in 2022 and 2023.

| Treatment | Available Si (mg kg ⁻¹) | Available B (mg kg ⁻¹) | Available Mo (μg kg ⁻¹) | Available S (mg kg ⁻¹) | Available Fe (mg kg ⁻¹) | Available Mn (mg kg ⁻¹) | Available Cu (mg kg ⁻¹) | Available Zn (mg kg ⁻¹) |
|-----------|--|---------------------------------------|--|---------------------------------------|--|--|--|--|
| 2022 | N-P | 131.49±4.17bC | 7.02±0.24aA | 72.83±58.45aA | 65.27±6.74bB | 10.41±2.35bB | 1.14±0.10cC | 3.64±0.12bB |
| | N-2/3P | 248.95±5.90aC | 7.43±0.28aA | 89.28±61.16aA | 87.23±9.80bB | 16.95±1.66aB | 3.11±0.03bB | 8.48±0.67aC |
| | N-1/3P | 249.87±14.53aA | 7.47±1.04aA | 91.01±51.88aB | 128.97±5.18aB | 16.15±1.65aA | 3.44±0.02aA | 4.30±0.72cC |
| | 2/3N-P | 175.50±12.58cA | 6.68±0.51aA | 70.86±65.21bA | 164.25±3.74aA | 16.15±1.18bA | 2.99±0.05bA | 7.03±0.48cA |
| | 2/3N-2/3P | 303.02±3.97aA | 6.80±0.08aA | 191.29±94.76bA | 143.97±6.06bA | 23.09±1.42aA | 3.40±0.06aA | 12.47±2.17aA |
| | 2/3N-1/3P | 199.21±11.99bB | 6.97±0.84aA | 312.89±36.36aA | 145.45±9.37bA | 16.24±1.64bA | 1.63±0.10cC | 7.62±1.08bA |
| | 1/3N-P | 158.58±6.93cB | 6.67±0.37aA | 29.91±18.34aA | 62.90±4.07bB | 8.90±0.65bB | 1.42±0.08bB | 2.51±0.88cC |
| | 1/3N-2/3P | 274.14±6.55aB | 6.56±0.47aA | 96.89±58.72aA | 68.01±3.96bC | 15.38±2.24aB | 2.63±0.05aC | 9.93±2.09aB |
| | 1/3N-1/3P | 201.02±3.14bB | 6.72±0.34aA | 74.55±105.92aB | 83.39±18.25aC | 17.32±2.28aA | 2.63±0.04aB | 7.23±0.92bA |
| | N-P | 183.83±7.86aC | 6.38±0.26aB | 38.00±4.35aA | 70.57±1.56aC | 16.70±1.47aA | 2.62±0.04aB | 7.35±0.21bA |
| | N-2/3P | 173.60±5.63abC | 6.39±0.19aB | 42.00±12.53aA | 72.47±1.55aC | 14.80±1.40aA | 2.63±0.06aC | 10.01±2.13aB |
| | N-1/3P | 160.87±12.32bB | 6.71±0.25aB | 39.17±1.73aA | 72.80±6.94aB | 9.30±0.75bB | 1.38±0.03bC | 2.73±0.25cC |
| 2023 | 2/3N-P | 199.50±9.47bB | 6.83±0.06aAB | 28.67±1.15aB | 113.97±10.18aB | 15.43±0.31aA | 1.62±0.05cC | 7.66±1.15bA |
| | 2/3N-2/3P | 308.43±11.60aA | 6.83±0.15aAB | 38.67±8.50aA | 101.30±2.95bB | 18.40±4.56aA | 3.35±0.02aA | 12.52±2.11aA |
| | 2/3N-1/3P | 164.60±2.60cB | 6.73±0.21aB | 32.33±6.81aA | 114.43±7.19aA | 16.20±1.31aA | 2.95±0.04bA | 7.05±1.08bA |
| | 1/3N-P | 249.27±10.52aA | 7.27±0.65aA | 28.67±9.07aB | 140.50±4.86aA | 15.53±1.03aA | 3.43±0.44aA | 4.48±1.18bB |
| | 1/3N-2/3P | 251.40±5.27aB | 7.01±0.10aA | 25.33±9.23aB | 143.25±5.43aA | 15.97±0.45aA | 3.13±0.74aB | 8.59±1.12aC |
| | 1/3N-1/3P | 239.00±10.82aA | 7.40±0.46aA | 15.67±5.13aB | 121.10±4.55bA | 10.10±1.25bB | 2.12±0.32bB | 3.84±0.20cB |
| N | ** | ** | * | ** | ** | ** | ** | ** |
| P | * | * | ns | ns | ** | ** | ** | ** |
| N*P | ** | ns | ns | ** | ns | ** | ** | ** |

*and ** indicated significance based on two-way ANOVA at 0.05 and 0.01 levels, respectively. Ns, no significance. Different letters in the same column indicate significant differences at the $P<0.05$ level. Lowercase letters denote statistically significant differences ($P<0.05$) in P fertilizer reduction effects on micro-elements contents under each N fertilizer reduction levels, whereas uppercase letters represent N fertilizer reduction impacts under each P fertilizer reduction levels.




Fig. 2. Impacts of chemical fertilizer reduction on fruit yield (a and d) and quality, including fruit diameter (b and e) and edible rate (c and f) within a bayberry orchard in 2022 and 2023. Lowercase letters denote statistically significant differences ($P < 0.05$) in P fertilizer reduction effects on soil nutrient contents under each N fertilizer reduction levels, whereas uppercase letters represent N fertilizer reduction impacts under each P fertilizer reduction levels.




Fig. 3. Impacts of chemical fertilizer reduction on fruit quality, including soluble solid (a and c) and titratable acidity (b and d) within a bayberry orchard in 2022 and 2023. Lowercase letters denote statistically significant differences ($P < 0.05$) in P fertilizer reduction effects on soil nutrient contents under each N fertilizer reduction levels, whereas uppercase letters represent N fertilizer reduction impacts under each P fertilizer reduction levels.

Table 4. Load matrix and Norm value of each soil property with principal component (PC) analysis.

| Soil indicators | PC1 | PC2 | PC3 | PC4 | PC5 | PC6 | Group | Norm value |
|--------------------------|--------|--------|--------|--------|--------|--------|-------|------------|
| Soil moisture | 0.385 | 0.656 | -0.035 | 0.459 | 0.256 | 0.179 | 2 | 1.690 |
| Bulk density | 0.238 | -0.123 | -0.518 | 0.406 | -0.040 | 0.607 | 6 | 1.309 |
| pH | 0.048 | -0.616 | 0.426 | 0.356 | 0.165 | -0.276 | 2 | 1.487 |
| Soil organic matter | 0.429 | -0.429 | 0.621 | 0.186 | 0.163 | 0.104 | 3 | 1.655 |
| Total N | 0.418 | 0.266 | 0.520 | -0.365 | 0.374 | 0.026 | 1 | 1.513 |
| Available N | 0.333 | 0.785 | 0.136 | -0.17 | 0.106 | -0.107 | 2 | 1.729 |
| Available P | -0.005 | 0.696 | 0.023 | 0.373 | 0.376 | 0.145 | 2 | 1.499 |
| Available K | 0.504 | 0.023 | 0.187 | 0.001 | -0.689 | -0.039 | 5 | 1.446 |
| Cation exchange capacity | 0.350 | 0.624 | 0.264 | 0.056 | -0.525 | -0.060 | 2 | 1.630 |
| Exchangeable Ca | 0.704 | 0.172 | -0.399 | 0.056 | 0.144 | -0.377 | 1 | 1.819 |
| Exchangeable Mg | 0.730 | 0.327 | -0.338 | 0.349 | -0.048 | -0.159 | 1 | 1.929 |
| Available Si | 0.656 | -0.285 | -0.247 | -0.459 | 0.144 | 0.277 | 1 | 1.784 |
| Available S | 0.682 | -0.186 | 0.260 | 0.191 | -0.307 | 0.254 | 1 | 1.734 |
| Available Mo | 0.367 | -0.449 | 0.638 | 0.223 | 0.196 | 0.117 | 3 | 1.625 |
| Available B | 0.203 | 0.394 | 0.327 | -0.537 | -0.048 | 0.395 | 4 | 1.311 |
| Available Fe | 0.777 | 0.233 | 0.305 | 0.013 | 0.006 | -0.240 | 1 | 1.917 |
| Available Mn | 0.810 | -0.426 | -0.177 | -0.036 | 0.006 | 0.044 | 1 | 2.051 |
| Available Cu | 0.672 | -0.111 | -0.390 | -0.348 | 0.238 | -0.096 | 1 | 1.760 |
| Available Zn | 0.629 | -0.526 | -0.348 | -0.038 | -0.013 | -0.093 | 1 | 1.851 |
| Eigenvalue | 5.260 | 3.705 | 2.544 | 1.675 | 1.400 | 1.081 | - | - |
| Explanation (%) | 27.682 | 19.498 | 13.389 | 8.815 | 7.37 | 5.687 | - | - |
| Cumulative variance (%) | 27.682 | 47.181 | 60.570 | 69.384 | 76.755 | 82.442 | - | - |

Discussion

The Reduction of N and P Fertilization Affected Soil Available Nutrient Contents and Improved Soil Quality in the Bayberry Orchard

Our results showed that chemical N fertilizer reduction increased soil pH (Table 2). Consistent with previous studies [2, 15], applying chemical N fertilizer significantly decreased soil pH due to ammonium-based fertilizers that release hydrogen ions when converted to nitrate by soil microorganisms. We further found that reducing chemical fertilizer enhanced soil organic matter (Table 2), possibly due to improved greenhouse gas emissions and organic carbon mineralization caused by long-term chemical fertilization [31, 32]. In addition, the reduction of chemical fertilizer significantly decreased soil available N and P (Table 2), indicating that long-term fertilization resulted in the accumulation of available nutrients in the soil with low efficiency of nutrient use [33]. The 2/3N treatment increased the availability of Mg, Ca (Fig. 1), and micro-elements (Table 3). This phenomenon could be explained by

the enhancement of soil pH [34], while there was no additional corresponding macro- and micro-elements input to the soil. Additionally, the input of inorganic P may largely affect the availability of Mg, Ca, and micro-elements due to the association of PO_4^{3-} and metal ions [35].

This study developed a comprehensive SQI by integrating key soil property indicators to systematically evaluate the effects of reducing chemical N and P fertilizer application on soil quality. We illustrated that soil pH, organic matter, bulk density, moisture, exchangeable Mg, and available N, P, K, Mn, and B from the whole data set, as the MDS could be used to evaluate soil quality under different chemical fertilizer reduction levels (Fig. 3). The results of SQI were higher with 2/3N applied treatment and lower within 1/3N applied treatment, compared to the whole N applied treatment (Fig. 3g) and h)). Consistent with previous studies [7, 36], long-term chemical fertilization resulted in soil degradation, and appropriate reduction of chemical fertilizer would improve soil quality. Nevertheless, it should be noted that the SQIs under 1/3N treatment were significantly decreased, which indicated it also had some




Fig. 4. Radar map of soil quality under chemical fertilizer reduction practices in 2022 (a, b, and c) and 2023 (d, e and f); and soil quality index (SQI) (g and h) response to chemical fertilizer reduction practices. Lowercase letters denote statistically significant differences ($P < 0.05$) in P fertilizer reduction effects on soil nutrient contents under each N fertilizer reduction levels, whereas uppercase letters represent N fertilizer reduction impacts under each P fertilizer reduction levels.




Fig. 5. Correlation analysis of soil quality index (SQI) with fruit yield (a) and quality (b and c) in a bayberry orchard; random forest model of SQI (d) and fruit yield (e) with soil properties.




Fig. 6. Structural equation model (SEM) illustrating relationships between soil properties (pH, available N, available B, and exchange Mg), soil quality index (SQI), and fruit yield. Red and blue lines indicate positive and negative effects, respectively. Correlation coefficients and significant level (*, $P < 0.05$; **, $P < 0.01$) are shown with lines.

negative effects on soil quality (Fig. 3). This may be due to the limited input of N and the inability to meet the demand for plant uptake [27]. Therefore, the effects of chemical N fertilizer reduction on soil quality depend on the reduction rates. It is recommended to combine with organic fertilizers or return straw to the soils [37] if a significant reduction of chemical N fertilizer is required.

Furthermore, reducing P fertilizer did not affect soil quality (Fig. 3). This could be attributed to the weaker effects of chemical P fertilizer reduction on soil nutrients and chemical properties than the reduction of N

chemical fertilizer (Table 2). Additionally, compared to the availability of P in southeast China [38], the available P contents in the bayberry orchard were relatively high; thereby, the reduction of P fertilizer did not affect soil quality. Notably, the differential mobility of N and P in soil systems may also explain their contrasting impacts: N losses via leaching or volatilization can rapidly deplete soil N pools, whereas P tends to remain sequestered in less labile forms, buffering against abrupt changes in availability [38]. The reduction of chemical N fertilizer, rather than that of chemical P fertilizer,




Fig. 7. Conceptual graph to elucidate effects of chemical N fertilizer reduction on soil properties, soil quality index (SQI), and fruit yield and quality.

drove the changes in soil quality.

The Appropriate Reduction of Chemical N and P Fertilization did not Decrease Bayberry Yield and Quality

In the present study, the appropriate reduction of chemical N and P fertilization did not decrease fruit yield and quality in a bayberry orchard (Fig. 2). Firstly, this could be explained by the improvement of soil pH. It is well documented that soil pH is an important factor affecting plant growth [39, 40], with a low pH reducing plant resistance to diseases [41], as well as root growth, which is detrimental to fruit yield and quality. This mechanism could be evidenced by the SEM and random forest model (Fig. 4 and 5). Moreover, a global meta-analysis showed that the average N-use efficiency was only 39% [42]. In long-term fertilized systems, residual soil N often accumulates, creating a buffer that sustains plant growth even with reduced inputs. Thus, the N was not the limiting factor of fruit yield and quality in long-term fertilized agroecosystems. Additionally, the availability of Mg, Ca, and microelements enhanced under 2/3N treatment could support fruit yield and quality. Li et al. [43] reported that the content of microelements could be the limiting factor of plant growth. Mg is integral to chlorophyll synthesis and photosynthetic efficiency, while Ca strengthens cell walls and prolongs fruit postharvest quality [44]. Micronutrients like Zn and Fe, which act as enzyme cofactors, may have bolstered metabolic processes critical for fruit development. The results of SEM supported the idea that available B had a significant contribution to fruit yield (Fig. 6). Furthermore, plants in N-reduced systems may activate stress-responsive pathways, elevating the synthesis of secondary metabolites (e.g., phenolics, anthocyanins) that improve fruit antioxidant capacity and color [45]. Balanced micronutrient supply [44] under reduced N can sustain or even improve quality by alleviating hidden hunger. Thus, strategic N reduction aligns with “less is more” agronomy, where precision in nutrient delivery – rather than quantity – drives quality outcomes.

Soil quality, an indicator of sustainable soil management, can help sustain biological productivity, maintain environmental quality, and promote plant and animal health [36]. Our results showed that SQI was an important driving factor in fruit yield (Fig. 5). The high positive correlation that typically exists between soil quality and crop yield [46, 47] allowed us to evaluate the accuracy of assessment methods [28]. Our results revealed that there was a positive and significant correlation between fruit yield and SQI (Fig. 4), which indicated that PCA analysis and the MDS method could be used to assess soil quality in a fruit orchard with chemical fertilizer reduction practice. These results were similar to those of Cheng et al. [48] and Chen et al. [49] in the navel orange production systems of the red soil region of China and the agricultural land

management systems of Northeast China, respectively. However, the threshold of N reduction warrants careful consideration. While moderate reductions (e.g., 33.3% in our study) preserved yield, steeper cuts might risk inducing nutrient deficits, particularly in high-demand growth stages [15]. Site-specific factors, including soil type, climate, and crop genotype, further modulate these outcomes. Thus, dynamic SQI monitoring, coupled with real-time nutrient diagnostics, is essential for tailoring fertilizer regimes to local conditions. Overall, our study suggested that the appropriate chemical N fertilizer reduction did not decrease fruit yield and quality by improving soil quality and soil chemical properties in a subtropical bayberry orchard.

Conclusions

Our results showed that chemical fertilizer reduction significantly increased soil pH and microelement availability in a subtropical bayberry orchard. The appropriate reduction of N fertilization (2/3N) did not decrease bayberry yield and quality, while it increased the soil quality index (SQI) through the 2-year field experiments. However, the 1/3N treatment (two-thirds N reduction (Fig. 7)) unexpectedly decreased SQI, highlighting the non-linear relationship between fertilization intensity and soil health. The reduction of chemical P fertilizer had no significant effects on SQI and fruit quality. Furthermore, SQI was mainly driven by the availability of nutrients and soil pH, while fruit yield was driven primarily by SQI and soil pH. These results deepen our understanding of the effects of chemical fertilizer reduction on soil properties, fruit yield and quality, and soil health, as well as reveal the role of chemical fertilizer reduction in the sustainability of agriculture and forestry.

Acknowledgements

This work was supported by the Wencheng Science and Technology Plan Program (2021NKY08 and 2023NKY04).

Conflict of Interest

The authors declare no conflict of interest.

References

1. LI C., WEI Z., WANG X., MA X., YANG P., SHAN J., YAN X. Long-term fertilization regulates dissimilatory nitrate reduction processes by altering paddy soil organic carbon components. *Soil Use and Management*. **40** (1), e12994, 2024.
2. CHEN J., WU Q., LI S., GE J., LIANG C., QIN H., XU Q., FUHRMANN J.J. Diversity and function of soil

- bacterial communities in response to long-term intensive management in a subtropical bamboo forest. *Geoderma*. **354**, 113894, **2019**.
3. HAN J., DONG Y., ZHANG M. Chemical fertilizer reduction with organic fertilizer effectively improve soil fertility and microbial community from newly cultivated land in the Loess Plateau of China. *Applied Soil Ecology*. **165**, 103966, **2021**.
 4. YE L., ZHAO X., BAO E., LI J., ZOU Z., CAO K. Bio-organic fertilizer with reduced rates of chemical fertilization improves soil fertility and enhances tomato yield and quality. *Scientific Reports*. **10** (1), 177, **2020**.
 5. HE P., LI S., JIN J., WANG H., LI C., WANG Y., CUI R. Performance of an Optimized Nutrient Management System for Double-Cropped Wheat-Maize Rotations in North-Central China. *Agronomy Journal*. **101** (6), 1489, **2009**.
 6. COLOMBO F., MACDONALD C.A., JEFFRIES T.C., POWELL J.R., SINGH B.K. Impact of forest management practices on soil bacterial diversity and consequences for soil processes. *Soil Biology & Biochemistry*. **94**, 200, **2016**.
 7. CHEN Y., JIANG Z., OU J., LIU F., CAI G., TAN K., WANG X. Nitrogen substitution practice improves soil quality of red soil (Ultisols) in South China by affecting soil properties and microbial community composition. *Soil & Tillage Research*. **240**, 106089, **2024**.
 8. BÜNEMANN E.K., BONGIORNO G., BAI Z., CREAMER R.E., DE DEYN G., DE GOEDE R., FLESKENS L., GEISSEN V., KUYPER T.W., MÄDER P., PULLEMAN M., SUKKE W., VAN GROENIGEN J.W., BRUSSAARD L. Soil quality – A critical review. *Soil Biology & Biochemistry*. **120**, 105, **2018**.
 9. MUÑOZ-ROJAS M. Soil quality indicators: critical tools in ecosystem restoration. *Current Opinion in Environmental Science & Health*. **5**, 47, **2018**.
 10. WAN P., ZHOU Z., YUAN Z., WEI H., HUANG F., LI Z., LI F.-M., ZHANG F. Fungal community composition changes and reduced bacterial diversity drive improvements in the soil quality index during arable land restoration. *Environmental Research*. **244**, 117931, **2024**.
 11. LALITHA M., KALAISELVI B., DHARUMARAJAN S., ANIL KUMAR K.S., RAMESH KUMAR S.C., SRINIVASAN R., RAMAMURTHY V., HEGDE R. Determining soil quality indicators for alluvial plains in the semi-arid tropics. *Soil use and management*. **40** (1), e12929, **2024**.
 12. REN H., GUO H., SHAFIQU L. ISLAM M., ZAKI H.E.M., WANG Z., WANG H., QI X., GUO J., SUN L., WANG Q., LI B., LI G., RADWAN K.S.A. Improvement effect of biochar on soil microbial community structure and metabolites of decline disease bayberry. *Frontiers in Microbiology*. **14**, **2023**.
 13. SUN C., HUANG H., XU C., LI X., CHEN K. Biological Activities of Extracts from Chinese Bayberry (*Myrica rubra* Sieb. et Zucc.): A Review. *Plant Foods for Human Nutrition (Dordrecht)*. **68** (2), 97, **2013**.
 14. REN H., WANG H., YU Z., ZHANG S., QI X., SUN L., WANG Z., ZHANG M., AHMED T., LI B. Effect of Two Kinds of Fertilizers on Growth and Rhizosphere Soil Properties of Bayberry with Decline Disease. *Plants*. **10**, 2386, **2021**.
 15. JIN W., GE J., SHAO S., PENG L., XING J., LIANG C., CHEN J., XU Q., QIN H. Intensive management enhances mycorrhizal respiration but decreases free-living microbial respiration by affecting microbial abundance and community structure in Moso bamboo forest soils. *Pedosphere*. **34** (2), 508, **2024**.
 16. WORLD REFERENCE BASE FOR SOIL RESOURCES. A framework for international classification, correlation and communication. Food and Agriculture Organization of the United Nations, Rome, **2006**.
 17. LU R. Analytical Methods for Soils and Agricultural Chemistry. China Agricultural Science and Technology Press, Beijing. **2000**.
 18. BRAY R.H., KURTZ L.T. Determination of Total, Organic, and Available Forms of Phosphorus in Soils. *Soil Science*. **59** (1), 39, **1945**.
 19. GU S., YANG X., CHEN H., JEYAKUMAR P., CHEN J., WANG H. Crawfish shell- and Chinese banyan branch-derived biochars reduced phytoavailability of As and Pb and altered community composition of bacteria in a contaminated arable soil. *The Science of the Total Environment*. **865**, 161284, **2023**.
 20. CHEN L., LI X., PENG Y., XIANG P., ZHOU Y., YAO B., ZHOU Y., SUN C. Co-application of biochar and organic fertilizer promotes the yield and quality of red pitaya (*Hylocereus polyrhizus*) by improving soil properties. *Chemosphere*. **294**, 133619, **2022**.
 21. MENG L.-L., LIANG S.-M., SRIVASTAVA A.K., LI Y., LIU C.-Y., ZOU Y.-N., KUČA K., HASHEM A., FATHI ABD ALLAH E., WU Q.-S. Easily Extractable Glomalin-Related Soil Protein as Foliar Spray Improves Nutritional Qualities of Late Ripening Sweet Oranges. *Horticulturae*. **7** (8), 228, **2021**.
 22. ANDREWS S.S., MITCHELL J.P., MANCINELLI R., KARLEN D.L., HARTZ T.K., HORWATH W.R., PETTYGROVE G.S., SCOW K.M., MUNK D.S. On-Farm Assessment of Soil Quality in California's Central Valley. *Agronomy Journal*. **94** (1), 12, **2002**.
 23. ARMENISE E., REDMILE-GORDON M.A., STELLACCI A.M., CICCARESE A., RUBINO P. Developing a soil quality index to compare soil fitness for agricultural use under different managements in the Mediterranean environment. *Soil & Tillage Research*. **130**, 91, **2013**.
 24. RAMESH T., BOLAN N.S., KIRKHAM M.B., WIJESEKARA H., KANCHIKERIMATH M., SRINIVASA RAO C., SANDEEP S., RINKLEBE J., OK Y.S., CHOUDHURY B.U., WANG H., TANG C., WANG X., SONG Z., FREEMAN II O.W. Chapter One - Soil organic carbon dynamics: Impact of land use changes and management practices: A review. In *Advances in Agronomy*, D.L. Sparks Ed. Academic Press: **156**, 1, **2019**.
 25. ZÖRB C., SENBAYRAM M., PEITER E. Potassium in agriculture – Status and perspectives. *Journal of Plant Physiology*. **171** (9), 656, **2014**.
 26. KUZUYAKOV Y., GUNINA A., ZAMANIAN K., TIAN J., LUO Y., XU X., YUDINA A., APONTE H., ALHARBI H., OVSEPYAN L., KURGANOVA I., GE T., GUILLAUME T. New approaches for evaluation of soil health, sensitivity and resistance to degradation. *Frontiers of Agricultural Science and Engineering*. **7** (3), 282, **2020**.
 27. CHEN Q., LIU Z., ZHOU J., XU X., ZHU Y. Long-term straw mulching with nitrogen fertilization increases nutrient and microbial determinants of soil quality in a maize–wheat rotation on China's Loess Plateau. *The Science of the Total Environment*. **775**, 145930, **2021**.
 28. CHERUBIN M.R., KARLEN D.L., FRANCO A.L.C., TORMENA C.A., CERRI C.E.P., DAVIES C.A., CERRI C.C. Soil physical quality response to sugarcane expansion in Brazil. *Geoderma*. **267**, 156, **2016**.

29. JIAO S., CHEN W., WANG J., DU N., LI Q., WEI G. Soil microbiomes with distinct assemblies through vertical soil profiles drive the cycling of multiple nutrients in reforested ecosystems. *Microbiome*. **6** (1), 146, **2018**.
30. SITU G., ZHAO Y., ZHANG L., YANG X., CHEN, LI S., WU Q., XU Q., CHEN J., QIN H. Linking the chemical nature of soil organic carbon and biological binding agent in aggregates to soil aggregate stability following biochar amendment in a rice paddy. *Science of the Total Environment*. **847**, 157460, **2022**.
31. ZHANG J., SAYER E.J., ZHOU J., LI Y., LI Y., LI Z., WANG F. Long-term fertilization modifies the mineralization of soil organic matter in response to added substrate. *Science of the Total Environment*. **798**, 149341, **2021**.
32. ZHOU J., QU T., LI Y., VAN ZWIETEN L., WANG H., CHEN J., SONG X., LIN Z., ZHANG X., LUO Y., CAI Y., ZHONG Z. Biochar-based fertilizer decreased while chemical fertilizer increased soil N₂O emissions in a subtropical Moso bamboo plantation. *Catena*. **202**, 105257, **2021**.
33. LASSALETTE L., EINARSSON R., QUEMADA M. Nitrogen use efficiency of tomorrow. *Nature Food*. **4** (4), 281, **2023**.
34. KICIŃSKA A., POMYKAŁA R., IZQUIERDO-DIAZ M. Changes in soil pH and mobility of heavy metals in contaminated soils. *European Journal of Soil Science*. **73** (1), e13203, **2022**.
35. ZHANG L., CHU Q., ZHOU J., RENGEL Z., FENG G. Soil phosphorus availability determines the preference for direct or mycorrhizal phosphorus uptake pathway in maize. *Geoderma*. **403**, 115261, **2021**.
36. LI P., WU M., KANG G., ZHU B., LI H., HU F., JIAO J. Soil quality response to organic amendments on dryland red soil in subtropical China. *Geoderma*. **373**, 114416, **2020**.
37. LIU H., DU X., LI Y., HAN X., LI B., ZHANG X., LI Q., LIANG W. Organic substitutions improve soil quality and maize yield through increasing soil microbial diversity. *Journal of Cleaner Production*. **347**, 131323, **2022**.
38. CHEN L., JIANG Y., WANG H., ZHAO Q., SUN B. Effects of long-term application of organic materials on phosphorus fractions and availability in red soil. *Soils*. **52** (3), 451, **2020**.
39. MAO Q., LU X., ZHOU K., CHEN H., ZHU X., MORI T., MO J. Effects of long-term nitrogen and phosphorus additions on soil acidification in an N-rich tropical forest. *Geoderma*. **285**, 57, **2017**.
40. PAN S., WANG Y., QIU Y., CHEN D., ZHANG L., YE C., GUO H., ZHU W., CHEN A., XU G., ZHANG Y., BAI Y., HU S. Nitrogen-induced acidification, not N-nutrient, dominates suppressive N effects on arbuscular mycorrhizal fungi. *Global Change Biology*. **26** (11), 6568, **2020**.
41. XING J., PENG L., CHEN J., HUANG J., JIANG P., QIN H. A new insight into spacing patterns of soil bacterial microbiome induced by root rot of *Carya cathayensis*. *Applied Soil Ecology*. **174**, 104416, **2022**.
42. YU X., KEITEL C., ZHANG Y., WANGECI A.N., DIJKSTRA F.A. Global meta-analysis of nitrogen fertilizer use efficiency in rice, wheat and maize. *Agriculture, Ecosystems & Environment*. **338**, 108089, **2022**.
43. LI Y., GONG H., LI S., ZHANG Y. Ecological Stoichiometry Homeostasis of Six Microelements in *Leymus chinensis* Growing in Soda Saline-Alkali Soil. *Sustainability*. **12** (10), 4226, **2020**.
44. AHMED N., ZHANG B., CHACHAR Z., LI J., XIAO G., WANG Q., HAYAT F., DENG L., NAREJO M.-U.-N., BOZDAR B., TU P. Micronutrients and their effects on Horticultural crop quality, productivity and sustainability. *Scientia Horticulturae*. **323**, 112512, **2024**.
45. PONI S., GATTI M., PALLIOTTI A., DAI Z., DUCHÊNE E., TRUONG T.-T., FERRARA G., MATARRESE A.M.S., GALLOTTA A., BELLINCONTRO A., MENCARELLI F., TOMBESI S. Grapevine quality: A multiple choice issue. *Scientia Horticulturae*. **234**, 445, **2018**.
46. LI P., WU M., KANG G., ZHU B., LI H., HU F., JIAO J. Soil quality response to organic amendments on dryland red soil in subtropical China. *Geoderma*. **373**, 114416, **2020**.
47. LI P., SHI K., WANG Y., KONG D., LIU T., JIAO J., LIU M., LI H., HU F. Soil quality assessment of wheat-maize cropping system with different productivities in China: Establishing a minimum data set. *Soil & Tillage Research*. **190**, 31, **2019**.
48. CHENG J., DING C., LI X., ZHANG T., WANG X. Soil quality evaluation for navel orange production systems in central subtropical China. *Soil & Tillage Research*. **155**, 225, **2016**.
49. CHEN Y.-D., WANG H.-Y., ZHOU J.-M., XING L., ZHU B.-S., ZHAO Y.-C., CHEN X.-Q. Minimum Data Set for Assessing Soil Quality in Farmland of Northeast China. *Pedosphere*. **23** (5), 564, **2013**.