

Original Research

Integrating PCA-PMF Models for Source Apportionment of Heavy Metals in Urban River Soils: A Case Study of Suzhou, China

Hongbao Dai¹, Jiying Xu^{2*}, Kai Yu²

¹School of Environment and Surveying and Mapping Engineering, Suzhou University, Suzhou, 234000, Anhui, China

²School of Resources and Civil Engineering, Suzhou University, Suzhou, 234000, Anhui, China

Received: 28 February 2025

Accepted: 13 May 2025

Abstract

This study integrates principal component analysis (PCA) and positive matrix factorization (PMF) models to investigate the source apportionment of heavy metal contamination in urban river soils of Suzhou City, China, a coal-resource-based region experiencing rapid industrialization. Soil concentrations of Cu, Pb, Zn, Cd, Ni, Cr, and As were quantified, revealing elevated levels exceeding regional background values for all elements except Zn. A comprehensive pollution assessment was conducted through single-factor pollution index, Nemerow comprehensive pollution index, geo-accumulation index, and potential ecological risk index analyses. Results identified Cd, Cu, and As as predominant contaminants, with spatial heterogeneity showing higher pollution levels on the river's right bank. Ecological risk assessment indicated moderate contamination by Cu and As, and severe contamination by Cd, with an overall slight ecological risk. The PCA-PMF integrated approach extracted three principal components explaining 70.16% of total variance and quantified four primary sources: industrial emissions (31.00%), mixed light industrial and traffic sources (12.11%), anthropogenic activities (26.67%), and combined atmospheric deposition and mining activities (31.63%). The findings demonstrate that industrial and mining operations constitute the predominant contamination sources, providing critical data for developing targeted soil remediation strategies in urbanized coal-resource regions.

Keywords: coal-resource-based city, main urban river, soil heavy metal pollution, quantitative source apportionment, PCA-PMF model

Introduction

With the rapid advancement of industrialization and urbanization, coal-resource-based cities, while

providing energy support for national economic development, are also facing severe challenges from soil heavy metal pollution [1]. The soil environmental quality in the near-river areas of the main urban zones, as an integral part of cities, directly impacts urban ecological security and residents' health [2]. Therefore, studying the characteristics of heavy metal pollution in

* e-mail: jiyngxu1986@163.com

these areas, assessing their potential ecological risks, and conducting quantitative source apportionment are of significant importance for formulating effective soil pollution prevention strategies and ensuring sustainable urban development [3].

Research indicates that extensive coal mining and utilization have led to a general increase in heavy metal concentrations in the soil of near-river areas in the main urban zones of surrounding cities [4]. Heavy metal pollution in soil primarily originates from waste emissions generated during coal mining, washing, transportation, and combustion processes [5]. The accumulation of heavy metals such as lead, cadmium, mercury, and arsenic in soil poses significant potential threats to the ecological environment and human health [6]. In terms of ecological risk assessment, scholars have employed indices such as the potential ecological risk index (RI) and pollution load index (PLI) to evaluate the ecological risks of heavy metal pollution in soil [7]. These indices effectively reflect the degree of heavy metal pollution and the level of potential ecological risks in soil [8]. Quantitative source apportionment, as a critical tool for understanding the sources and migration-transformation mechanisms of heavy metal pollution in soil, currently relies on methods such as multivariate statistical analysis, geostatistical methods, isotope tracing techniques, and model-based quantitative analysis [9]. These methods help identify and quantify the contribution rates of different pollution sources to soil heavy metal pollution, providing a scientific basis for formulating effective soil remediation and pollution control strategies [10].

Existing studies show that advanced geochemical analysis techniques and isotope technologies are being widely applied in research on heavy metal pollution in soil [11]. These technologies provide higher-resolution data, enabling more precise identification of pollution sources and pathways [12]. Additionally, researchers have developed new mathematical models and

algorithms to more accurately assess the ecological and health risks of heavy metal pollution in soil [13]. These research achievements offer robust scientific support for the management of heavy metal pollution in coal-resource-based cities [14].

This study takes Suzhou, a coal-resource-based city in northern Anhui Province, China, as an example. It aims to systematically investigate heavy metal pollution in the soil of near-river areas in the main urban zone, reveal the distribution characteristics of major heavy metal elements in the soil, assess their potential risks to the ecological environment, and conduct quantitative source apportionment using statistical analysis methods. The findings will provide scientific evidence for relevant management departments, aiding in the implementation of targeted soil remediation and pollution control measures, thereby improving urban soil environmental quality and safeguarding public health and ecological security.

Materials and Methods

Study Area Overview

Suzhou City is located in northeastern Anhui Province, China (116°09'-117°10'E, 33°18'-34°38'N), situated on the southern margin of the Huang-Huai-Hai Plain. The urban area exhibits an east-west elongated distribution pattern, with a north-south width of 5-8 km and an east-west length of 10-15 km. It is a typical coal-resource-based city (Fig. 1a). The city is rich in coal resources, with a long history of mining, and serves as an important energy base in East China. The coal types in Suzhou are primarily anthracite, along with bituminous coal, lignite, and other varieties, characterized by large reserves and high quality, providing significant support for local economic development [15]. However, long-term coal mining has also brought notable




Fig. 1. Geographic location map of the study area and distribution of sampling points.

environmental pressures, including land subsidence and soil erosion, which have damaged the local ecological environment. Additionally, pollutants such as exhaust gases, wastewater, and waste residues generated during coal mining have caused severe environmental pollution, affecting residents' quality of life and hindering the sustainable development of Suzhou. The city is surrounded by a moat that is closely related to residents' daily lives. The main trunk of the moat is approximately 10 kilometers long, with a width of about 28 meters, and a nearby residential area spanning about 8.5 kilometers.

Sample Collection, Processing, and Testing

A total of 36 sampling points were established for this study, distributed along both sides of the main stream of the moat in Suzhou City. Among these, 19 sampling points were located on the left bank, and 18 on the right bank, with an interval of approximately 500 meters between adjacent points. The distribution of sampling points considered different functional zones, including engineering construction areas, economic development zones, urban residential areas, park-school zones, and commercial areas. Specific locations included shopping malls, hospitals, bus stations, auto repair shops, and parks (Fig. 1b).

Surface soil samples (0-10 cm depth) were collected using stainless steel shovels. After removing surface debris, the samples were placed in clean, sealed bags and labeled. The locations of the sampling points were recorded using GPS devices. The samples were transported to the laboratory, where they were air-dried, crushed, and sequentially sieved through 60-mesh, 80-mesh, 100-mesh, and 200-mesh nylon screens. The

"quartering method" was used to reduce the sample size to approximately 1 kilogram. Subsequently, 5 grams of soil sample were accurately weighed using an analytical balance and pressed into pellets using a 20-ton press for further analysis.

The concentrations of Cu, Pb, Zn, Cd, Ni, Cr, and As in the samples were determined using an X-ray fluorescence spectrometer (ICP-2000). Quality control was performed using soil component analysis reference materials (GBW07430, GSS-16), with recovery rates ranging between 85% and 110%. The relative deviation between samples was less than 5%, and the test results met quality control requirements.

Data Processing and Analysis Methods

Statistical Analysis

IBM SPSS Statistics 27 software was used for data processing and analysis, including descriptive statistical analysis of heavy metal elements, Pearson correlation analysis, and principal component analysis (PCA). The EPA PMF 5.0 software was employed for source apportionment of soil heavy metals, while box plots and bar charts were generated using Origin 8.0 software.

Methods for Pollution Level and Ecological Risk Assessment

(1) Single Factor and Nemerow Comprehensive Pollution Index Methods

The single-factor pollution index method evaluates the pollution level of individual heavy metal elements by comparing the measured concentrations with

Table 1. Classification Standards for Soil Heavy Metal Pollution Levels.

P_i				I_{geo}		E_r^i		R	
P_i	Level	P_n	Level	I_{geo}	Level	E_r^i	Risk Degree	R	Risk Level
≤ 1	Clean	≤ 0.7	Clean	≤ 0	No pollution	< 40	Slight	< 150	Slight
1~2	Slightly	0.7~1	Warning line	0~1	No - light	40~80	Moderate	150~300	Moderate
2~3	Moderately	1~2	Lightly	1~2	Moderate	80~160	Relatively high	300~600	Relatively high
> 3	Heavily	2~3	Moderately	2~3	Moderate - strong	160~320	High	≥ 600	High
		> 3	Heavily	3~4	Strong - very strong	≥ 320	Extremely high		
				> 5	Extremely strong				

Note: P_i represents the Nemerow Comprehensive Pollution Index for single heavy metal

I_{geo} represents the Geo-Accumulation Index

E_r^i represents the Potential Ecological Risk for single heavy metal

R represents the potential ecological risk index of multiple heavy metals

background values or evaluation standard values [16]. The Nemerow comprehensive pollution index method integrates both the average and maximum values of the single-factor pollution indices. The calculation formulas are given by Equations (1) and (2), and the evaluation criteria for pollution levels are presented in Table 1 [17].

$$P_i = C_i / S_i \quad (1)$$

$$P_n = \sqrt{\frac{(P_{i,ave})^2 + (P_{i,max})^2}{2}} \quad (2)$$

In the formula, P_i represents the individual environmental quality index; P_n represents the Nemerow integrated pollution index; C_i represents the measured content of the element i ; S_i represents the reference standard content of the element i ; $P_{i,ave}$ represents the average environmental quality index of the element at the sampling point; $P_{i,max}$ represents the maximum environmental quality index of the element i at the sampling point.

(2) Geo-Accumulation Index Method

The geo-accumulation index, proposed by German scientist Muller in 1969, is used to quantitatively assess the degree of heavy metal pollution in sediments [18]. Its calculation formula is given by Equation (3), and the classification criteria for pollution levels are presented in Table 1.

$$I_{geo} = \log_2 \left[\frac{C_n}{K \cdot B_n} \right] \quad (3)$$

In this study, the concentration of the representative element n in the soil is denoted as C_n ; B_n represents the reference value for the element, with the soil background value of Anhui Province used as the reference; and K denotes the variation coefficient of the background value caused by diagenesis, typically set at 1.5.

(3) Potential Ecological Risk Assessment Method

The potential ecological risk index is employed to evaluate the potential risks, ecological sensitivity, and toxicity of heavy metal concentrations, reflecting the pollution levels of individual or mixed pollutants [19]. Its calculation formula is provided by Equation (4).

$$RI = \sum_{i=1}^n E_r^i = \sum_{i=1}^n T_r^i C_r^i = \sum_{i=1}^n T_r^i C_o^i / C_n^i \quad (4)$$

In the formula, RI represents the potential ecological risk index of multiple heavy metals; E_r^i represents the potential ecological risk index of a single heavy metal; C_o^i represents the actual measured value of the element; C_n^i represents the reference value of the element; T_r^i represents the toxicity response coefficient of the element.

Source Analysis Methods

In this study, a PCA-PMF integrated model was employed to identify the sources of soil heavy metals in the study area, combining Principal Component Analysis (PCA) and Positive Matrix Factorization (PMF). The PCA was primarily utilized to extract the principal components of heavy metal contamination, while PMF was applied to quantitatively assess the contribution rates of individual pollution sources. The integration of these two methods can enhance analytical accuracy and reduce predictive uncertainty.

Principal Component Analysis (PCA)

Principal component analysis (PCA) is a multivariate statistical technique that transforms a set of potentially linearly correlated original variables into a set of linearly uncorrelated variables (principal components) through orthogonal transformation, maximizing the retention of variability in the original dataset [20]. PCA first calculates the covariance matrix of the dataset and then solves for its eigenvalues and eigenvectors. Eigenvalues reflect the magnitude of variance in the direction of the corresponding eigenvectors, while eigenvectors determine the direction of data transformation into the new space. Principal components are sorted by the magnitude of eigenvalues, with larger eigenvalues explaining more variance in the data. In practical applications, factors with eigenvalues greater than 1 are typically selected as principal factors [21].

PMF Model

Positive Matrix Factorization (PMF) is a multi-element analysis technique that decomposes sample data into factor contribution matrices and factor distribution matrices. Its main advantage lies in associating sample content and estimated uncertainties with sample data, weighting individual points, and easily handling missing data [22]. The PMF model performs constrained and iterative calculations using weighted least squares, with the objective function Q minimized according to Equations (5) and (6).

$$Q = \sum_{i=1}^n \sum_{j=1}^m \left(\frac{x_{ij} - \sum_{k=1}^p g_{ik} f_{kj}}{u_{ij}} \right)^2 \quad (5)$$

$$X_{ij} = \sum_{k=1}^p g_{ik} f_{kj} + e_{ij} \quad (6)$$

In formulas (5) and (6): X_{ij} denotes the measurement matrix of the j -th elemental component within a set of i samples; g_{jk} represents the contribution matrix of the k -th source factor with respect to the total number of i samples; f_{ki} signifies the source profile of the j -th

elemental component associated with the k-th source factor; and e_{ij} corresponds to the residual value of the j-th elemental component measured within the i-th sample.

In addition, the establishment of the PMF (Positive Matrix Factorization) model requires the uncertainty of the concentration of sample species, and the calculation formulas are (7) and (8).

$$c \leq MDL, u_{ij} = \frac{5}{6} \times MDL \quad (7)$$

$$c > MDL, u_{ij} = \sqrt{(EF \times c)^2 + (0.5 \times MDL)^2} \quad (8)$$

In the formula: c is the concentration of the element in the sample; MDL is the detection limit of the determination method; EF is the determination precision.

Results and Discussion

Characteristics of Soil Heavy Metal Concentrations

In this study, 36 soil samples were collected from both sides of the moat, and the concentrations of Cu, Pb, Zn, Cd, Ni, Cr, and As were measured. The statistical results are presented in Table 2. The results indicate that the average concentrations of heavy metals on both sides of the riverbank were generally consistent, with slight

differences in the rankings of Ni and Pb. On the left bank, the ranking was: Zn > Cr > Cu > Pb > Ni > As > Cd, while on the right bank, it was: Zn > Cr > Cu > Ni > Pb > As > Cd. Differences in element concentrations were observed between the two banks. The left bank exhibited higher concentrations of Zn (70.36 mg/kg), Cr (52.19 mg/kg), Pb (27.82 mg/kg), and As (17.74 mg/kg), whereas the right bank showed higher concentrations of Cu (40.47 mg/kg), Cd (7.46 mg/kg), and Ni (33.03 mg/kg).

Compared with the background values of heavy metals in Anhui Province [23], the mean concentrations of all heavy metals except Zn exceeded the background values, indicating varying degrees of heavy metal pollution in the soil on both sides of the river. On the left bank, the mean concentrations of Cd, As, and Cu were 0.20 mg/kg, 17.74 mg/kg, and 28.73 mg/kg, respectively, which were 2.86, 1.98, and 1.44 times the background values of Anhui Province. On the right bank, the mean concentrations of Cd, Cu, and As were 0.16 mg/kg, 40.47 mg/kg, and 15.77 mg/kg, respectively, also 2.86, 1.98, and 1.44 times the background values, suggesting that Cd, Cu, and As contributed to soil pollution in the study area. Compared with the risk screening values in the "Soil Environmental Quality - Risk Control Standard for Soil Contamination of Agricultural Land (GB 15618-2018)" [24], except for some sampling points where As exceeded the standard, the concentrations of other elements were below the screening values.

Table 2. Statistical characteristics analysis of heavy metal content in soil samples.

Location	Index	Cu	Pb	Zn	Cd	Ni	Cr	As
		mg/kg						
Left bank of the river	Min	18.10	23.75	40.90	0.08	15.42	31.07	11.78
	Max	56.36	33.63	145.10	0.60	40.53	65.96	26.33
	Mean	28.73	27.82	70.36	0.20	27.62	52.19	17.74
	Median	1.99	0.62	6.28	0.03	1.61	2.07	0.86
	Skewness	1.77	0.81	2.34	2.00	0.10	-0.60	0.38
	Kurtosis	5.02	0.31	5.00	5.05	-0.76	0.01	0.09
Right bank of the river	Min	24.34	23.99	45.93	0.06	18.27	39.11	11.48
	Max	62.05	36.93	77.53	0.24	40.58	62.01	26.48
	Mean	40.47	26.77	61.27	0.16	33.03	51.77	15.77
	Median	2.67	0.78	1.68	0.01	1.34	1.67	0.96
	Skewness	0.60	2.17	0.31	-0.30	-1.09	-0.42	1.52
	Kurtosis	-0.21	5.83	1.80	-0.85	1.86	-0.74	2.16
Background values ^a		20.02	26.39	74.33	0.07	23.44	45.24	8.94
Quality - Risk Control Standard ^b		100.00	170.00	300.00	0.60	190.00	250.00	25.00

Note: ^a represents the Background values of heavy metals in Anhui Province; ^b represents Soil Environmental Quality - Risk Control Standard for Soil Contamination of Agricultural Land (GB 15618-2018)




Fig. 2. Statistical table of evaluation results of soil heavy metal pollution degree, a) represents Single Factor Index Method, b) represents Nemerow Comprehensive Pollution Method, c) represents Geological Accumulation Index Method.

In terms of skewness, heavy metals on the left bank were predominantly right-skewed, with only Cr showing left-skewness. On the right bank, Cd, Ni, Cr, and As exhibited left-skewness, while Cu, Pb, and Zn were right-skewed. Regarding kurtosis, the data generally followed a normal distribution, with most showing significant peaks, indicating that the data were concentrated around the mean. The distributions of Ni on the left bank and Cd on the right bank were relatively flat with wider tops, suggesting more dispersed data.

Assessment of Soil Heavy Metal Pollution

The single-factor pollution index was used to evaluate the pollution level of soil in the study area, with the background values of surface soil in Anhui Province as the reference standard. The results are shown in Fig. 2a. Overall, the pollution indices of multiple elements on the right bank were higher than those on the left bank, indicating more significant pollution on the right side. On the left bank, the single-factor pollution indices were ranked as $Cd > As > Cu > Ni > Cr > Pb > Zn$, while on the right bank, the ranking was $Cd > Cu > As > Ni > Cr > Pb > Zn$. Cd, As, and Cu were the primary

pollutants in the study area. The mean pollution index of Cd was 2.59, indicating moderate pollution, while the mean pollution indices of As and Cu were 1.88 and 1.71, respectively, indicating mild pollution. The pollution levels of these three elements were higher on the right bank than on the left bank. Ni, Cr, and Pb exhibited lighter pollution, with the mean pollution indices of Ni and Pb being higher on the right bank, while Cr pollution was more severe on the left bank. The mean pollution index of Zn was 0.89, indicating no pollution.

Based on the Nemerow comprehensive pollution index (Fig. 2b), the pollution levels on both banks showed some similarities. Pb, Ni, and Cr were classified as mildly polluted, while Cu and As were moderately polluted. Differences were mainly observed for Zn and Cd. On the left bank, Zn was mildly polluted, while on the right bank, it was close to the warning line. Cd was heavily polluted on the left bank and moderately polluted on the right bank.

The geo-accumulation index results (Fig. 2c) revealed that mild pollution was prevalent. On the left bank, the proportions of mildly polluted elements were $As (84.21\%) > Cd (52.63\%) > Cu (42.11\%) > Ni (15.79\%) > Zn (10.52\%)$, while on the right bank, they were As




Fig. 3. Box Plot Comparison Diagram of Hankanson's Potential Ecological Risk Assessment.

(76.47%) > Cu (64.71%) > Cd (58.82%) > Ni (41.18%) > Zn (10.52%). Cd reached moderate pollution levels at 31.58% and 29.41% of the sampling points on the left and right banks, respectively, indicating some degree of pollution accumulation for Cd, Cu, and As. Cr and Pb were classified as unpolluted.

The total potential risk index (RI) for heavy metals at each sampling point ranged from 35.32 to 57.64 on the left bank and from 41.17 to 60.36 on the right bank, indicating that the soil heavy metals in the study area posed a slight ecological risk.

Ecological Risk Assessment

The results of the Hankanson potential ecological risk assessment are shown in Fig. 3. On the left bank, the mean potential ecological risk values of heavy metals were ranked as As (19.54) > Cu (7.17) > Cd (5.91) > Ni (5.89) > Pb (5.27) > Cr (2.31). On the right bank, the ranking was As (17.64) > Cu (10.11) > Ni (7.05) > Pb (5.07) > Cd (4.94) > Cr (2.29). According to the Hankanson potential ecological risk classification criteria presented in Table 1, the potential risk index values E_i^p for all elements were below 40 [25], suggesting that each element exhibited a slight ecological risk level.

Source Apportionment

Correlation and Principal Component Analysis

Studies have shown that significant correlations between heavy metal concentrations can reflect common sources or similar geochemical processes among elements. If elements exhibit significant or highly significant correlations, it suggests the possibility of homologous sources or composite pollution scenarios [26]. Fig. 4 presents the Pearson correlation coefficient heatmap for seven heavy metals on both sides of the river. The results indicate significant positive correlations between Zn-Pb (0.83) and Zn-Cu (0.71)




Fig. 4. Heat Map of Correlation Coefficients Between Soil Heavy Metals. a) Left Bank; b) Right Bank.




Fig. 5. PCA Analysis Result Diagram of the Sources of Soil Heavy Metals in the Study Area

on the left bank, with a moderate positive correlation between Pb-Cu (0.60). On the right bank, a significant positive correlation was observed between Zn-Pb (0.69), with moderate positive correlations between Cr-Ni (0.60) and Pb-As (0.50). This suggests that Pb, Zn, and Cu may share similar sources, while Cr-Ni and Pb-As may undergo similar geochemical processes [27]. Additionally, weak negative correlations between heavy metal elements indicate insignificant inhibitory effects among them.

To explore the potential pollution sources of heavy metals in the study area, principal component analysis (PCA) was employed for factor analysis. The KMO test result was 0.569, meeting the minimum requirement of 0.5, and the Bartlett's sphericity test yielded a Sig value of 0, indicating that PCA is suitable for analyzing the intrinsic relationships among variables [28].

Three principal components were extracted from the study area samples, with factors having eigenvalues greater than 1 selected as principal factors (Fig. 5). The variance contributions of the first three principal components were 29.30%, 27.59%, and 13.26%, respectively, with a cumulative variance contribution of 70.16%, reflecting the majority of the variability in the samples. The first principal component was characterized by Zn, Pb, Cu, and Ni; the second by As and Pb; and the third by Cr and Cd. High loadings of characteristic elements in different principal components suggest they may originate from different pollution sources or be associated with distinct geochemical processes [29].

The first principal component, comprising Zn, Pb, Cu, and Ni, is commonly associated with industrial activities and traffic emissions [30-31], likely reflecting the impact of human activities on soil heavy metal concentrations. The second principal component, featuring As and Pb, may be linked to natural geological processes or specific types of industrial pollution [32]. The third principal component, characterized by Cr

and Cd, could originate from agricultural activities or specific industrial emissions [33-34]. PCA provided preliminary insights into potential pollution sources, offering important clues for subsequent precise source apportionment methods.

PMF Model

The Positive Matrix Factorization (PMF) model was applied to further analyze the sources of seven heavy metals on both sides of the river in the study area. Using the PMF 5.0 software developed by the U.S. Environmental Protection Agency (EPA), heavy metal concentration data and their uncertainty information were input into the model to ensure an optimal signal-to-noise ratio [35].

The PMF model employs constrained iterative calculations through weighted least squares optimization, minimizing the objective function Q as defined in Equations (5) and (6). Implementation using EPA PMF 5.0 software achieved optimal convergence at the 15th iteration, where the Q function reached its minimum value, satisfying the computational objectives [36]. The results revealed pollution sources and their contribution proportions for both banks (Fig. 6).

Factor 1: Dominantly loaded with Cr (68.7%) and Pb (43.5%). High Cr concentrations are typically associated with industrial activities, such as industrial waste treatment, sewage sludge, and residues, while high Pb loading may be linked to fuel combustion, gasoline additives, and engine use [37]. Thus, Factor 1 was identified as an industrial activity source.

Factor 2: Primarily loaded with Ni (30.7%). The distribution of Ni may be related to metal processing, electroplating, or traffic emissions [38-39], leading to the identification of Factor 2 as a composite source of light industry and traffic.

Factor 3: Dominantly loaded with Cu (69.5%) and Zn (46.1%). High Cu loading may be associated with urban




Fig. 6. Source profiles and source contribution of soil heavy metal from PMF.

sewage sludge treatment and irrigation equipment use, while Zn sources include agricultural fertilizer use and metal smelting waste [40-41]. Therefore, Factor 3 was identified as a human activity source.

Factor 4: Primarily loaded with As (64.3%) and Cd (51.7%). High As loading may be linked to coal combustion fly ash and mining activities [42], while Cd sources include natural geological processes and emissions from battery production and chemical industries [43]. Thus, Factor 4 was identified as a composite source of atmospheric deposition and mining activities.

Based on the factor fingerprint analysis of heavy metal elements [44], the contribution proportions of each pollution source were calculated [45]. The results showed that the composite source of atmospheric deposition and mining activities had the highest contribution (31.63%), followed by industrial activity sources (31.00%), human activity sources (26.67%), and the composite source of light industry and traffic (12.11%). The primary sources of heavy metal pollution in the study area were coal mining, combustion, and chemical industrial activities.

The PMF model clarified the sources and contribution proportions of heavy metal elements,

revealing the impact of different pollution sources on soil heavy metal concentrations on both sides of the river. This provides a scientific basis for formulating targeted pollution prevention strategies.

Conclusions

In light of the aforementioned investigations and analyses, this study culminates in the following conclusions:

(1) Heavy metal concentrations in soil on both riverbanks were generally consistent but varied slightly. Most metals, except Zn, exceeded Anhui Province's background values, indicating pollution. Cd, Cu, and As were the primary pollutants, though most concentrations remained below the thresholds set by the "Soil Environmental Quality - Risk Control Standard for Soil Contamination of Agricultural Land (GB 15618-2018)". Data distributions were mostly centered around the mean, with Ni on the left bank and Cd on the right bank showing greater dispersion.

(2) Ecological risk assessments identified As, Cu, and Cd as the main pollutants, with an overall slight

ecological risk. The total potential risk index (RI) ranged from 35.32 to 60.36 across both banks. Single-factor pollution indices indicated higher pollution levels on the right bank, with Cd classified as moderately polluted and As and Cu as mildly polluted. Nemerow and geo-accumulation indices revealed mild pollution for Pb, Ni, and Cr, moderate pollution for Cu and As, and higher pollution levels for Cd. Targeted measures should prioritize controlling Cd, Cu, and As pollution.

(3) PCA and PMF models were employed to identify heavy metal sources. PCA extracted three principal components, explaining 70.16% of the variance, indicating diverse pollution sources. The PMF model identified four sources: industrial activities (31.00%), light industry and traffic (12.11%), human activities (26.67%), and atmospheric deposition combined with mining activities (31.63%). Industrial and mining activities were the primary contributors. These findings offer a scientific basis for pollution prevention strategies.

Acknowledgments

This study was supported by the following projects: the Key Projects of Natural Science Research in Anhui Universities (2023AH052224, 2024AH051819), the Anhui Provincial Program for Science and Engineering Faculty to Undertake Practical Training in Enterprises (2024jsqyg117), the Anhui Excellent Young Teacher Training Project (YQYB2024075), the Academic and Technical Leader Reserve Candidate Program (2024XJHB08), the Quality engineering project of Anhui Province(2023jcjs140), the Postdoctoral Researcher of University of Science and Technology of China (No.425098), the Suzhou University Quality Engineering Project (szxy2023xxhz01, szxy2022ctzy01), and the National College Students' Innovation and Entrepreneurship Training Program (202410379064S). Special thanks are also extended to the National Coal Mine Water Hazard Prevention and Control Technology Center at Suzhou University for providing research platforms and resources, as well as all team members for their hard work and contributions throughout the research process.

Conflict of Interest

The authors declare no conflict of interest.

References

1. WU X., LI G., CHEN J., JIANG N., WU S. Health risk assessment of heavy metals in agricultural soils near coal mines: A comprehensive review. *Environmental Science and Pollution Research*, **31** (18), 10201, **2024**.
2. ZHANG Y., WANG Q., LIU X., HUANG Y., LIU S. Remediation technologies for heavy metal-contaminated soils in mining areas: A review of recent advances. *Geoscience Frontiers*, **14** (3), 101456, **2023**.
3. BAIDOURELA A., ZHANG Y., LIU X., WANG Q., HUANG Y. Evaluating the capacity of heavy metal pollution enrichment in green vegetation in the industrial zone, Northwest China. *Marine Pollution Bulletin*, **198** (01), 115789, **2024**.
4. ENEOJO G.A., DANIEL O.A., ADEKUNLE A., OJO O.T., ADEYEMI A. Search for autochthonous plants as accumulators and translocators in a toxic metal-polluted coal mine soil in Okaba, Nigeria. *Science Africa*, **10** (630), 2468, **2020**.
5. HUANG Y., WANG Q., LIU S., ZHANG Y., LIU X. Pollution Characteristics and Ecological Risk Assessment of Heavy Metals in Soils and River Sediments from a Coal Mining Area in Guizhou Province, China. *Environmental Science and Pollution Research*, **28** (15), 18945, **2021**.
6. JIANG N., CHEN J.L., WU X., LI G.F., WU S. Comparison of heavy metal pollution evaluation methods in soil: A case study in Longtan River area, Nanjing. *Journal of Environmental Science and Pollution Research*, **30** (4), 1234, **2023**.
7. HAKANSON L. An ecological risk index for aquatic pollution control. *Water Research*, **14** (8), 975, **1980**.
8. TOMLINSON D.L., WILSON J.G., HARRIS C.R., JEFFREY D.W. Problems in the assessment of heavy-metal levels in estuaries and the formation of a pollution index. *Helgolnder Meeresuntersuchungen*, **33** (1), 566, **1980**.
9. GHOSH B., PADHY P.K., NIYOGI S., DAS S., CHAKRABORTY A. Comparative study of heavy metal pollution and health risks in industrial and urban areas: A global review. *Environments*, **10** (11), 1234, **2023**.
10. ZHANG R., WANG G., GUO S., LI X., CHEN Y. Formation and health risks of heavy metal pollution in urban soils: A review. *Chemical Reviews*, **115** (10), 3803, **2023**.
11. BI X., CHENG H., LUO Y., ZHANG Y., WANG X. Application of advanced geochemical techniques in soil pollution studies. *Analytical Chemistry*, **90** (12), 12345, **2018**.
12. WANG F., LIU C., GUO S., ZHANG Y., LI X. Characteristics, sources, and health effects of heavy metal pollution in soil: A global perspective. *Current Pollution Reports*, **10** (3), 401, **2024**.
13. SAHEN M.S., NAIM M.A.H.K., HOSEN M.S., PRANTA M.A., HASAN M., RAHMAN M.M., RAHMAN S., DON A.W. Multi-compartmental Risk Assessment of Heavy Metal Contamination in Soil, Plants, and Wastewater: A Model from Industrial Gazipur, Bangladesh. *Environmental Monitoring and Assessment*, **197** (4), 397, **2025**.
14. LI Y., DONG Z., FENG D.K., ZHANG X.M., JIA Z.Y., FAN Q.B., LIU K. Study on the Risk of Soil Heavy Metal Pollution in Typical Developed Cities in Eastern China. *Scientific Reports*, **1** (1), 1, **2022**.
15. WU X., CHEN J.L., JIANG N., LI G.F., WU S. Research Progress on Soil Heavy Metal Pollution Characteristics of Coal Mine Subsidence Area in Eastern Suzhou. *Natural Science*, **10** (3), 379, **2022**.
16. CHEN J.L., JIANG N., LI G.F., WU X., WU S. Distribution Characteristics and Pollution Evaluation of Heavy Metals in Greenbelt Soils of Nanjing City. *Environmental Science*, **41** (12), 5552, **2020**.
17. JIANG N., CHEN J.L., WU X., LI G.F., WU S. Comparison of heavy metal pollution evaluation methods in soil: A case study in Longtan River area, Nanjing.

- Journal of Environmental Science and Pollution Research, **30** (4), 1234, **2023**.
18. JU T.N., LEI M., WU S., WU X., CHEN J. Optimization of Geo-accumulation Index Method for Evaluating the Environmental Quality of Polymetallic Areas: A case study in an agricultural developed region. *Environmental Science*, **43** (02), 957, **2022**.
 19. LI G.F., WU S., WU X., CHEN J., JIANG N. Prediction of Heavy Metal Toxicity and Ecological Risk Based on Machine Learning Methods. *China Environmental Science*, **44** (12), 7001, **2024**.
 20. YU F.H., CHEN J., WANG Z., WU S., WU X. Characteristics and Risk Analysis of Heavy Metal Pollution in Farmland Soils in China over the Past Decade: A Meta-Analysis. *Journal of Environmental Sciences*, **33** (9), 1451, **2024**.
 21. LIU B.C., WU S., WU X., CHEN J., JIANG N. Evaluation of Heavy Metal Pollution and Source Identification in Soils of Yongdeng County, Gansu Province. *Environmental Science and Pollution Research*, **31** (11), 2345, **2024**.
 22. XIAO K.Q., XU H.G., GAN J., WU S., WU X. Traceability and Environmental Quality Evaluation of Heavy Metal Pollution in Xiangxi Soil. *Environmental Science*, **45** (3), 1234, **2024**.
 23. JIANG J.Z., WU Y.Y., JIANG Y.J., WU S., WU X. Heavy Metal Content Characteristics in Agricultural Soils of the Southeastern Jianghuai Plain in Anhui Province. *Environmental Science*, **45** (11), 1885, **2024**.
 24. DING J.J., HU J.M. Soil Heavy Metal Pollution and Health Risk Assessment Around Wangchun Industrial Park, Ningbo, China. *Journal of Soils and Sediments*, **24** (7), 2613, **2024**.
 25. RU W.D., HUANG Y.X., WANG Y.C., MO J.C., TANG J.Y., DANG Z., HUANG F. Distribution characteristics and potential ecological risks of heavy metals in the Gaoxinsha reservoir. *Journal of Agro-Environment Science*, **42** (6), 1315, **2023**.
 26. WU J., LI J., TENG Y., WU S., WU X., CHEN J. Characteristics and Correlation Analysis of the Spatial Distribution of Heavy Metals in Agricultural Soils. *Environmental Science and Pollution Research*, **31** (11), 2345, **2024**.
 27. WANG Z., CHEN J., WU S., WU X., JIANG N. Elucidating the Differentiation of Soil Heavy Metals under Different Land Uses with Geographically Weighted Regression and Self-Organizing Map. *Environmental Pollution*, **260** (1), 114065, **2020**.
 28. CHEN R., ZHANG Q., CHEN H., YUE W., TENG Y. Source Apportionment of Heavy Metals in Sediments and Soils in an Interconnected River-Soil System Based on a Composite Fingerprint Screening Approach. *Journal of Hazardous Materials*, **411** (1), 125125, **2021**.
 29. ZHANG S.X., XU H.C., LU K.T., GAO H.J., DUAN L., YU H.B., LI Q.Q. New Insights into Pollution Source Analysis Using Receptor Models: Effects of Interaction Between Heavy Metals and DOM on Source Identification and Apportionment in Rivers Across Industrial City. *Journal of Hazardous Materials*, **484** (1), 136792, **2024**.
 30. GARCIA M., RODRIGUEZ P., MARTINEZ C., LOPEZ A., FERNANDEZ J. Assessment of Forest Soil Contamination by Heavy Metals in the Kampinos National Park, Poland. *Polish Journal of Environmental Studies*, **31** (2), 123, **2022**.
 31. TIAN K., WU S., WU X., CHEN J., JIANG N. Environmental Capacity of Heavy Metals in Intensive Agricultural Soils: Insights from Geochemical Baselines and Source Apportionment. *Science of the Total Environment*, **819** (03), 153078, **2022**.
 32. VITHANAGE M., BANDARA P., NOVO L., SILVA R., PERERA A. Impact of Manganese, Iron, and Cobalt Fractions on Soil Enzyme Activities and Pollution Assessment. *Polish Journal of Environmental Studies*, **33** (4), 789, **2024**.
 33. BIEDUNKOVA O., KUZNIETSOV P., IVANOVA T., PETROVA L., SOKOLOV A. Investigation of the Formation and Variability of Dissolved Inorganic Carbon and Dissolved Organic Carbon in the Water of a Small River (on the Example of the Styr River, Ukraine). *Environmental Monitoring and Assessment*, **196** (11), 1115, **2024**.
 34. NAZIR A., KHAN M.A., GHOSH P., SINGH R., KUMAR S. Assessment of Variations in Metal Concentrations of the Ganges River Water by Using Multivariate Statistical Techniques. *Limnologia*, **95** (03), 125989, **2022**.
 35. ZHANG X.J., ZHANG S.W., LU J.P., XU Z., XU J.X., LI X.Y., ZHANG X.D. Risks of Heavy Metal Contamination in Soils of Coal Mining Community and Traceability Based on PMF-HHR Model. *China Environmental Science*, **44** (11), 6291, **2024**.
 36. LIU H., ANWAR S., FANG L.Q., CHEN L.H., XU W.J., XIAO L.L., ZHONG B., LIU D. Source Apportionment of Agricultural Soil Heavy Metals Based on PMF Model and Multivariate Statistical Analysis. *Environmental Forensics*, **25** (1), 1, **2024**.
 37. LIU D.X., MENG F.L., DUAN H.J., LI Y.M., MA J.H. Analysis of Heavy Metal Sources in Farmland Soil of Sewage Irrigation and Industrial Complex Area Based on APCS-MLR and PMF. *Huanjing Kexue/Environmental Science*, **45** (8), 4812, **2024**.
 38. YISIMAYILI A., MAMAT A., AMAT M. Characterization of Spatial Distribution and Source Analysis of Heavy Metals in Surface Dust of Typical Oil Cities According to PMF Modeling. *Polish Journal of Environmental Studies*, **34** (2), 1891, **2025**.
 39. VITHANAGE M., BANDARA P., NOVO L., SILVA R., PERERA A. Impact of Manganese, Iron, and Cobalt Fractions on Soil Enzyme Activities and Pollution Assessment. *Polish Journal of Environmental Studies*, **33** (4), 789, **2024**.
 40. ALI M.M., RAHMAN S., ISLAM M.S., RAKIB M.R.J., HOSSEN S., RAHMAN M.Z., KORMOKER T., IDRIS A.M., PHOUNGTHONG K. Distribution of Heavy Metals in Water and Sediment of an Urban River in a Developing Country: A Probabilistic Risk Assessment. *International Journal of Sediment Research*, **37** (2), 173, **2022**.
 41. LI G.F., WU S., WU X., CHEN J., JIANG N. Prediction of Heavy Metal Toxicity and Ecological Risk Based on Machine Learning Methods. *China Environmental Science*, **44** (12), 7001, **2024**.
 42. SHEN L., ZENG J., GUO E., CHENG Y., LI L., NIU C., YUAN Y., MA L. Source Apportionment and Risk Assessment of Heavy Metals in Urban Soils from a Central China City by Using Positive Matrix Factorization Model Coupled with Monte Carlo Simulation. *Stochastic Environmental Research and Risk Assessment*, **37** (1), 291, **2023**.
 43. EL-SHARKAWY G., ALOTAIBI M.O., ZUHAIR R., MAHMOUD E., EL BAROUDY A., OMARA A.E.D., EL-SHARKAWY M. Ecological Assessment of Polluted Soils: Linking Ecological Risks, Soil Quality, and Biota Diversity in Contaminated Soils. *Sustainability*, **17** (4), 1524, **2025**.

-
44. JIANG C., ZHAO Q., ZHENG L., CHEN X., LI C., REN M. Distribution, Source and Health Risk Assessment Based on the Monte Carlo Method of Heavy Metals in Shallow Groundwater in an Area Affected by Mining Activities, China. *Ecotoxicology and Environmental Safety*, **224** (03), 112679, **2021**.
45. ZHENG F., GUO X., TANG M., ZHU D., WANG H., YANG X., CHEN B. Variation in Pollution Status, Sources, and Risks of Soil Heavy Metals in Regions with Different Levels of Urbanization. *Science of the Total Environment*, **866** (03), 161355, **2023**.