Original Research

Changes in Soil Fungal Communities in the Rhizosphere of *Platycladus orientalis* Plantation Forests of Different Stand Ages

Yue-xin Zhang¹**, Xiu-zhi Ma¹*, Yun-xia Ma², Cui-wei Li²

¹College of Forestry, Inner Mongolia Agricultural University, Hohhot, 10010, China ²College of Desert Governance, Inner Mongolia Agricultural University, Hohhot, 10010, China

Received: 25 March 2025 Accepted: 31 July 2025

Abstract

Soil fungi play a vital role in nutrient cycling and plant growth within forest ecosystems. Understanding the dynamics of soil fungal communities in Platycladus orientalis forests of different forest ages can provide a theoretical basis for the management, restoration, and sustainable development of oriental leopard plantation forests. In the present study, rhizosphere soil samples were collected from 10-, 23-, 35-, and 46-year-old P. orientalis plantations to investigate the diversity and community structure of soil fungi using Illumina MiSeq sequencing technology. The results showed that soil available nitrogen, total nitrogen, nitrate nitrogen (NN), total potassium, available potassium, and soil organic matter contents increased significantly with increasing stand age, while total phosphorus (TP) content decreased significantly. Furthermore, available phosphorus (AP) content initially increased and then decreased; the relative abundance of pathotrophs in the soil gradually increased and then stabilized, the saprotrophs increased gradually, and the symbiotrophs initially increased and then decreased. Furthermore, distance-based redundancy analysis identified soil pH, NN, and AP contents as the primary factors influencing fungal community composition. Structural equation modeling revealed a significant positive correlation between pH and symbiotrophs. Contrastingly, pH and TP were significantly negatively correlated with pathotrophs. P. orientalis forest plantation significantly altered soil physicochemical properties, which subsequently affected the soil fungal community. Increasing the soil pH and TP content in the study area would promote the growth and development of P. orientalis forests.

Keywords: Platycladus orientalis, forest plantation, stand age, soil fungal community

Introduction

The rhizosphere, a microregion a few millimeters from the surface of the root axis, serves as a unique area where the plant root system interacts with the soil microbial community [1]. These interactions

*e-mail: 1668681793@qq.com **e-mail: 15248178840@163.com Tel.: +86-417-15248178840

between rhizosphere microbes and the plant root system ultimately determine nutrient uptake, resource cycling, and plant health [2]. Rhizosphere microorganisms inhabit the rhizosphere soil of various plant species, playing a pivotal role in plant growth [3]. Specifically, rhizosphere soil fungi are important decomposers that regulate plant nutrient uptake; thereby, promoting growth and development, enhancing host plant disease resistance, and maintaining microecosystem balance within the rhizosphere.

Soil fungal community diversity maintains soil productivity and is an important indicator for evaluating the stability, health, and sustainability of soil ecosystems [4]. Previous studies have demonstrated that changes in the soil environment can considerably affect the composition of soil fungal communities [5-7]. Additionally, plant community succession can indirectly affect the soil environment, thereby altering fungal composition [8]. In monospecific plantation forests, the forest age is a significant factor influencing changes in soil fungal communities [9]. For example, Wang et al. [10] revealed that the development of Pinus tabuliformis plantation forests altered the soil nutrient content in rhizosphere soil, indirectly affecting rhizosphere fungal composition and diversity. Moreover, Zhang et al. [11] identified that the cycling and allocation capacity of soil elements also change significantly with stand age, further affecting fungal community structure. However, fungal species are specific and complex and respond differently to different soil environments even within the same ecosystem [12]. Determining soil microbial community dynamics that reflect changes in forest age and soil properties is important when assessing interactions between above- and below-ground communities [13]. Therefore, it is necessary to understand the changes in soil fungal communities during the development of plantation forests and to identify the main factors affecting these changes.

Platycladus orientalis, commonly known as lateral cypress and incense cedar, is an evergreen tree in the Cupressaceae family. It is widely distributed in China, with favorable characteristics, such as a well-developed root system, drought and barrenness tolerance, a long growth cycle, and the ability to maintain strong forest stability. Therefore, its seedlings are preferred for the afforestation of barren mountains, playing a key role in artificial vegetation construction [14]. The rhizosphere soil microenvironment of P. orientalis may undergo changes with forest age, which could potentially affect the interactions between rhizosphere microorganisms and lateral cypress. Therefore, the present study investigates the soil environment and soil fungal community structure of P. orientalis plantation forests of different ages in Jungar Banner, China. Specifically, we aimed to explore the rhizosphere soil environment and community diversity of P. orientalis plantation forests across different stand ages and to determine the correlations between rhizosphere soil and soil fungal community succession during the development of P. orientalis plantation forests.

Materials and Methods

Study Site

The study site was located in Jungar Banner (110°05′-111°27′E, 39°16′-40°20′N), Ordos City, Inner Mongolia Autonomous Region, China (Fig. 1). The study site receives average annual sunlight, temperature, and precipitation of over 3,000 h, 6.2–8.7 °C, and 400 mm, respectively. The soil types present in the study area include chestnut, calcium, and windy sandy soils. *P. orientalis* has been planted extensively for over 40 years in the study area, and many planted *P. orientalis* forests of different ages have been created through effective




Fig. 1. Geographical locations of the sampling sites.

Table 1. Sampling plot details.

Stand age (years)	Tree height (m)	Diameter at breast height (cm)	Altitude (m)	Slope degree (°)	Slope aspect (°)
10	5.40 ± 0.19	10.41 ± 0.24	1,315	23	30 west-northwest
23	8.33 ± 0.25	14.32 ± 0.32	1,294	23	50 north-east
35	13.8 ± 0.36	17.65 ± 0.55	1,305	27	30 west-northwest
46	18.0 ± 0.51	20.05 ± 0.97	1,297	25	60 north-east

Note: Data is expressed as mean \pm standard deviation.

management practices. The study area has sparse herbaceous plants, with dominant *Puccinellia tenuiflora* and *Leymus chinensis* species.

Soil Sample Collection and Analysis

In August 2024, we utilized the spatiotemporal intergenerational method to investigate forest stands in the area. Stands under similar conditions, including management style, slope, slope direction, and soil type, were selected to avoid introducing variation due to differences in ecological conditions. We identified four ages of P. orientalis plantations based on records from the local forestry department: 10 years old (10a), 23 years old (23a), 35 years old (35a), and 46 years old (46a). Detailed information is provided in Table 1. Three plants of *P. orientalis* (growth condition, diameter at breast height were similar) were randomly selected in the sample plots of each forest age, and within 1 meter from the trunks of the trees, a 0-30 cm soil profile was dug, and roots with fine roots were taken from near the main root system, the large pieces of soil attached to the roots were shaken off, and the soil adhering to the fine roots was collected into self-sealing bags with a sterile brush. The soil from the 4 directions was thoroughly mixed to obtain this P. orientalis rhizosphere soil sample, and a total of 12 soil samples were obtained. The soil samples were halved, and one half was placed into a sterile selfsealing bag and stored at -80 °C for DNA extraction. The other half was air-dried and passed through a 2-mm sieve for soil chemical analysis.

Total nitrogen (TN) and effective N were determined by the Kjeldahl method [15] and NaOH alkaline diffusion method [16], respectively, and soil effective N included available nitrogen (AN) and nitrate nitrogen (NN). Total phosphorus (TP) and available phosphorus (AP) were determined by NaHCO₃ leaching-molybdenum-antimony resistivity [17]; total potassium (TK) was determined by NaOH melting-molybdenum-antimony resistivity [18]; available potassium (AK) was determined by NH₄OAc leaching-flame photometry [19]; soil organic matter (SOM) by K₂Cr₂O₇ oxidation [20]; and soil pH by pH meter.

DNA Extraction and High-Throughput Sequencing

ITS1F We employed the (5'-GGAAGTAAAAGTCGTAACAAGG-3') and ITS2R (5'-GCTGCGTTCTTCATCGATGC-3) primers the amplification of the fungal ITS1-ITS2 region. The polymerase chain reaction (PCR) system comprised 25 μL of reaction buffer (5 μL [Thermo Fisher Scientific]), GC high-fidelity buffer (5 µL [Thermo Fisher Scientific]), Q5DNA polymerase (0.25 µL), deoxynucleotide triphosphate (10 µM, 2 µL), and double-distilled H₂O (8.75 µL). PCR cycling program was performed as follows: pre-denaturation at 98 °C for 5 min, followed by 30 cycles of denaturation at 98 °C for 30 s, annealing at 55 °C for 45 s, and extension at 72 °C for 45 s. Subsequently, the separation of PCR products by gel electrophoresis was performed. Then, target fragments were excised and recovered using the Axygen Gel Recovery Kit (Thermo Fisher Scientific). The Shanghai Personalbio Technology Co., Ltd., Shanghai, China, performed Illumina MiSeq sequencing analysis.

Data Preprocessing and Bioinformatics Analysis

The raw sequencing data were processed using QIIME2 software (Jacobs School of Engineering, La Jolla, CA, USA). First, the cutadapt plug-in (Jacobs School of Engineering) was used for primer excision. Next, quality control, denoising, removal of low-quality sequences and chimeras, amplicon sequence variant clustering with 100% similarity, and comparative annotation against the Greengenes database were performed using the DADA2 plug-in (Jacobs School Engineering). Alpha diversity indices were calculated using QIIME2 software (Jacobs School of Engineering). Subsequently, samples were analyzed using linear discriminant analysis based on taxonomic composition under different grouping conditions using Lefse software (Huttenhower Lab, Boston, MA, USA). Functional classification of each sequence, including trophic mode and growth status, was annotated using the FUNGuild online database (http://github.com/ UMNFuN/FUNGuild).

Table 2. Soil properties of Platycladus orientalis across different stand ages.

Hd	8.32±0.73a	8.30±0.02a	8.28±0.01a	8.20±0.02b	NS	
SOM (g/kg)	5.86±0.05d	6.67±1.94c	9.95±2.12b	12.73±1.27a	* *	(F = 84.55; P < 0.01)
AK (mg/kg)	74.75±4.34b	59.32±6.66c	96.29±3.99b	121.58±11.84a	* *	(F = 35.97; P < 0.01)
TK (g/kg)	14.53±0.64b	14.96±1.39b	16.42±0.47ab	17.55±0.73a	*	(F = 4.72; P < 0.05)
AP (mg/kg)	20.45±0.42d	24.51±0.39b	30.84±0.03a	22.21±0.93c	* *	(F = 46.99; P < (F = 137.68; P < (F = 4.72; P < 0.01)
TP (g/kg)	1.29±0.04a	1.12±0.07a	0.73±0.09b	0.52±0.08c	* *	(F = 46.99; P < 0.01)
NN (g/kg)	5.79±0.78b	10.54±0.98b	13.76±0.91a	15.97±1.12a	*	(F = 62.81; P < 0.01)
TN (g/kg)	0.87±0.02b	1.43±0.02a	1.56±0.06a	1.76±0.12a	**	(F = 62.81; P < 0.01)
AN (g/kg)	50.13±1.73d	55.87±0.88c	64.31±2.60b	77.83±1.75a	* *	(F = 83.30; P < (F = 62.81; P < 0.01)
Stand age	10a	23a	35a	46a	Soil properties along the chronosequence	

Note: Data are expressed as mean \pm standard deviation. Different lowercase letters indicate significant differences at p < 0.05 level of significance based on analysis of variance. NS: Non-significant regressions along the chronosequence; *: p < 0.05; **: P < 0.01. AN-available nitrogen, TN-total nitrogen, NN-nitrate nitrogen, TP-total phosphorus, AP-available phosphorus, TK-total potassium, AK-available potassium, SOM-soil organic matter

Data Analysis

Significance differences were evaluated using SPSS 22.0 software (SPSS, Chicago, IL, USA). The soil fungal community alpha diversity index was analyzed using Mothur software (National Institutes of Health, Bethesda, MD, USA). Additionally, the similarity in soil fungal community abundance at different forest ages was assessed using cluster analysis in the Vegan program package (https://cran.r-project.org/web/pac kages/vegan/ index.html) in R software (Version 3.3.1 [RStudio, Boston, MA, USA]). Principal coordinate analysis (PCoA) and distance-based redundancy analysis (db-RDA) were performed to explore community structure. Furthermore, structural equation modeling (SEM) was performed using AMOS software (Version 20.0, Amos Development Corporation, Meadville, PA, USA). Oneway analysis of variance was employed to identify significant correlations between forest age and soil physicochemical properties and fungal alpha diversity.

Results

Soil Properties

The physicochemical properties of the rhizosphere soil in P. orientalis plantation forests exhibited significant changes with increasing forest age (Table 2). Specifically, results revealed that increasing forest age significantly (p<0.05) increased the TN, NN, AN, TK, and SOM contents in the rhizosphere soil. Contrastingly, the TP content significantly decreased; whereas, AP initially increased and then decreased, reaching its peak in the 35a forest stand. In addition, no significant correlation between rhizosphere soil pH and stand age in P. orientalis.

General Characterization of Rhizosphere Fungal Communities

Illumina MiSeq sequencing revealed 690,400 fungal sequences, which were classified into 740 operational taxonomic units (OTUs). Among these, 385 OTUs were identified at the genus level, distributed across 9 phyla, 38 classes, 89 orders, 213 families, and 312 genera. Rhizosphere soil samples from 10a, 23a, 35a, and 46a *P. orientalis* plantations contained 149, 150, 178, and 198 fungal genera, respectively (Fig. 2). Among these genera, 71 were shared across the 4 forest ages. Additionally, the rhizosphere soil samples from 10a, 23a, 35a, and 46a *P. orientalis* plantations contained 25, 18, 36, and 57 endemic genera of fungi, respectively.

Fungal Diversity and Structure

The results of the diversity analysis showed that forest age significantly affected the Chaol and Pielou's evenness indices (p < 0.05) but had no significant effect




Fig. 2. Venn diagram of the soil fungal composition of *Platycladus orientalis* plantation forests at the genus level across different stand ages.

on the Shannon and Simpson indices (p > 0.05) (Table 3). Diversity indices showed a trend of first increasing and then decreasing with forest age. Among these, the Chaol index of soil fungi at 35a was significantly higher than that of the other three forest ages (p < 0.05); the Pielou index of soil fungi at 23a and 35a was significantly higher than that at 46a (p < 0.05); the Shannon index of soil fungi at 35a was significantly higher than that at 46a (p < 0.05), while the Simpson index of soil fungi at 46a was significantly lower than that of the other three forest ages (p < 0.05).

Significant differences in the rhizosphere soil fungi at the phylum level were observed in P. orientalis plantation forests of varying stand ages (Fig. 3). The abundance of soil fungi from the phyla Basidiomycota and Mortierellomycota increased initially and then declined with increasing stand age of P. orientalis plantations, reaching their highest levels in the 23a stand. Conversely, the Ascomycota abundance decreased initially and then increased, reaching its peak in the 46a stand. Furthermore, soil fungi from P. orientalis plantation forests of varying stand ages differed significantly (p < 0.05) at the genus level (Fig. 4). Penicillium, Pseudogymnoascus, and Mortierella

were the top three dominant genera in the 10a stand considering relative abundance. In the 23a stand, *Penicillium*, *Mortierella*, and *Pseudogymnoascus* were the dominant genera; whereas, *Paraboeremia*, *Bradymyces*, and *Penicillium* were dominant in the 35a age stand. In the 46a stand, *Knufia*, *Niesslia*, and *Penicillium* were the dominant genera. *Penicillium* was distributed in four forest stands and had a relatively high relative abundance.

The functional classification annotation results demonstrated that the rhizosphere soil fungi of *P. orientalis* plantations predominantly exhibited three trophic types: pathotrophs, symbiotrophs, and saprotrophs (Fig. 5). The proportions of these functional groups varied significantly across different forest ages. Specifically, as forest age increased, the relative abundance of pathotrophs gradually increased and then stabilized. Meanwhile, the saprotrophs exhibited a gradual increase, and symbiotrophs initially increased and decreased, reaching their peak at 35a. Additionally, a higher relative abundance of fungi not identified as functional taxa was observed in the soil of the 46a stand than in the soil of the other stand ages.

The PCoA results revealed significant differences in the rhizosphere soil community composition across different stand ages of P. orientalis plantation forests $(R^2 = 0.346, p = 0.018)$ (Fig. 6). Notably, the soil fungal communities of the 10a and 23a stands were notably distant from those of the 35a and 46a stands in the PCoA results. Contrastingly, the soil fungal communities between the 10a and 23a, and those between the 35a and 46a stands, were in close proximity, indicating a greater similarity in their community compositions. Furthermore, Hierarchical cluster analyses also showed more similarity in soil fungal community structure between the 10a and 23a stands, and great similarity between 35a and 46a stands (Fig. 7). The differences in the soil fungal composition among P. orientalis plantation forests of different stand ages primarily presented as changes in the relative abundances of Paraboeremia, Mortierella, Pseudogymnoascus, and Niesslia.

Table 3. Diversity index of rhizosphere soil fungi associated with <i>Platycladus orientalis</i> plantation forests at different stand ages.
--

	Chao1	Pielou	Shannon	Simpson
10a	360.69±12.67b	0.66±0.02ab	5.57±0.14ab	0.94±0.02a
23a	364.01±17.95b	0.69±0.03a	5.83±0.23ab	0.96±0.01a
35a	415.82±46.00b	0.69±0.03a	5.96±0.17a	0.96±0.08a
46a	462.59±20.12a	0.62±0.02b	5.49±0.14b	0.93±0.02b
P value	p < 0.05	P < 0.05	0.087	0.246

Note: Data are expressed as mean \pm standard deviation. Different lowercase letters indicate significant differences at p < 0.05, based on analysis of variance.




Fig. 3. Composition of soil fungal communities at the phylum level across different stand ages of Platycladus orientalis plantation forests.





Fig. 4. Composition of soil fungal communities at the genus level across different stand ages of Platycladus orientalis plantation forests.



R²=0.346, P=0.018 ● 10a 23a 0.3 35a 0.2 0.1 PC02 (13.6%) 0.0 -0.1 -0.2-0.3-0.4 -0.3 -0.2 -0.1 0.0 0.1 0.2 0.3 0.4 PCo1 (20.2%)

Fig. 5. Functional taxonomic composition of soil fungal communities across different stand ages of *Platycladus orientalis* plantation forests.

Fig. 6. Principal coordinates analysis of soil fungal communities in *Platycladus orientalis* plantation forests across different stand ages.




Fig. 7. Hierarchical clustering analysis of soil fungal communities in Platycladus orientalis plantation forests across different stand ages.

Correlation Among Fungal Community Composition and Soil Nutrients

The db-RDA results showed that there were relationships between the distribution of rhizosphere soil fungal communities and soil physicochemical properties across different stand ages of P. orientalis plantation forests (Fig. 8). The first principal axis accounted for 18.3% of the variance, with the first and second explanatory axes jointly accounting for 30%. Notably, TP content (p < 0.01), pH (p < 0.01), and NN content (p < 0.05) significantly influenced soil fungal communities.

The results of correlation analysis showed that soil fungal genera were significantly correlated with soil physicochemical properties at different forest ages (Fig. 9). The relative abundance of *Penicillium*, *Roesleria*, and *Fusarium* showed significant positive correlation with soil pH (p < 0.05), *Humicola* showed highly significant positive correlation with soil pH (p < 0.01), *Xenopolyscytalum* and *Exophiala* showed significant negative correlation with pH (p < 0.05); *Cadophora* and *Tetracladium* were significantly positively correlated with TP content (p < 0.05), *Cladophialophora* and *Cyphellophora* were significantly negatively




Fig. 8. Distance-based redundancy analysis of the soil fungal community structure in *Platycladus orientalis* plantation forests across different stand ages.

correlated with TP content (p < 0.05); Fusarium and Cladophialophora were significantly positively correlated with the content of TN and NN (p < 0.05); Scutellinia showed significant positive correlation with soil TN (p < 0.05); Pseudogymnoascus and Cadophora were significantly negatively correlated with NN content (p < 0.05).

Finally, the SEM results showed that increasing stand age significantly increased soil NN content and Chao 1 index diversity while reducing TP content (Fig. 10). Additionally, an increase in pH and TP content significantly decreased the relative abundance of pathotrophs, while the increase in pH also significantly increased the relative abundance of symbiotrophs and the Pielou index.




Fig. 9. Correlation analysis between soil fungal community composition and soil properties in *Platycladus orientalis* plantation forests. *: p < 0.05, **: p < 0.01.




Fig. 10. Structural equation model of plantation age, soil physicochemical properties, and soil fungal community composition and diversity in *Platycladus orientalis* plantation forests. The numbers provided for each component in the model reflect the coefficient of determination R^2 ; meanwhile, the numbers adjacent to the arrows indicate the path coefficients. Red, green, and gray arrows represent positive, negative, and no significant correlations, respectively. NN: nitrate nitrogen, TP: total phosphorus. *: p < 0.05, **: p < 0.01.

Discussion

The compositional structure of forest ecosystems, environment, and soil physicochemical properties have been found to change with increasing forest age. The present study revealed that soil AN, TN, NN, TK, and SOM contents increased with increasing age of P. orientalis. As the age of P. orientalis forests increases, the amount of litter under the canopy also increases. Litter on the ground is decomposed by microorganisms and root secretions and converted into organic nutrients [21-22]. Soil SOM and N elements mainly come from ground litter, resulting in a continuous increase in AN, NN, TN, and SOM content in the soil. The primary means by which soil acquires P comes from the weathering of rocks and leaching, and the method of acquisition is relatively singular [23]. Soil TP content decreased with increasing stand age; whereas, AP content initially increased and then decreased. TP content significantly decreased between 23a and 35a, but AP content increased significantly during this period. However, although the TP content in the soil of the 35a stand was low, conducive soil conditions contributed to the release of P from apoplast or soil minerals, ultimately increasing AP content and enhancing plant growth. Therefore, P. orientalis plantation succession ameliorates P deficiency in plants in the study area [24]. However, the soil TP contents of 35a and 46a stands were 0.73 g/kg and 0.52 g/kg, respectively, which is considered severely phosphorusdeficient soil. Furthermore, the study area had no supplementation of the P element by weathering of mineral rocks. Therefore, ecosystem management in Jungar Banner *P. orientalis* plantation forests should focus on increasing P content in the soil to meet the future demand for forest productivity.

In this study, the Chaol, Pielou-e, Shannon, and Simpson indices unexceptionally increased with the extension of forest age and decreased subsequently. This is similar to the research findings of Cao et al. [25] on the characteristics of soil fungal communities in Cunninghamia lanceolata forests at dissimilar stages of development. The reason may reside in that as the forest ages, the ecosystem gradually matures throughout the early stages, providing fungi with diverse habitats and resources, ultimately giving rise to an increase in the diversity index. As the ecosystem matures further, certain fungal species may exhibit a downturn trend due to competition, and the community structure may stabilize, resulting in a decline in the diversity index [26]. In addition, both PCoA analysis and hierarchical clustering analysis results suggest that the soil fungal community compositions of the 35a and 46a stands are more similar, which demonstrates that the fungal community in the rhizosphere soil of *P. orientalis* tends to stabilize as forest age increases. The aforementioned phenomenon illustrates that the underground ecosystem has reached a relatively stable equilibrium. Nonetheless, this study focused attention on 10a to 46a P. orientalis plantation forests. In addition, it is imperative to conduct more profound research on the soil fungal community succession, so as to understand the future succession process of *P. orientalis* plantation forests.

In this study, Ascomycota, Basidiomycota, and Mortierellomycota were dominant in all forest ages, which is consistent with the results of the study by Xu et al. [27]. It is noteworthy that Ascomycota demonstrated exceptional abundance in all four forest ages, all exceeding 65%. Ascomycota is the dominant fungal phylum globally, occupying the largest proportion of soil fungal species. It plays a paramount role as an important decomposer in soil, thereby contributing tremendously to the degradation of organic matter and soil nutrient cycling [28]. Basidiomycota is a diverse group of fungi that can form symbiotic relationships with plant roots to form mycorrhizae, which contribute to plant stress tolerance. Basidiomycota displays superior degradation capabilities for difficult-todegrade substances such as cellulose and lignin. Most species of Mortierella within the Mortierellomycota are beneficial to plants [29]. The abundant presence of these three types of fungi can not only heighten the rate of litter decomposition in the soil, ameliorate soil fertility and utilization, but also bring more nutrients for plant growth, and push ahead soil restoration subsequent to afforestation [30]. In this study, there were differences between dominant genera of different forest ages. The dominant genera in the 10a and 20a stands were Penicillium and Mortierella. The dominant genera in the 35a stand were Pseudogymnoascus and Penicillium. The predominant genera in the 40a stand were Knufia and Penicillium. Penicillium was present in all four forest ages. Penicillium is extensively acknowledged for its production of enzymes and antibiotics, and is crucial for biodegradation, nitrogen metabolism, and promoting plant growth. The abundance of Penicillium was higher in 10a and 23a stands, but conspicuously decreased in 35a and 46a stands. The abundance of dominant fungal genera in the soil reveals noticeable fluctuations and succession patterns with the extension of forest age. This phenomenon may be attributable to the disparities in the adaptive capacity of various fungal groups to physicochemical environmental conditions, with changes in understory microenvironmental factors becoming progressively disadvantageous for their growth and reproduction, eventually triggering a reduction in the abundance or replacement by other

Soil fungal communities employ various nutritional strategies to adapt to environmental changes. In the present study, the rhizosphere soil fungi of *P. orientalis* plantation forests were primarily categorized into three groups: pathotrophs, symbiotrophs, and saprotrophs [32]. The relative abundance of saprotrophs was the highest among the four stand ages, aggregating in large numbers in the 46a stand. Notably, in the 46a stand, there was a notable increase in the amount of apoplastic and necrotic tissues, which provided a substantial pool of SOM for saprotrophic fungi. The apoplastic decomposition by saprotrophs resulted in the release of easily accessible carbon sources into the rhizosphere soil environment, resulting in a peak in N

and SOM contents, indicating that the decomposition of apoplastic matter by specific fungal taxa became more efficient over time [33, 34]. Ascomycota fungi, which are predominantly saprotrophs, decompose a wide range of difficult-to-degrade substances and are important decomposers of SOM [35]. The results of the present experiment similarly indicated that Ascomycota abundance reached a maximum in stand 46a. Moreover, there are mutual restraints between different trophic fungi, with the large enrichment of symbiotrophs having antagonistic effects on plant pathotrophs; thereby, reducing their colonization and playing a positive role in the development of *P. orientalis* plantation forests [36].

Results from db-RDA analysis illustrate that soil pH, TP, and NN stand as primary factors influencing soil fungal communities. In detail, soil pH serves as a pivotal indicator of soil acidity or alkalinity. Apart from that, dissimilar fungal species display varying degrees of adaptability to pH conditions. For this reason, soil pH potentially imposes direct influences on the species composition and abundance of fungal communities in the soil [37]. Numerous studies have shown that soil pH is an important driver of fungal community composition [38, 39]. Apart from that, soil pH is not only bound up with multiple genera, but also reveals a conspicuous and positive correlation with the dominant genus Humicola, further demonstrating that soil pH plays a paramount role in altering soil fungal community composition [40, 41]. NN may indirectly influence fungal communities by altering soil microenvironments (e.g., pH or nitrogen form). Some studies have put forth a viewpoint that NN nutrients can inhibit the growth of pathogens (such as Fusarium), potentially through resource competition or activation of plant resistance [42]. Correlation analysis also suggested a striking and negative association between NN and Fusarium. Soil TP is a key nutrient element that can drive fungal community structure by influencing different fungal functional groups. Other investigations have also drawn a conclusion that TP indirectly affects fungal communities by regulating soil pH and organic matter [43].

As suggested by SEM results, forest age has a positive effect on soil NN and TP content and negatively impacts soil pH. Consistent with the results of studies on Pinus sylvestris var and Cunninghamia lanceolata plantations, pH values exhibited a remarkable decline with the increase of forest age [44, 45]. The reason is that as forest age increases, vegetation cover increases, litter decomposition and root exudate accumulation occur, thus giving rise to soil acidification [46]. As the age of the forest increases, the number and activity of ammonia-oxidizing archaea (AOA) in the soil may change accordingly. Associated studies have revealed that AOA in acidic soils have a higher affinity for ammonia and are better suited to surviving in acidic environments [47]. As the acidity of the rhizosphere soil of P. orientalis increases, the activity of AOA gets more dramatic, thereby elevating the production of NN. As trees age, their root systems become more

developed, and root exudates and root residues (such as root hairs and fine roots) also add phosphorus to the soil. These root inputs function as one of the paramount sources of phosphorus in the soil [48]. An abatement in soil pH may trigger the binding of phosphorus with elements such as iron and aluminum, thereby elevating the content of available phosphorus in the soil. On top of that, a remarkable and positive correlation was detected between pH and the relative abundance of symbiotrophs, while pH and TP demonstrated a striking and negative correlation with pathotrophs. The relationship between soil pH and fungal community structure is complex, as it may influence fungal growth and activity through both direct and indirect mechanisms. Optimal pH levels may provide more advantageous growth conditions for symbiotic fungi while inhibiting the activity of pathogenic fungi [49]. TP content in soil imposes direct influence on plant growth, development, and health. High TP levels may facilitate plant growth, reinforce plants' resistance to pathogenic fungi, and thereby lessen the relative abundance of pathogenic fungi [50]. As a consequence, we speculate that increasing the soil pH and TP content in this region could not only propel the growth and development of P. orientalis but also lower the abundance of pathotrophs while lessening the abundance of symbiotrophs. It is crucial to figure out these details for strengthening the management of P. orientalis plantation forests.

Conclusions

In this study, the long-term growth of *P. orientalis* plantation changed rhizosphere soil nutrients as well as fungal community composition. Soil pH, NN, and AP content were the main factors affecting fungal community composition. Among them, pH was significantly positively correlated with the relative abundance of symbiotrophs, and pH and TP content were significantly negatively correlated with the relative abundance of pathotrophs. In conclusion, the results of the present study revealed the relationship between fungal composition and soil properties in *P. orientalis* plantation forests, and provided a theoretical basis for the management and sustainable development of *P. orientalis* plantation forests in this region.

Acknowledgments

This study was supported by the Third Xinjiang Scientific Expedition Program (2022xjkk0403) and the Natural Science Foundation of Inner Mongolia Autonomous Region, "Structural and Functional Analyses of *Xanthoceras sorbifolium* rhizosphere Arbuscular Mycorrhizal Fungi Community in Changing Environments and Research on Effect Mechanisms" (2023QN03049).

Conflict of Interest

The authors declare no conflicts of interest.

References

- 1. MENDES R., GARBEVA P., RAAIJMAKERS J.M. The rhizosphere microbiome: Significance of plant beneficial, plant pathogenic, and human pathogenic microorganisms. FEMS Microbiology Reviews, **37** (5), 634, **2013**.
- CASTAÑO C., LINDAHL B.D., ALDAY J.G., HAGENBO A., MARTÍNEZ DE ARAGÓN J., PARLADÉ J. Soil microclimate changes affect soil fungal communities in a Mediterranean pine forest. New Phytologist, 220 (4), 1211, 2018
- EDWARDS J., JOHNSON C., SANTOS-MEDELLÍN C., LURIE E., SUNDARESAN V. Structure, variation, and assembly of the root-associated microbiomes of rice. Proceedings of the National Academy of Sciences, 112 (8), 911, 2015.
- WANG J.X., GAO J., ZHANG H.Q., TANG M. Changes in rhizosphere Soil Fungal Communities of Pinus tabuliformis Plantations at Different Development Stages on the Loess Plateau. International Journal of Molecular Sciences, 23 (12), 6753, 2022.
- LIU C., WANG S., YAN J., HUANG Q., SHEN Q. Soil fungal community affected by regional climate played an important role in the decomposition of organic compost. Environmental Research, 197, 111076, 2021.
- SUN K., FU L., SONG Y., YUAN L., WANG K. Effects of continuous cucumber cropping on crop quality and soil fungal community. Environmental Monitoring and Assessment, 193 (7), 436, 2021.
- BASTIDA F., LÓPEZ-MONDÉJAR R., BALDRIAN P., ANDRÉS-ABELLÁN M., JEHMLICH N., TORRES I.F. When drought meets forest management: Effects on the soil microbial community of a Holm oak forest ecosystem. Science of The Total Environment, 662, 276, 2019.
- 8. MARTÍNEZ-DIZ M.P., ANDRÉS-SODUPE M., BUJANDA R., DÍAZ-LOSADA E., GRAMAJE D. Soilplant compartments affect fungal microbiome diversity and composition in grapevine. Fungal Ecology, **41**, 234, 2019
- DONG H.Y., GE J.F., SUN K., WANG B.Z., XUE J.M., WAKELIN S.A., WU J.S., SHENG W.X., LIANG C.F., XU Q.F., JIANG P.K., CHEN J.H., QIN H. Change in root-associated fungal communities affects soil enzymatic activities during Pinus massoniana forest development in subtropical China. Forest Ecology and Management, 482, 118817, 2021.
- WANG J.X., GAO J., ZHANG H.Q., TANG M. Changes in rhizosphere Soil Fungal Communities of Pinus tabuliformis Plantations at Different Development Stages on the Loess Plateau. International Journal of Molecular Sciences, 23 (12), 6753, 2022.
- 11. ZHANG J., ZHAO N., LIU C., YANG H., LI M., YU G. C:N:P stoichiometry in China's forests: from organs to ecosystems. Functional Ecology, **32** (1), 50, **2018**.
- 12. KENNEDY P.G., MIELKE L.A., NGUYEN N.H. Ecological responses to forest age, habitat, and host vary by mycorrhizal type in boreal peatlands. Mycorrhiza, 28 (3), 315, 2018.
- 13. REICHERT J.M., MORALES M.S., LIMA C.L.R., DE BASTOS F., SANTANA C.A., DE ARAUJO E.F. Soil

- morphological, physical and chemical properties affecting Eucalyptus spp. productivity on entisols and ultisols. Soil & Tillage Research, **226**, 105575, **2023**.
- 14. CHEN X.D. Study on the Community Structure of Arbuscular Mycorrhizal Fungal under Different Afforestation Types and Their Effects on Litter Decomposition. Northwest A&F University, 2018.
- BREMNER J.M., MULVANEY C.S. Methods of soil analysis: Chemical and microbiological properties. American Society of Agronomy Inc & Soil Science Society of America Inc, 595, 1982.
- NELSON D.W., SOMMERS L.E. Total carbon, organic carbon and organic matter, in: Methods of Soil Analysis Part 2. Chemical and Microbial Properties, 539, 1982.
- 17. PARKINSON J.A., ALLEN S.E. A wet oxidation procedure suitable for the determination of nitrogen and mineral nutrients in biological material. Communications in Soil Science and Plant Analysis, 6 (1), 1, 1975.
- 18. BAO S.D. Soil Agrochemical Analysis, third ed.[M]. China Agriculture Press. 2005.
- 19. WU S., ZHANG Y., TAN Q., SUN X., HU C. Biochar is superior to lime in improving acidic soil properties and fruit quality of satsuma mandarin. Science of The Total Environment, 714, 136722, 2020.
- WALKLEY A., BLACK I.A. An examination of the Degtjareff method for determining soil organic matter, and A proposed modification of the chromic acid titration method. Soil Science, 37 (1), 29, 1934.
- FUSS C.B., LOVETT G.M., GOODALE C.L., OLLINGER S.V., LANG A.K., OUIMETTE A.P. Retention of nitrate-N in mineral soil organic matter in different forest age classes. Ecosystems, 22 (6), 1280, 2019.
- 22. BURKE D.J., WEINTRAUB M.N., HEWINS C.R., KALISZ S. Relationship between soil enzyme activities, nutrient cycling and soil fungal communities in a northern hardwood forest. Soil Biology and Biochemistry, 43 (4), 795. 2011.
- 23. LEI X.T., ZHANG H., CHEN M., GUO L., ZHANG X.G., JIANG Z.H. The efficiency of sequential extraction of phosphorus in soil and sediment: Insights from the oxygen isotope ratio of phosphate. Journal of Soils and Sediments, 20 (3), 1332, 2020.
- 24. SCHLATTER D.C., HANSEN J.C., SCHILLINGER W.F., SULLIVAN T.S., PAULITZ T.C. Common and unique rhizosphere microbial communities of wheat and canola in a semiarid Mediterranean environment. Applied Soil Ecology, 144, 170, 2019.
- CAO Y.X. Characterization of soil microbial communities in Cunninghamia lanceolata plantations of different ages. Central South University of Forestry and Technology, 2022.
- 26. DICKIE I.A., MARTÍNEZ-GARCÍA L.B., KOELE N., GRELET G.A., RICHARDSON S.J. Mycorrhizas and mycorrhizal fungal communities throughout ecosystem development. Plant and Soil, 367 (1-2), 11, 2013.
- XU G.L., MI Y.T., LIU L.Z., WEN L., LI Y.J., XU J.F. Study on the structure and driving factors of soil fungal community in different grassland types in Inner Mongolia. Acta Agrestia Sinica, 31 (7), 1977, 2023.
- HOU Q., WANG W.X., YANG Y., HU J., BIAN C.S., JIN L.P., LI G.C., XIONG X.Y. Rhizosphere microbial diversity and community dynamics during potato cultivation. European Journal of Soil Biology, 96, 100, 2020
- 29. HANNULA S.E., HEINEN R., HUBERTY M., BEZEMER T.M. Persistence of plant-mediated microbial

- soil legacy effects in soil and inside roots. Nature Communications, 12 (1), 5686, 2021.
- ZHANG J., LIU Q., WANG D.Z., ZHANG Z.D. Soil Microbial Community, Soil Quality, and Productivity along a Chronosequence of Larix principis-rupprechtii Forests. Plants, 12 (16), 2913, 2023.
- 31. DONG G.Q., ZHANG Y.T., ZHANG H.J., CHAI X., ZHOU S.Z., TONG Z. Effects of Chinese fir plantations with different densities on understory vegetation and soil microbial community structure. Chinese Journal of Plant Ecology, 45 (9), 1006, 2021.
- 32. NGUYEN N.H., SONG Z., BATES S.T., BRANCO S., TEDERSOO L., MENKE J., SCHILLING J.S., KENNEDY P.G. FUNGuild: An open annotation tool for parsing fungal community datasets by ecological guild. Fungal Ecology, 20, 241, 2016.
- 33. AMARASINGHE A., KNOX O.G.G., FYFE C., DE BRUYN L.L., WILSON B.R. Response of soil microbial functionality and soil properties to environmental plantings across a chronosequence in south eastern Australia. Applied Soil Ecology, 168, 104100, 2021.
- 34. CHU H., WANG C., WANG H., CHEN H., TANG M. Pine wilt disease alters soil properties and root-associated fungal communities in Pinus tabulaeformis forest. Plant and Soil, 404 (1-2), 237, 2016.
- 35. BEIMFORDE C., FELDBERG K., NYLINDER S., RIKKINEN J., TUOVILA H., DORFELT H., GUBE M., JACKSON D.J., REITNER J., SEYFULLAH L.J., SCHMIDT A.R. Estimating the Phanerozoic history of the Ascomycota lineages: Combining fossil and molecular data. Molecular Phylogenetics and Evolution, 78, 386, 2014.
- 36. FREW A., POWELL J.R., GLAUSER G., BENNETT A.E., JOHNSON S.N. Mycorrhizal fungi enhance nutrient uptake but disarm defenses in plant roots, promoting plant-parasitic nematode populations. Soil Biology and Biochemistry, 126, 123, 2018.
- 37. KYASCHEVKO J., CLEMMENSEN K.E., HAGENBO A., KARLTUN E., LINDAHL B.D. Shift in fungal communities and associated enzyme activities along an age gradient of managed Pinus sylvestris stands. ISME Journal, 11 (4), 863, 2017.
- 38. FAROOQ T.H., KUMAR U., SHAKOOR A., ALBASHER G., ALKANTANI S., RIZWANA H. Influence of intraspecific competition stress on soil fungal diversity and composition in relation to tree growth and soil fertility in subtropical soils under Chinese fir monoculture. Sustainability, 13 (19), 10688, 2021.
- 39. DANG P., VU N.H., SHEN Z., LIU J., ZHAO F., ZHU H. Changes in soil fungal communities and vegetation following afforestation with Pinus tabulaeformis on the Loess Plateau. Ecosphere, 9 (8), e02401, 2018.
- 40. ROUSK J., BÅÅTH E., BROOKES P.C., LAUBER C.L., LOZUPONE C., CAPORASO J.G. Soil bacterial and fungal communities across a pH gradient in an arable soil. The ISME Journal, 4 (10), 1340, 2010.
- GEML J., PASTOR N., FERNANDEZ L., PACHECO S., SEMENOVA T.A., BECERRA A.G. Large-scale fungal diversity assessment in the Andean Yungas forests reveals strong community turnover among forest types along an altitudinal gradient. Molecular Ecology, 23 (10), 2452, 2014
- 42. QIAO R.Y., SONG Z., CHEN Y.M., XU M.Z., YANG Q.Q., SHEN X.L., YU D.Y., ZHANG P.D., DING J.H., GUO H. Planting density effect on poplar growth traits and soil nutrient availability, and response of microbial

- community, assembly and function. BMC Plant Biology, 24 (1), 1, 2024.
- 43. LIU J., ZHU Z.C., CHEN J.F. Effects of soil total phosphorus and available phosphorus on soil fungal functional groups. Fujian Forestry Science and Technology, 50 (1), 1, 2023.
- 44. YUAN G., ZHENG Y., SUN X. Unveiling Microbial Dynamics: How Forest Aging Shapes the Microbial Communities of Pinus massoniana. Ecology and Evolution, 15 (3), e71132, 2025.
- 45. ZHANG Y., CHEN Y., AA B., MA X., ZHANG H., LIU Q., MAO R. The succession patterns and drivers of soil bacterial and fungal communities with stand development in Chinese fir plantations. Plant and Soil, **500** (1-2), 547, **2024**.
- 46. ZHAO H., ZHOU Y.C., REN Q.F. Evolution of soil microbial community structure and functional diversity in Pinus Massoniana plantations with age of stand. Acta

- Pedologica Sinica, 57 (1), 227, 2020.
- 47. HE J.Z., HU H.W., ZHANG L.M. Current insights into the autotrophic thaumarchaeal ammonia oxidation in acidic soils. Soil Biology and Biochemistry, 55, 146, 2012.
- 48. CHEN H., TANG H.Y., GUO J.H., PAN C., WANG R.H., WU Y.H., YU Y.C. Root exudates' roles and analytical techniques progress. Soil, 55 (2), 225, 2023.
- 49. JI K.P., WEI Y.Q., WANG X., LIU Y., SUN R., LI Y.W., LAN G.Y. Distribution patterns of fungal community diversity in the dominant tree species Dacrydium pectinatum and Vatica mangachapoi in tropical rainforests. Microbiology Spectrum, 13 (1), e03104-24, 2025.
- 50. LI L.L., LUO Z.Z., LI L.L., NIU Y.N., ZHANG Y.Q., HE R.Y., LIU J.H., NIAN L.L. Long-term phosphorus fertilization reveals the phosphorus limitation shaping the soil micro-food web stability in the Loess Plateau. Frontiers in Microbiology, 15, 1472473, 2024.