

Original Research

Investigating the Impact of Some Habitat Characteristics on Distribution of *Stachys pilifera* Benth Using the BMLR Model in Iran

Shahab Mirinejad¹, Esfandiar Jahantab^{2*}, Mohammad Reza Mahmoudi³,
Mehrdokht Najafpour Navaei⁴, Mohammad Mehdi Rahimi⁵,
Mohsen Sharafatmandrad⁶

¹Research Division of Natural Resources, Kohgiluyeh and Boyerahmad Agricultural and Natural Resources Research and Education Center, AREEO, Yasooj, Iran

²Department of Range and Watershed Management, Faculty of Agricultural Sciences, Fasa University, Fasa, Iran

³Department of Statistics, Faculty of Science, Fasa University, Fasa, Iran

⁴Research Institute of Forests and Rangelands, Agricultural Research, Education and Extension Organization (AREEO), Tehran, Iran

⁵Department of Agronomy, Arsanjan Branch, Islamic Azad University, Arsanjan, Iran

⁶Faculty of Natural Resources, University of Jiroft, Jiroft, Iran

Received: 6 July 2017

Accepted: 2 November 2017

Abstract

Stachys pilifera is an endemic plant to Iran. The main purpose of this study was to investigate the impacts of some habitat characteristics on the distribution of *S. pilifera* using the BMLR model in Kohgiluyeh and Boyer-Ahmad Province, Iran. Two transects were randomly established in each region. 10 quadrats sized 1 x 1 m² were established along the transect. In each plot, canopy cover (%) of *S. pilifera* was recorded. 25 soil samples were gathered in the region to evaluate soil characteristics. The associated pixel of each plot was extracted and added to a prepared spreadsheet of a field database. The data gathered from experiments were fed into the computer item by item according to their own values. To determine the effect of environmental factors on the distribution of *S. pilifera* species, the researchers run a set of backward multiple linear regressions. The results of MLR to investigate the significance of was not significant (P-Value >0.05). Then the most insignificant term in the first run (silt) is removed and the operation run again. After removing silt, the results of the BMLR model are seen by considering the effect of other factors, and some of terms are insignificant again. The process is continued step by step to the point that there are only significant terms. Results showed that the effects of rain, sand, P, Fe, Zn, Mn, and evaporation on the distribution of *S. pilifera* species were significant. Rain, P, and Fe have positive effects (positive coefficient), while sand, Zn, Mn, and evaporation have negative effects (negative coefficient).

Results showed that the BMLR appropriately modeled the canopy cover of *S. pilifera* based on RMSE, R^2 , and , which were 0.999, 0.998, and 2.1395, respectively.

Keywords: *Stachys pilifera*, habitat characteristics, BMLR model

Introduction

The genus *Stachys* (Lamiaceae) contains about 300 species and is extensively distributed in tropical and subtropical countries [1]. This genus in Iran is represented by 34 species, 13 of which are endemic. *Stachys pilifera* Benth is one of the endemic species to Iran. *Stachys pilifera* is a fruticose and pregnant perennial herbaceous plant of the Lamiaceae family. This plant species can be found in mountainous habitats, especially moist areas that are affected by humidity resulting from surface or subsurface waters. The plant has a short stem covered with simple and slender leaves, pink to white flowers that all aerial organs have a pleasant and distinctive smell (an aromatic compound). There are many studies on essential oil of this plant. Cis-chrysanthenyl acetate (25.2%) and trans-verbenol (19.7%) are the main compounds in the essential oil of *Stachys pilifera*.

The aerial organs of the plant are used in Iranian traditional medicine as herbal tea for the treatment of various diseases such as asthma and rheumatoid arthritis, plus infections [2]. A biological study has shown the potent antioxidant, antimicrobial, and antitumor activities of the n-butanolic extract of *S. pilifera* in in-vitro conditions [3].

Although there are many studies about environmental factors controlling plant distribution of endemic and endangered plant species [4-7], this kind of knowledge on *S. pilifera* is rare. For example, the distribution map of *S. pilifera* in Central Zagros in the Kohgiluyeh Region is prepared, and its autecology is determined [8]. Studying the cultivation possibility of *S. pilifera* showed that lowland ecotypes have more capability for greenhouse cultivation and that these ecotypes have more essential oil [9]. Phonological studies indicate that *S. pilifera* starts to germinate in mid-April and continues to vegetative growth until June; inflorescences appears in late June and seeds will mature until late September [8].

As mentioned above, there are not many studies about the relationship between *S. pilifera* and controlling environmental factors. Therefore, studying the environmental requirements of this plant species is important. The main purpose of this study is to evaluate the impact of habitat characteristics on the distribution of *S. pilifera* using the BMLR model in Kohgiluyeh and Boyer-Ahmad Province, Iran.

Material and Methods

Study Area

This study was carried out on Kohgiluyeh and Boyer-Ahmad Province in central Zagros in southwestern Iran. The *S. pilifera* habitats were determined by asking experts and local people and also field investigations. Thus, *S. pilifera* habitats were clarified and separated as one of the sampling points. Detailed information about each habitat is presented in Table 1.

Data Collection

Habitat characteristics of the *S. pilifera* species were determined in order to achieve behavior and environmental conditions. Therefore, vegetation and different environmental factors including topography, pedology, and climatic characteristics were measured.

Tow transects were randomly established in each region. Ten 1x1 m quadrats were established along each transect. The recorded parameter in each quadrat was *S. pilifera* canopy cover (%). The location of each quadrat was recorded with GPS. 25 soil samples were taken (5 samples in each habitat) in order to identify soil characteristics such as EC [10], pH [11], organic matter [12], soil texture [13], P [14], N [10], K [15], Mn, Cu, Zn, and Fe [16]. For climatological and land shape data, the height, slope and aspect maps were generated by creating DEM. The associated pixel values of each plot

Table 1. Detailed information about *S. pilifera* habitats in Kohgiluyeh and Boyer-Ahmad Province in central Zagros.

Habitat	Longitude	Latitude	Mean elevation (m asl)	Dominant vegetation type
Pazanan Sepidar	30°32'9.3"	51°29'8.1"	2575	<i>Prangos uloptera</i>
Delon Sisakht	30°52'11.1"	51°28'54.9"	2585	<i>Daphne mucronata</i>
Road of Margon to Lodab	30°58'35.3"	51°2'32.1"	2090	<i>Gundelia turnefortii</i>
Veze	30°30'24.5"	51°42'33.4"	2420	<i>Dorema aucheri</i>
Chin village	31°5'20.8"	51°48'38.2"	2415	<i>Prangos Sp</i>

were extracted and added to a prepared spreadsheet of field database.

Data Analysis

A set of backward multiple linear regressions were run to determine the effect of environmental factors on the distribution of *S. pilifera* species. The Kolmogorov-Smirnov test was applied to verify the normality of canopy cover. This condition is an essential condition for using the parametric methods as BMLR. Canopy cover had normal distribution. Therefore, parametric methods can be applied for statistical analysis. Regression analysis generates an equation to describe the statistical relationship between canopy cover and environmental factors. Regression results indicate the direction, size, and statistical significance of the relationship between canopy cover and environmental factors. The sign of each coefficient indicates the direction of the relationship. Coefficients represent the mean change in canopy cover for one unit of change in one environmental factor while holding other environmental factors in the model constant. P-value for each coefficient tests the null hypothesis that the coefficient is equal to zero (no effect). Therefore, low p-values suggest that one environmental factor is a meaningful addition to the model.

In MLR, variables were removed and added to the regression model for the purpose of identifying a useful subset of environmental factors (the predictors). Multiple linear regression (MLR) is a flexible method of data analysis that may be appropriate whenever a quantitative variable (the dependent or response variable) is to be examined in relationship to any other factors (X_1, X_2, \dots, X_k – the independent or explanatory or predictor variables). The general equation of MLR is presented by

$$Y_i = \beta_0 + \beta_1 X_{1i} + \beta_2 X_{2i} + \dots + \beta_k X_{ki} + \varepsilon_i$$

...for n observations $n = 1, \dots, n$,

...where $\beta_0, \beta_1, \dots, \beta_k$ are model parameters (coefficients) and $\varepsilon_i, i = 1, \dots, n$, are the random components of the model that follow independent normal distributions with mean 0 and variance σ^2 .

The dataset is used to estimate the coefficients $\beta_0, \beta_1, \dots, \beta_k$. The general equation of the predictive MLR model is presented by

$$\hat{Y}_i = b_0 + b_1 X_{1i} + b_2 X_{2i} + \dots + b_k X_{ki}$$

...where b_0, b_1, \dots, b_k are estimations of model parameters, and \hat{Y}_i is the predicted value of Y_i .

The MLR model can be rewritten in matrix form

$$Y = X\beta + \varepsilon$$

...where $Y = (y_1, \dots, y_n)^T$ is vector of responses, X is a $n \times (k + 1)$ full rank design matrix with the first

column given by $(1, \dots, 1)^T$, and the l^{th} ($2 \leq l \leq k + 1$) column given by, $(x_{l-1,1}, \dots, x_{l-1,n})^T$, $\beta = (\beta_0, \dots, \beta_k)^T$ is vector of unknown parameters, and $\varepsilon = (\varepsilon_1, \dots, \varepsilon_n)^T$ is vector of random errors. Also $\hat{Y} = X\hat{\beta}$, where $\hat{Y} = (\hat{y}_1, \dots, \hat{y}_n)^T$ is vector of predicted values, and $\hat{\beta} = (b_0, \dots, b_k)^T$ is vector of coefficients. We note that in the model without interception ($\beta_0 = 0$), column $(1, \dots, 1)^T$ should be removed from matrix X .

The ordinary least squares (Maximum Likelihood) estimation of the coefficient vector β is given by:

$$\hat{\beta} = (X^T X)^{-1} X^T Y$$

This is a common approach, and it assumes that there are enough measurements to say something meaningful about β .

As can be seen, the MLR model contains linear effects of X_1, X_2, \dots, X_k . But perhaps some of these effects are not significant (P-Value >0.05). In this case, the backward method (BMLR) is used and the ineffective parameters are removed gradually. The final model has parsimonious accurate parameters. The statistical BMLR model was applied using R and Minitab software.

Table 2. Descriptive statistics of environmental variables for all *S. pilifera*.

	N	Minimum	Maximum	Mean	Std. Deviation
Elevation	25	2,078	2,595	2,424.68	181.794
Rain	25	766	964	887.68	69.916
Temperature	25	11	20	14.60	4.113
PH	25	7.10	7.35	7.1888	.07020
EC	25	.32	1.88	.7364	.49942
OC (organic matter)	25	1.07	2.97	2.0092	.60643
Clay	25	28	46	36.28	5.144
Silt	25	50	65	56.08	3.696
Sand	25	3	15	7.76	2.554
K	25	194	529	327.28	112.316
P	25	4.22	19.96	13.0208	6.25604
N	25	.10	.85	.3091	.25617
Fe	25	8.17	19.80	14.1928	4.03759
Zn	25	.19	1.50	.6336	.47668
Cu	25	.40	1.22	.8072	.24234
Mn	25	5.37	14.49	10.9112	3.00740
Humidity	25	52	63	58.80	4.537
Evaporation	25	1,458	1,803	1,597.96	149.574
Canopy cover	25	43	68	55.36	6.720

Results and Discussion

Descriptive Statistics

Descriptive statistics of measured variables are shown in Table 2.

Important Factors on Distribution of *S. pilifera* Species

In order to evaluate the importance of different environmental factors on the distribution of *S. pilifera*, its canopy cover was considered as the response variable. Also, the other variables were continuous predictor variables. The general equation of MLR was presented by

$$\text{Canopy Cover}_i = \beta_0 + \beta_1 \text{Elev}_i + \dots + \beta_{18} \text{Evapo}_i + \varepsilon_i$$

First, all terms were entered and created the model. Table 4 shows the results of MLR to investigate the significance of β_0 . As can be seen in Table 4, intercept (constant) was not significant (P-Value >0.05). In other words, the model didn't need β_0 .

Table 4. Results of MLR for investigating the significance of β_0 .

	Coefficient	P-Value
Constant	-24.6421	0.993
Elevation	0.0636	0.631
Rain	0.7317	0.132
Temperature	3.5585	0.918
PH	4.3846	0.853
EC	18.3102	0.650
OC	-1.3487	0.963
Clay	-0.2384	0.882
Silt	-0.0610	0.971
Sand	-1.7221	0.389
K	-0.2491	0.365
P	5.8969	0.263
N	-30.1243	0.636
Fe	3.4602	0.509
Zn	-33.0304	0.445
Cu	-17.3814	0.693
Mn	-9.6971	0.433
Humidity	-1.2606	0.954
Evaporation	-0.3835	0.720

Table 5. The results of the first run of BMLR.

	Coefficient	P-Value
Elevation	0.0629	0.508
Rain	0.7299	0.075
Temperature	3.7060	0.896
PH	4.3749	0.840
EC	18.1040	0.557
OC	-1.2537	0.960
Clay	-0.2382	0.872
Silt	-0.0586	0.969
Sand	-1.7188	0.342
K	-0.2484	0.307
P	5.8871	0.214
N	-29.8105	0.545
Fe	3.4709	0.461
Zn	-33.0846	0.401
Cu	-17.2099	0.638
Mn	-9.7372	0.361
Humidity	-1.4385	0.872
Evaporation	-0.3917	0.465

Table 5 shows the results of the BMLR model after removing β_0 . As Table 5 indicates, by considering the effects of other variables, the effect of some factors was not significant (P-Value >0.05).

Then the most insignificant term (term with the most P-Value) in the first run (Silt) is removed and the operation run again. Table 6 shows the results of the BMLR model after removing Silt. As can be seen in Table 6, by considering the effect of other factors, some terms are not significant again.

Then the most insignificant term in the second run (OC) is removed and run the operation again. The process is continued step by step to the point that there are only significant terms. Table 7 indicates the summary of excluded variables in the BMLR method in each step.

Table 8 shows the results of the last run. As can be seen in Table 8, the effects of rain, sand, P, Fe, Zn, Mn, and evaporation on the distribution of *S. pilifera* species were significant. Rain, P, and Fe have positive effects (positive coefficient), and sand, Zn, Mn, and evaporation have negative effects (negative coefficient).

Therefore, the equation of the final BMLR model was

$$\begin{aligned} \text{Canopy Cover} = & 0.3301\text{Rain} - 1.3765\text{Sand} \\ & + 6.1675\text{P} + 2.6558\text{Fe} - 54.4292\text{Zn} - \\ & 6.3337\text{Mn} - 0.1511\text{Evaporation} \end{aligned}$$

Table 6. Results of the second run of BMLR.

	Coefficient	P-Value
Elevation	0.0623	0.476
Rain	0.7242	0.041
Temperature	4.2758	0.849
PH	4.4191	0.827
EC	17.5547	0.491
OC	-0.7520	0.970
Clay	-0.1835	0.651
Sand	-1.6559	0.024
K	-0.2448	0.240
P	5.9520	0.150
N	-29.1160	0.495
Fe	3.3913	0.389
Zn	-33.7229	0.312
Cu	-16.3938	0.557
Mn	-9.9082	0.273
Humidity	-1.3221	0.860
Evaporation	-0.4009	0.370

Now, the goodness of the fitted model was assessed by the coefficient of determination (R^2), adjusted coefficient of determination (R^2_{adj}), root mean square error ($RMSE$), and residual analysis. A lower $RMSE$ value and higher R^2 and R^2_{adj} values and independent normal residuals with stable variance are regarded as showing goodness of fitted predictive model. As can be

Table 7. Summary of excluded variables in the BANCOVA method.

Step	Excluded variable	P-Value
1	All variables are included	
2	Silt	0.969
3	OC	0.970
4	Humidity	0.853
5	PH	0.761
6	Clay	0.520
7	Temperature	0.331
8	Elevation	0.388
9	EC	0.208
10	Cu	0.443
11	N	0.267
12	K	0.158

Table 8. The results of the last run of BMLR.

	Coefficient	P-Value
Rain	0.3301	0.002
Sand	-1.3765	<0.001
P	6.1675	0.005
Fe	2.6558	0.017
Zn	-54.4292	0.006
Mn	-6.3337	0.001
Evaporation	-0.1511	0.018

Table 9. Indexes for the goodness of fitted BMLR models.

R Square	Adjusted R Square	RMSE
0.999	0.998	2.1395

seen in Table 9, it seems that the BMLR nicely modeled the canopy cover based on $RMSE$, R^2 , and R^2_{adj} .

To investigate the normality of residuals, probability plot and different statistical tests (Anderson-Darling, Kolmogorov-Smirnov, Shapiro-Wilk) were used. As can be seen in Figure 1, the normal probability plot (2) satisfied the normality of residuals, since the points are close to the line. The normality was also satisfied with different statistical tests (P -Value >0.05).

As Fig. 2 indicates, the plot of residuals versus observations' order was completely random about horizontal axis. Therefore, the independence of residuals is satisfied.

Fig. 3 shows the plot of the residuals versus the fitted values. As can be seen, the points are completely random about of the horizontal axis, and the stability of the variance is satisfied. Therefore, the BMLR nicely modeled the canopy cover.

Cognition of vegetation communities and evaluating their interaction with environments is known as a crucial factor for achieving sustainable rangeland management in order to introduce the appropriate species for reclaiming a degraded area. Furthermore, knowledge about environmental factors can be used to predict success and failure of growing species.

In order to recognize the various native species and vegetation cover associated with them, these species can be used in ecosystem improvement and pastoral ecosystem revival. The present study examined the relationship between environmental variables and *S. pilifera* species distribution in Kohgiluyeh and Boyer-Ahmad Province in central Zagros, Iran.

Results showed that the effects of rain, sand, P, Fe, Zn, Mn, and evaporation, on the distribution of *S. pilifera* species were significant. Rain, P, and Fe have positive effects (positive coefficient), and sand, Zn, Mn, and evaporation have negative effects (negative coefficient). Results showed that the BMLR nicely




Fig. 1. Normal probability plot of residuals for BMLR model. If the points on the plot depart from a straight line, the normality assumption may be invalid.

modeled the canopy cover based on $RMSE$, R^2 , and F , which were 0.999, 0.998, and 2.1395, respectively.

The results are in line with the findings of Darvishi et al. (2013). In this regard, Darvishi et al. (2013) investigated a study on the environmental factors contributing to the distribution of *Thymus kotschyanus* in Taleghan Basin, Iran. Results demonstrated that factors involving slope, altitude, organic matter, lime content, nitrogen content, and soil texture have the highest impact on vegetation characteristics [5].

Our results showed that the most significant factors are rain and evaporation. In this regard Jahantab et al. (2015) have shown that the relationship between number (density) and canopy cover percentage of *Kelussia odoratissima* mozaft with height, slope, rainfall, and evaporation were statistically meaningful at 95% confidence interval. Then the factors such as rain and evaporation were important factors on the distribution of plant species [17].

Soil factors play a major role in ecological processes and are closely associated with the growth and distribution of plants. Our results indicated that soil properties such as sand, P, Fe, Zn, and Mn were effective in distributing *S. pilifera* species. In our study area, the differences of climate and topographical features are relatively small, so distribution of *S. pilifera* species may be potentially affected by soil properties. In this regard Zare et al. (2011) stated that soil texture is one of the most effective factors in distributing *Z. eurypterum*, *P. aucheri*, and *A. spinosa* species [18]. Gholinejad et al. (2012) stated that soil texture is one of the most effective factors in the distribution of *As. gossypinus* – *Gu. tournefortii*, *Br. tomentellus*-*Fe. ovina* and *Fe. hausknechtii* – *Pr. Ferulacea* [19].

This study shows that organic carbon was not effective on the distribution of *S. pilifera* species. Based

on our results, Komae et al. (2012) [20], Fahimi-Pour et al. (2010) [21], and Ahmadi et al. (2010) [22] introduced organic matter and depth as the most important soil factors that influence the separation of vegetation types in the study area. Tao et al. (2017) stated that the distribution of herbs on sand dunes was dominantly influenced by sand dune topography (slope position, convexity, and relative height), soil nutrients (total nitrogen and phosphorus), and deep-rooted shrubbery (*Ephedra distachya*) [23].

Plant species distribution over a high geographical range is controlled by climatic factors — mainly temperature and rainfall. In a small range, species distribution is related to edaphic factors [24]. Therefore, these environmental factors are important not only in detecting plant species distribution variations with spatial scale, but also for providing insight into the environmental requirements of the species needed for




Fig. 2. Plot of residuals versus observation order for BMLR model; this is a plot of all residuals in the order that the data were collected and can be used to find non-random error.




Fig. 3. Plot of residuals versus fitted values for BMLR model.

successful ecological restoration and the establishment of plantations [25].

Our results showed that elevation was not effective distribution of *S.pilifera*, while some researchers reported that topographic factors are related to vegetation communities, as found previously by many studies [26]. Zhong-hua et al. (2013) showed that topographical factors were more important than edaphic ones in affecting local plant distribution on steep slopes with extensive rock outcrops, while edaphic factors were more influential on gentle slope and relatively thick soil over rock in subtropical karst forest [27].

Schumann et al. (2016) reported that elevation and precipitation were the most influencing environmental factors on potential distributions of both juvenile and adult richness [28]. The appearance of a plant group in a given area isn't accidental, but occurs in response to changes in climatic, topographic, edaphic, and biotic parameters. In fact, vegetation groups are determined by the combined effects of a variety ecological factors.

In this research we investigated the impact of some habitat characteristics on the distribution of *S. pilifera* Benth using the BMLR model. In this regard, Akhter et al. (2017) stated that the application of the species distribution model may provide policymakers and conservationists with a useful tool for predicting future distribution (at both local and regional scales) of poorly known species with high preservation concerns [29].

Bobrowski et al. (2017) stated that that analyzing and understanding the environmental factors driving the current distribution of *Betula utilis* is crucial for the prediction of future range shifts of *B. utilis* and other treeline species, and for deriving appropriate climate change adaptation strategies [30].

Conclusions

Our study investigated the impact of some habitat characteristics on the distribution of *Stachys pilifera* Benth, an endemic medicinal plant in Iran, using the BMLR model in Kohgiluyeh and Boyer-Ahmad

Province. The BMLR model showed that environmental factors such as soil (sand, P, Fe, Zn, and Mn) and climate (rain and evaporation) are effective for the distribution of *S. pilifera* species. Rain, sand, P, Fe, Zn, Mn, and evaporation have significant effects on the distribution of *S. pilifera*. Rain, P, and Fe have positive effects, which means that *S. pilifera* will increase with their increase. Sand, Zn, Mn, and evaporation have negative effects (negative coefficient), in which case *S. pilifera* will decrease with their increase. The present study addressed some aspects of relationships between environmental factors and the distribution of *S.pilifera* species in Kohgiluyeh and Boyer-Ahmad Province. It was anticipated that this finding could be used as a tool for predicting the probability of existence and the absence of this plant species in similar ecosystems.

Acknowledgements

This work was supported by the Kohgiluyeh-va-Boyerahmad Agricultural and Natural Resources Research and Education Center. The authors are grateful for all their help and assistance in all aspects of manuscript preparation.

References

1. KUMAR D., BHAT Z.A., KUMAR V., KHAN N.A., CHASHOO I.A., ZARGAR M.I., SHAH M.Y. Effects of *Stachys tibetica* essential oil in anxiety. *European Journal of Integrative Medicine*, **4** (2), e169, **2012**.
2. ZARGARI A. Medicinal plants. Tehran: Tehran University Press; **1992**.
3. FARJAM M.H., KHALILI M., RUSTAYIAN A., JAVIDNIA K., IZADI S. Biological activity of the n-butanolic extract of *Stachys pilifera*. *African Journal of Microbiology Research*, **5** (28), 5115, **2011**.
4. JAHANTAB E., SEPEHRI A., BARANI H., QASEMI ARYAN Y., FARAJOLAH A. An Introduction on the Endangered Medicinal Species of Mountain's Kelavs (*Kelussia odoratissima* Mozaff.) in Central Zagros in Kohgiluyeh Region, Iran. *Journal of Rangeland Science*, **2** (1), 409, **2011**.
5. DARVISHI L., ZARE CHAHOUKI M.A., JAFARI M., AZARNIVAND H., YOUSEFI VALIKCHALI M. Study on the Environmental Factors Contributing to Distribution of *Thymus kotschyanus* in Taleghan Basin, Iran. *Journal of Rangeland Science*, **24**; **4** (1), 82, **2013**.
6. AHMADI A., SHAHMORADI A.A., ZAREKIA S., AHMADI E., NATEGHI S. Autecological study of *Astragalus effusus* in rangelands of west Azarbaijan province, Iran. *Iranian Journal of Range and Desert Research*, **20** (1), 172, **2013**.
7. SHARIFI YAZDI M., SHAHMORADI A.A., ZAREKIA S. Autecology of *Ferula ovina* Boiss in Kerman Province. *Journal of redintegrate natural resources*, **5** (2), 57, **2014**.
8. MIRIRNEJAD S.H., NAJAFPORNAVEEI M., MOHTASHAMI R., DASTYARAN M. Investigation of Atacology of Extinction species *Stachys pilifera* L. in Kohgiluyeh and Boyer Ahmad Province. *National*

- Conference on Natural Products and Medicinal Plants, Iran, **2012**.
9. HEIDARI H. Identification of Morphological Characteristics and Determination of Essential Oil of *Stachys pilifera* L. in different habitats of Kohgiluyeh and Boyer Ahmad Province and Study on the Possibility of Potting in Greenhouse. Master's Thesis, faculty of agriculture, university of Yasuj. **2015**.
 10. BREMNER J.M., MULVANEY C.S. Nitrogen – total. Methods of soil analysis. Part 2. Chemical and microbiological properties. (methodsofsoilan2) 595, **1982**.
 11. KLUTE A. (ed.) Methods of soil analysis. Part I. physical and micrological methods. Second edition. Agronomy series **9**, ASA. Madison, Wis, **1986**.
 12. WALKLEY A., BLACK I.A. An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. Soil science, **1**; **37** (1), 29, **1934**.
 13. GEE G.W., BAUDER J.W. Particle-size analysis. Methods of soil analysis: Part 1 – Physical and mineralogical methods. (methodsofsoilan1) 383, **1986**.
 14. OLSEN S.R. Estimation of available phosphorus in soils by extraction with sodium bicarbonate. United States Department of Agriculture; Washington; **1954**.
 15. HELMKE P.A., SPARKS D.L. Lithium, sodium, potassium, rubidium, and cesium. Methods of Soil Analysis Part 3 – Chemical Methods. (methodsofsoilan3) 551, **1996**.
 16. PAGE A.L., MILLER R., KEENEY D.R. Part 2: Chemical and microbiological properties. Methods of soil analysis, 595, **1982**.
 17. JAHANTAB E., SEPEHRY A., MESDAGHI M., BARANI H. Impact of topographic and climatologic factors on Canopy cover percentage and the number of species *Kelussia odoratissima* mozaff in Kohgiluyeh. Journal of Iranian Plant Ecophysiological Research, **9** (36), 1, **2015**.
 18. ZARE S., JAFARI M., TAVILI A., ABBASI H., ROSTAMPOUR M. Relationship between environmental factors and plant distribution in arid and semiarid area (Case Study: Shahriyar Rangelands, Iran). American-Eurasian Journal of Agriculture & Environmental Science, **10** (1), 97, **2011**.
 19. GHOLINEJAD B., FARAJOLLAHI A., POUZESH H. Environmental factors affecting on distribution of plant communities in semiarid areas (Case study: Kamyaran rangelands, Iran). Annals of Biological Research, **3** (8), 3990, **2012**.
 20. KOMAEE F., HESHMATI GH., CHAHOOKY ZARE M.A. Effect of soil characteristics on the distribution of plant communities in Taleghan pastures. Third National Student Conference on Rangeland, Watersheds and Desert. Tehran University, Tehran. **2012**.
 21. FAHIMI POOR A., CHAHOOKY ZARE M.A., TAVILI A. Study of relation some of range plants to environmental factors (case study: Rangelands of Central Taleghan). Jour. Range Management, **4** (1), 32, **2010**.
 22. AHMADI H., KAMALI N., SALAJEGHE A., JAFARI M., SADEGHIPOUR A. Investigation on some environmental factors influencing distribution of plant species (Case study: Ghara Aghach watershed, Semrom, Iran). Watershed Management Researches Journal, (Pajouhesh & SAZANDEGI), **88**, 55, **2010**.
 23. TAO Y., WU G.L., ZHANG Y.M. Dune-scale distribution pattern of herbaceous plants and their relationship with environmental factors in a saline – alkali desert in Central Asia. Science of the Total Environment, **15**, 576, 473, **2017**.
 24. AYYAD M.A., EL-GHAREEB R. Micro variations in edaphic factors and species distribution in a Mediterranean salt desert. Oikos, **1**, 125, **1972**.
 25. TOLEDO M., PEÑA-CLAROS M., BONGERS F., ALARCÓN A., BALCÁZAR J., CHUVIÑA J., LEAÑO C., LICONA J.C., POORTER L. Distribution patterns of tropical woody species in response to climatic and edaphic gradients. Journal of Ecology, **1**; **100** (1), 253, **2012**.
 26. CUI B.S., ZHAI H.J., DONG S.K., CHEN B., LIU S.L. Multivariate analysis of the effects of edaphic and topographical factors on plant distribution in the Yilong Lake Basin of Yun-Gui Plateau, China. Canadian Journal of Plant Science, **1**; **89** (1), 211, **2009**.
 27. ZHANG Z.H., HU G., NI J. Effects of topographical and edaphic factors on the distribution of plant communities in two subtropical karst forests, southwestern China. Journal of Mountain Science, **1**; **10** (1), 95, **2013**.
 28. SCHUMANN K., NACOULEMA B.M., HAHN K., TRAORÉ S., THIOMBIANO A., BACHMANN Y. Modeling the distributions of useful woody species in eastern Burkina Faso. Journal of Arid Environments, **31** (135), 104, **2016**.
 29. AKHTER S., MCDONALD M.A., VAN BREUGEL P., SOHEL S., KJER E.D., MARIOTT R. Habitat distribution modeling to identify areas of high conservation value under climate change for *Mangifera sylvatica* Roxb. of Bangladesh. Land Use Policy, **31** (60), 223, **2017**.
 30. BOBROWSKI M., GERLITZ L., SCHICKHOFF U. Modeling the potential distribution of *Betula utilis* in the Himalaya. Global Ecology and Conservation, **31** (11), 69, **2017**.