

Further Studies on Aquatic Fungi in the River Biebrza within Biebrza National Park

B. Czeczuga, B. Kiziewicz*, B. Mazalska

Department of General Biology, Medical University, Kilińskiego 1, 15- 089 Białystok, Poland

Received: 7 February, 2003

Accepted: 18 March, 2003

Abstract

Zoosporic and conidial fungi species and environmental factors in 12 various sites of the River Biebrza in Biebrza National Park were studied. Samples of water were collected in autumn 1999 and spring 2000 for hydrochemical analysis in order to determine the number of fungal species. Buckwheat-seeds, hemp-seeds, cellophane and snake exuviae were used as bait during exposure in a laboratory. 199 lower aquatic fungus species were found in various sites of the River Biebrza. 16 species were recorded for the first time from Poland.

Keywords: aquatic fungi, hydrochemistry, river, Biebrza National Park

Introduction

The River Biebrza as a right tributary of the River Narew flows in the peaty valley and within Biebrza National Park forms diverse meanders and flooding, thus providing a wide environmental variety for flora and fauna. No wonder that the river valley is a habitat of rare plants and animal species. Preliminary investigations [1-4] have shown the presence of a number of aquatic fungus species in the waters of the park, which are new to Polish waters and rare in general. Therefore, we have decided to carry out a more detailed study of this water area, including different aquatic environments created by the River Biebrza within the Park.

Area of Study

For the purpose of our studies 12 different sites were chosen on the river Biebrza:

Site I - Dolistowo, river Biebrza in localities Dolistowo, round rivers stretch meadows, bank

grown *Phragmites australis* (Cav.) Trien. ex Steudel and here and there steps out *Schoenoplectus lacustris* (L.) Pala.

Site II - Dolistowo, old river closed, stagnant water and covered *Lemna minor* L. here and there steps out *Phragmites australis* (Cav.) Trien. ex Steudel and *Nuphar lutea* (L.) Sibth. et Sm.

Site III - Goniądz, river at bridge, not near rivers of building, at bank grows *Phragmites australis* (Cav.) Trien. ex Steudel and *Sagittaria sagittifolia* L.

Site IV - Osowiec, river at bridge, slowly flows, muddy bottom, at bank grows *Phragmites australis* (Cav.) Trien. ex Steudel.

Site V - Osowiec moat, stagnant water, covered *Lemna minor* L., here and there steps out *Phragmites australis* (Cav.) Trien. ex Steudel and *Nuphar lutea* (L.) Sibth. et Sm.

Site VI - Kanał Rudzki, very quick water flows, stony bottom.

Site VII - Olszowe Pole, river Biebrza in localities Olszowe Pole, muddy bottom, at bank grows *Phragmites australis* (Cav.) Trien. ex Steudel and *Hydrocharis morsus-ranae* L.

*Corresponding author

Table 1. Chemical composition (in mg l⁻¹) of water from the different sites (n=3) (autumn 1999).

XII Specification	Sites											
	I	II	III	IV	V	VI	VII	VIII	IX	X	XI	XI
Temperature, °C	12.3	10.5	12.5	13.0	12.7	12.5	13.0	12.5	12.5	13.0	12.7	11.5
pH	7.46	7.64	7.66	7.81	7.61	7.95	7.72	7.72	7.61	7.92	7.92	8.00
COD (Chemical Oxygen Demand)	6.47	21.56	7.45	9.40	10.58	7.45	7.84	8.43	8.23	9.02	11.17	6.20
CO ₂	11.0	17.6	11.0	8.8	13.2	11.0	8.8	11.0	11.0	11.0	15.4	13.2
Alkalinity in CaCO ₃ (mval l ⁻¹)	4.6	6.2	4.5	4.5	5.5	4.2	4.2	4.3	4.0	4.4	5.9	5.1
N-NH ₃	0.200	0.685	0.365	0.230	0.460	0.350	0.220	0.365	0.180	0.310	0.720	0.145
N-NO ₂	0.014	0.012	0.145	0.013	0.006	0.016	0.015	0.015	0.002	0.015	0.007	0.014
N-NO ₃	0.020	0.070	0.030	0.150	0.034	0.160	0.210	0.002	0.010	0.020	0.050	0.430
P-PO ₄	1.290	0.800	0.900	1.450	1.000	2.600	1.245	1.100	1.900	1.305	0.400	1.450
Sulphates	25.50	11.93	26.33	18.92	11.93	23.64	19.75	24.27	8.64	22.63	7.40	13.16
Chlorides	22.0	28.0	26.0	32.0	32.0	22.0	30.0	21.0	30.0	27.0	21.0	22.0
Total hardness in Ca	72.00	88.56	70.56	72.00	84.96	70.56	66.96	69.84	61.20	70.56	88.56	79.92
Total hardness in Mg	20.64	22.36	19.35	19.78	21.07	17.63	20.21	20.64	18.49	26.66	33.97	23.95
Fe (total)	0.60	1.33	0.45	0.45	0.73	0.45	0.40	0.45	0.20	0.45	1.33	0.48
Dry residue	298	429	279	273	342	266	291	263	232	273	372	287
Dissolved solids	296	389	241	266	342	264	264	258	229	251	370	266
Suspended solids	2	44	38	7	0	2	27	5	3	22	2	21

Site VIII - Olszowe Pole, old river joint with river Biebrza, steps out *Phragmites australis* (Cav.) Trien. ex Steudel, *Nuphar lutea* (L.) Sibth. et Sm. and *Nymphaea alba* L.

Site IX - Olszowe Pole, old river closed, stagnant water steps out *Phragmites australis* (Cav.) Trien. ex Steudel, *Nuphar lutea* (L.) Sibth. et Sm., *Nuphar pumila* (Timm) DC and *Nymphaea alba* L.

Site X - Biały Grąd - river Biebrza, muddy bottom, banks grown *Phragmites australis* (Cav.) Trien. ex Steudel and here and there steps out *Lemna trisulca* L.

Site XI - Biały Grąd, old river joint with river Biebrza, steps out *Phragmites australis* (Cav.) Trien. ex Steudel and *Nuphar lutea* (L.) Sibth. et Sm.

Site XII - Burzyn, dike on river Biebrza - in place of conscription of tests sampling banks concreted.

Materials and Methods

Samples of water were collected in autumn (October) 1999 and spring (May) 2000 for hydrochemical analysis

and the order to determine the number of fungal species. From site IV Osowiec and site VI Kanał Rudzki samples of water to determine zoosporic fungi species were collected every month from October 1999 to May 2000 and in autumn 2002. Nineteen parameters were determined in each water (Table 1) according to the generally accepted methods [5].

The water for analysis was coured into 3 containers for each sites. Water from each site was transferred to three 1.0 litre vessel and added of the baits and placed in the laboratory at ambient temperature. Buckwheat-seeds, hemp-seeds, cellophane and snake exuviae were used as bait during exposure in the laboratory.

The following procedures for the determination of the presence of fungus species on the fragments of baits were employed: during one month of exposure the fragments of baits were examined under a light microscope (once or twice a week) and mycelium of aquatic fungi growing on the baits was recorded. Identification of respective species was based on morphology and biometric data of antheridia and oogonia of the zoosporic fungi and conidiophores and conidia of the hyphomycetes. The following keys were used for zoosporic fungi - Johnson [6],

Table 2. Chemical composition (in mg l-1) of water from the different sites (n=3) (spring 2000).

Specification	Sites											
	I	II	III	IV	V	VI	VII	VIII	IX	X	XI	XI
Temperature, °C	19.0	23.0	17.0	18.0	16.5	16.5	19.0	21.0	21.0	19.8	18.5	17.0
pH	7.45	7.82	8.08	8.01	7.70	8.38	7.76	7.89	7.77	8.14	8.23	8.24
COD (Chemical Oxygen Demand)	9.11	23.76	10.30	9.50	12.28	8.71	9.70	11.88	12.67	14.62	26.14	13.07
CO ₂	13.2	11.0	4.4	6.6	13.2	4.4	8.8	11.0	11.0	10.5	6.6	8.8
Alkalinity in CaCO ₃ (mval l ⁻¹)	5.5	7.0	4.5	4.5	5.0	4.2	4.2	4.4	4.7	5.2	5.4	4.8
N-NH ₃	0.610	1.050	0.400	0.400	0.370	0.280	0.310	0.500	0.570	0.840	0.970	0.530
N-NO ₂	0.005	0.011	0.016	0.010	0.003	0.015	0.015	0.013	0.003	0.008	0.006	0.010
N-NO ₃	0.050	0.160	0.100	0.240	0.100	0.260	0.262	0.221	0.042	0.125	0.150	0.132
P-PO ₄	1.850	0.650	0.310	0.310	0.470	0.751	0.530	0.605	0.530	0.740	0.650	1.094
Sulphates	51.84	11.93	16.87	9.87	13.99	14.81	21.80	14.81	16.46	10.82	5.35	9.05
Chlorides	27.0	37.0	26.0	15.0	19.0	19.0	21.0	34.0	19.0	21.5	48.0	20.0
Total hardness in Ca	63.36	141.58	61.92	60.48	90.00	64.08	66.96	64.80	76.32	80.62	83.52	68.40
Total hardness in Mg	29.24	20.64	28.81	19.35	12.90	17.20	16.77	17.20	12.04	13.02	15.05	12.47
Fe (total)	1.00	2.15	0.50	1.20	0.45	0.30	0.50	0.70	0.70	0.70	1.00	0.85
Dry residue	252	453	252	314	359	235	340	315	317	320	336	272
Dissolved solids	226	437	215	294	318	211	237	308	301	298	326	268
Suspended solids	26	16	37	20	41	14	103	7	16	22	10	4

Sparrow [7], Waterhouse [8, 9], Seymour [10], Bedenek [11], Batko [12], Karling [13], Dick [14], Pystina [15] and for Hyphomycetes - Nilsson [16], Dudka [17, 18], Ingold [19], Carmichael et al. [20], Bräthen [21], Matsushima [22], and works of the authors who were the first to describe the respective species.

The investigate parameter data of water and fungal flora for these investigations were processed by cluster analysis [23].

Results

The physico-chemical properties of water at particular sites in the study period are presented in Tables 1, 2. Substantial differences were revealed in such water trophicity indices as oxidability and nutrients content (nitrogen, phosphates). In October oxidability was lowest at site XII Burzyn and in May at site VI Kanał Rudzki. Its maximum values in October were observed at site II Dolistowo, and in May at site X Biały Grąd and site II Dolistowo. The lowest ammonium nitrogen content in October was revealed at site XII Burzyn, in May at site VI Kanał Rudzki. Its highest content in October was

noted at site XI Biały Grąd, in May at site II Dolistowo. The content of nitrates in October was the lowest at site VIII Olszowe Pole, in May at site IX Olszowe Pole, and the highest in October at site XII Burzyn and in May in sites IV Osowiec and VI Kanał Rudzki. The lowest concentrations of phosphates were noted in October at site XI Biały Grąd and in May at sites III Goniądz and the IV Osowiec. The maximum values of phosphates in October were observed at site VI Kanał Rudzki, while in May at site I Dolistowo. In general, in the autumn the water at the sites examined showed higher oxidability and higher content of ammonium nitrate and phosphates, compared to the spring.

At 12 sites in Biebrza National Park 199 lower aquatic fungus species were found, including 108 zoosporic and 91 conidial belonging to Hyphomycetes (Table 3,4, Fig. 1). Most zoosporic fungus species were found to grow in water at sites IV Osowiec and VI Kanał Rudzki, fewest at site II Dolistowo. Most Hyphomycetes species were found at site I Dolistowo and site VIII Olszowe Pole (30 at each), fewest at site III Goniądz [18]. Some of the zoosporic and conidial fungi found in the present study are new to the waters of northeastern Poland. The group

Table 3. Zoosporic fungi found in water of particular sites of Biebrza National Park (a-autumn, s-spring, w-winter).

Species of fungi	Site											
	I	II	III	IV	V	VI	VII	VIII	IX	X	XI	XI
Chytridiomycetes												
Olpidiales												
<i>Rozella septigena</i> Cornu				w		a						
Chytridiales												
<i>Chytrium poculatus</i> Willoughby et Townley						w						
<i>Endochytrium ramosum</i> Sparrow	s	s										
<i>Karlingia hyalina</i> Karling						w						
<i>Nowakowskiella elegans</i> (Nowak.) Schr.						w						
<i>Phlyctochytrium aureliae</i> Ajello				w								
<i>Phlyctochytrium parasitans</i> Sparrow						w						
<i>Phlyctochytrium reinboldtae</i> Persiel				w								
<i>Polyphagus euglenae</i> Nowakowski						a,w						
<i>Rhizidium richmondense</i> Willoughby						w						
<i>Rhizophydium ampullaceum</i> (Braun) Fischer						a,w			s			
<i>Rhizophydium apiculatum</i> Karling		s										
<i>Rhizophydium carpophilum</i> (Zopf) Fischer	s		s	s			s	s	s	s		
<i>Rhizophydium coronum</i> Hanson				w								
<i>Rhizophydium elyensis</i> Sparrow				s								
<i>Rhizophydium globosum</i> (Braun) Rabenhorst	s		s	s	s		s		s	s	s	
<i>Rhizophydium keratinophilum</i> Karling						w						
<i>Septochytrium variabile</i> Berdan			s								s	
Blastocladales												
<i>Allomyces moniliformis</i> Coker et Braxton						a						
<i>Blastocladiopsis parva</i> (Whiffen) Sparrow	s		s	s,a,w	s	a,w						s
<i>Catenaria anguillulae</i> Sorokin	s	s	s	s,a		a						
<i>Catenaria verrucosa</i> Karling	s	s		w		a						
<i>Catenomyces persicinus</i> Hanson						a						
<i>Catenophlyctis variabilis</i> (Karling) Karling	s	s		s,w	s	s,a,w		s		s	s	s
Plasmodiophoromycetes												
Plasmodiophorales												
<i>Woronina polycystis</i> Cornu						a,w						
Oomycetes												
Saprolegniales												
<i>Achlya americana</i> Humphrey			s	a,w		a,w		s		s	s	s
<i>Achlya debaryana</i> Humphrey	s					w						s
<i>Achlya diffusa</i> Harvey ex Johnson				a								
<i>Achlya dubia</i> Coker						w						

Table 3 continues on next page...

<i>Achlya flagellata</i> Coker						a								
<i>Achlya glomerata</i> Coker				a		a,w								
<i>Achlya hypogyna</i> Coker et Pemberton				w		w								
<i>Achlya klebsiana</i> Pieters						a,w								
<i>Achlya megasperma</i> Humphrey				w										
<i>Achlya orion</i> Coker et Couch				w										
<i>Achlya polyandra</i> Hildebrand				s,w		w		s			s			
<i>Achlya prolifera</i> Coker				s										
<i>Achlya proliferoides</i> Coker				w		w								
<i>Achlya racemosa</i> Hildebrand	s			w		w				s				
<i>Achlya treleaseana</i> (Humphr.) Kauffman				w		s,w								
<i>Aphanodictyon papillatum</i> Huneycutt		s	s			s		s	s	s				
<i>Aphanomyces amphigynus</i> Cutter								s						
<i>Aphanomyces irregularis</i> Scott	s	s	s	s,w	s	w								s
<i>Aphanomyces laevis</i> de Bary	s	s	s	s,a,w	s	s,a,w	s	s			s	s	s	s
<i>Aphanomyces ovidestruens</i> Gickelh				w		w								
<i>Aphanomyces parasiticus</i> Coker		s		a,w		w								
<i>Aphanomyces stellatus</i> de Bary				a	s	s,a,w				s	s	s		
<i>Brevilegnia unisperma</i> (Coker et Braxton) Coker et Braxton	s			w		a,w		s	s	s	s			
<i>Calyptrolegnia achlyoides</i> (Coker et Couch) Coker						w								
<i>Cladolegnia unispora</i> (Coker et Couch) Johannes				w		w								
<i>Dictyuchus monosporus</i> Leitgeb				a,w		a,w				s	s	a	s	
<i>Isoachlya anisospora</i> (de Bary) Coker				s,w										
<i>Isoachlya monilifera</i> (de Bary) Kauffman						s								s
<i>Isoachlya toruloides</i> Kauffman et Coker				w										
<i>Leptolegnia caudata</i> de Bary	s				s									s
<i>Leptolegniella keratinophila</i> Huneycutt	s			a,w	s	s,a,w	s	s	s	s	s	s	s	s
<i>Protoachlya paradoxa</i> (Coker) Coker				w										
<i>Protoachlya polyspora</i> (Lindst.) Apinis		s		s,w		s,w								
<i>Pythiopsis cymosa</i> de Bary				w	s	a,w								
<i>Saprolegnia anisospora</i> de Bary				w		s,a,w	s			s				
<i>Saprolegnia asterophora</i> de Bary				w										
<i>Saprolegnia diclina</i> Humphrey				w		w								
<i>Saprolegnia ferax</i> (Gruih.) Thuret		s	s	s,a,w		a,w		a				s		
<i>Saprolegnia furcata</i> Maurizio				w		w								
<i>Saprolegnia glomerata</i> (Tiesenh.) Lund	s			s,w	s	w	s	s	s	s				
<i>Saprolegnia hypogyna</i> (Pringhs.) de Bary				w										
<i>Saprolegnia litoralis</i> Coker				w	s	w								
<i>Saprolegnia megasperma</i> Coker				w										

Table 3 continues on next page...

<i>Saprolegnia mixta</i> de Bary				w		a,w						
<i>Saprolegnia monoica</i> Pringsh.				w		w						
<i>Saprolegnia parasitica</i> Coker	s			a,w	s	s,a,w	s		s	s		
<i>Saprolegnia semihypogyna</i> S. Inaba et Tokumasu				a								
<i>Saprolegnia torulosa</i> de Bary						a						
<i>Traustotheca clavata</i> (de Bary) Humph.			s			s,w					s	s
Leptomitales												
<i>Apodachlya brachynema</i> (Hildebr.) Pring.				W								
<i>Leptomitus lacteus</i> (Roth) Agardh						w						
Lagenidiales												
<i>Lagenidium destruens</i> Sparrow								s			s	
<i>Lagenidium humanum</i> Karling				w								
<i>Olpidiopsis saprolegnia</i> (Braun) Cornu						a				a		
Peronosporales												
<i>Phytophthora cactorum</i> (Leb. et Cohn) Schröter						a						
<i>Pythiogeton autossytum</i> Drechsler							s					
<i>Pythiogeton nigricans</i> Batko				a								
<i>Pythiogeton ramosum</i> Minden							s				s	
<i>Pythiogeton uniforme</i> Lund				a			s	s		s		
<i>Pythium afertile</i> Kanouse et Humphrey				w								
<i>Pythium aquatile</i> Höhnk								s				
<i>Pythium artotrogus</i> Sideris				w		w						
<i>Pythium butleri</i> Subram.				a,w		a,w				a	a	
<i>Pythium carolinianum</i> Matthews				w								
<i>Pythium catenulatum</i> Matthews				w			s					
<i>Pythium debaryanum</i> Hesse				a,w		w					a	
<i>Pythium dissotocum</i> Drechsler				w		a,w						
<i>Pythium echinulatum</i> Matthews						a						
<i>Pythium helicandrum</i> Drechsler				w								
<i>Pythium imperfectum</i> Höhnk				a,w		a,w						
<i>Pythium inflatum</i> Matthews						s,w		s		s		
<i>Pythium intermedium</i> de Bary				w								
<i>Pythium multisporum</i> Poitras								s			s	
<i>Pythium myriotylum</i> Drechsler								s	s			
<i>Pythium palingenes</i> Drechsler	s		s		s							s
<i>Pythium perniciosum</i> A.A. Yachevskij et P.A. Yachevskij						w						
<i>Pythium proliferum</i> Schenk		s	s	w	s		s	s			s	
<i>Pythium rostratum</i> Butler				s,w		a,w	s			s	a	
<i>Pythium tenue</i> Gobi				w								
<i>Pythium torulosum</i> Coker et Patterson	s			w								

Table 3 continues on next page...

<i>Pythium undulatum</i> Petersen				w								
<i>Pythium vanterpoolii</i> V. Kouyeas et H. Kouyeas						w						
Zygomycetes												
Zoopagales												
<i>Zoophagus insidians</i> Sommerstorff				a,w		s,a,w	s			a		
Total number	18	12	13	69	14	66	13	15	16	20	18	12

Table 4. Hyphomycetes species found in water of particular sites of Biebrza National Park (a-autumn, s-spring).

Species of fungi	Site											
	I	II	III	IV	V	VI	VII	VIII	IX	X	XI	XI
<i>Acrodictys bambusicola</i> M. B. Ellis	a	a	a	s,a	a	a	a	a	s,a	a	s,a	a
<i>Acrodictys elaeidicola</i> M. B. Ellis	a			a	a	s				a		
<i>Acrodictys martinii</i> Crane et Dumont		s		a	s,a	a				a		
<i>Alatospora acuminata</i> Ing.					s						a	
<i>Anguillospora longissima</i> (Sacc. et Syd.) Ing.	s,a	s	s,a	s,a	a	s,a	s,a		s,a		s	a
<i>Angulospora aquatica</i> S. Nilss.	s,a	s,a	s,a	s,a	s,a	s,a	s,a	s,a	s,a	s,a	s,a	s,a
<i>Arborispora palma</i> Ando	a											s
<i>Arbusculina fragmentans</i> Marvanová et Marvan	a	s,a			s	a				a	a	s
<i>Arthrobotrys silbacea</i> Fres.					a		a					
<i>Bacillispora aquatica</i> S. Nilss.				s								
<i>Beverwykella pulmonaria</i> (v. Beverw.) Tubaki						s				a		
<i>Bidenticula cannae</i> Deighton				s	s							
<i>Blongettia indica</i> Subramanian				a	a	s		a	a	a	a	
<i>Calcarispora hiemalis</i> Marvanová et Marvan	s,a		a	s	s	s			a	a	s	
<i>Canalisporium caribense</i> (Hol.-Jech. et Merc.) Naw. et Kath.				a	s			a	s,a	a	a	
<i>Centrospora aquatica</i> Iqbal	a	a		s,a	a		s,a		a			s
<i>Centrospora filiformis</i> (Greath.) Petersen	s,a	s	s,a	s	s,a	s	s	s,a	s,a	s,a	s,a	s,a
<i>Cladosporium peruamazonicum</i> Matsushima							a					
<i>Clavatospora longibrachiata</i> (Ing.) S. Nilss.	s											
<i>Clavatospora tentacula</i> (Umph.) S. Nills.												a
<i>Colispora elongata</i> Marvanová							a					
<i>Condylospora gigantea</i> Nawawi et Kuth.						s						
<i>Condylospora spumigena</i> Nawawi		s,a				s		s				s
<i>Corynespora cubensis</i> Hol.-Jech.			s,a	s	a		a					
<i>Cylindrocarpon aequatoriale</i> Matsushima							s					
<i>Cylindrocarpon aquaticum</i> Matsushima												s
<i>Dactylaria candidula</i> (v. Höhnelt) Bhatt et Kendrick												a
<i>Dactylaria gracilis</i> Duddington									a			

Table 4 continues on next page...

<i>Dactylaria inaequilatera</i> Matsushima	a												
<i>Dactylella submersa</i> (Ing.) S. Nilss.	a		a					a	s			a	
<i>Didymobotrym verrucosum</i> Hino et Katum				a									
<i>Filosporella exilis</i> Gulis et Marvanová		s						s	s				
<i>Flagellospora stricta</i> S. Nilss.	s												
<i>Gyoeffyyella myrmecophagiformis</i> Mielnik et Dudka									s	s			
<i>Heliscus lugdunensis</i> Sacc. et Therry	s	a						a			a	a	a
<i>Heliscus submersus</i> Hudson	s,a	s	s,a		s,a	s,a	a	s,a	a		s	s,a	
<i>Helminthosporium bigenum</i> Matsushima								s		a	a		
<i>Hyalobelemnospora amazonica</i> Matsushima								a					
<i>Ingoldiella hamata</i> Shaw										s			
<i>Kontospora halophila</i> Roldan et Honr.	s	s,a	s	s	a	s,a	s,a	s,a	s,a	a			
<i>Kylindria keitae</i> Ramb. et Onofri					a					s		s	
<i>Kylindria peruamazonensis</i> Matsushima											a		
<i>Lemonniera aquatica</i> de Wild.	s,a	s	s		s,a	s,a	s,a	a		a		s,a	
<i>Lemonniera filiforme</i> Petersen			a										
<i>Leuliisinea amazonensis</i> Matsushima			s			a							
<i>Lunulospora curvula</i> Ing.	s,a	s,a	s,a			a	s,a	s,a	a	a	a	s,a	a
<i>Margaritipora aquatica</i> Ing.			a										
<i>Melanocephala manuensis</i> Matsushima											a		
<i>Malanocephala</i> sp.									a				
<i>Mirandina corticola</i> G. Arnaud	s,a	s,a	s	s,a	s,a	s	a	s	a	a	a	a	a
<i>Monodictys peruviana</i> Matsushima		a				a		a	s	a	s		
<i>Monodisma fragilis</i> Alcorn										a			
<i>Paraarthrocladium amazonense</i> Matsushima				s,a	s		s	s				s	
<i>Paracryptophiale kamarudolinii</i> Kuth. et Nawawi					s					s			
<i>Paradactylella peruviana</i> Matsushima	s												
<i>Paraepicoccum amazonense</i> Matsushima								a					
<i>Phaedactylum acutisporum</i> Matsushima	s												
<i>Phialogeniculata multiseptata</i> Matsushima							s					s,a	
<i>Piricauda cubensis</i> Hol.-Jech. et Merc.												a	
<i>Pithomyces obscuriseptatus</i> Matsushima												s	
<i>Pleiochaeta amazonensis</i> Matsushima													a
<i>Polycladium equiseti</i> Ing.			a	a									
<i>Polystratorictus fusarioideus</i> Matsushima	a					a	s	a	a				
<i>Pseudaegerita corticalis</i> (Peck.) Crane et Schok.			a			a							
<i>Pseudoanguillospora gracilis</i> Sincl. et Morgan-Jones		s		s									
<i>Pseudohansfordia dimorpha</i> Matsushima				a									
<i>Pseudospiropes lotorus</i> Morgan-Jones								s,a					s,a
<i>Ramichloridium clavulisporum</i> Matsushima								a					

Table 4 continues on next page...

<i>Scolecobasidium lanceolatum</i> Matsushima	s	s											a	
<i>Sigmoidea prolifera</i> (Peter.) Crane	a	s,a			s,a			s	s					
<i>Sporidesmium acutifusiforme</i> Matsushima								a	a	a			s	
<i>Sporidesmium inflatum</i> (Berk. et Br.) M.B. Ellis						s								
<i>Sporidesmium moniliforme</i> Matsushima				a	a			a	s					
<i>Stachybotrys theobromae</i> Hansf.								a			a			
<i>Stagonospora macropycnidia</i> Cunnell								a				s		
<i>Sterigmatobotrys uniseptata</i> Chang												a	a	
<i>Tetracladium marchalianum</i> de Wild.	a	s	s,a	s,a	s,a	a	s,a	a	s	a	s,a	s,a		
<i>Tetracladium maxilliformis</i> (Rostr.) Ing.					s									
<i>Titaea clarkeae</i> Ellis et Everh.													s	
<i>Tricellula aquatica</i> Webster	s,a	s		s		s		a	s				s	
<i>Tricladium angulatum</i> Ing.		a				s								
<i>Tricladium attenuatum</i> Iqbal	s			s				s				s		
<i>Tricladium marylandicum</i> Crane		a			a									
<i>Tricladium patulum</i> Marvanová et Marvan	s							s					s	
<i>Trinacrium subtile</i> Riess ap. Fres.		s						s			s	s,a		
<i>Tripospermum camelopardus</i> Ing. et al.					a		s				s	s		
<i>Tripospermum prolongatum</i> Sincl. et Morgan-Jones										a				
<i>Triscelophorus monosporus</i> Ing.	a			s		a							s	
<i>Varicosporium delicatum</i> Iqbal	a			a			a				a			
<i>Veronea bothryosa</i> Cif. et Montem.						a								
<i>Volucrispora graminea</i> Ing. et al.								a					a	
Total number	30: s-18 a-21	25: s-19 a-13	18: s-11 a-14	26: s-17 a-15	29: s-17 a-20	28: s-17 a-16	27: s-14 a-20	30: s-15 a-21	27: s-15 a-17	22: s-0 a-22	27: s-18 a-16	28: s-17 a-18		

of zoosporic fungi new to Polish waters contains *Phlyctochytrium parasitans*, *Rhizophyidium coronum*, *Rhizidium richmondense*, *Phytophthora cactorum*, *Endochytrium ramosum*, *Karlingia hyalina*, *Saprolegnia semihypogyna* and *Lagenidium destruens*. The Hyphomycetes, also new to Polish waters, include *Bidentacula cannae*, *Dactylaria gracilis*, *Dactylaria candidula*, *Filosporella exilis*, *Hyalobeleemospora amazonica*, *Monodisma fragilis*, *Pseudoanguillospora gracilis* and *Ramichloridium clavuliformis*.

The cluster analysis of the investigate parameters carried out in water of the River Biebrza during investigations of fungus species has revealed that in spring and autumn is different (Fig. 2,3).

Discussion

The present study has revealed that two sites - the River Biebrza in Osowiec (site IV) and Kanał Rudzki (site VI) are the most abundant in zoosporic fungus species. It

does not indicate, however, that there exist the optimum environmental conditions, but may be due to the fact that water samples from these sites were additionally collected. The most common zoosporic fungi at most sites were *Aphanomyces irregularis*, *Aphanomyces laevis*, *Catenophlyctis variabilis* and *Leptolegniella keratinophilla*. These species have been quite frequently encountered in our studies on keratin-containing substrates in different limnological types. The conidial fungus species common in that region and found in the water at all sites included *Acrodictis bambusicola*, *Angulospora aquatica*, *Centrospora filiformis*, *Mirandina corticola* and *Tetracladium marchalianum*. They are commonly encountered in our waters, the first two can be even found during heavy cyanobacterial blooming [24].

Most fungus species revealed in the water of Biebrza National Park had already been found in other rivers of northeastern Poland [25-39].

Eight zoosporic species and nine conidial species found in the present study are new to Polish waters. *Endochy-*




Fig. 1. Some zoosporic (A-D) and conidial (E,F) fungus species from river Biebrza (x 200).

A - *Achlya klebsiana*- oogonia (25-85 μm) with oospores (15-50 μm); B - *Aphanomyces stellatus* -hyphae from oogonia (25-30 μm); C - *Pythium echinulatum* - sporangia (10-30 μm); D - *Pythium vanterpoolii* - sporangia; E - *Paradactylella peruviana* - conidium (37-67 x 3.5-5.5 μm); F - *Pithomyces obscuriseptatus* - conidium (16-22.5 x 11-15.5 μm)

trium ramosum, a zoosporic species, was found to grow in the water collected at site I Dolistowo. This fungus was first described by Sparrow [40] in North America from dead aquatic green algae. *Karlingia hyalina*, known as an aquatic plant saprophyte, was first described by Karling [41] from Brazilian waters. We observed this fungus at site VI Kanał Rudzki in the winter.

Phlyctochytrium parasitans was found to grow on *Achlya dubia* specimens in water samples collected at site VI Kanał Rudzki in December. It was first described by Sparrow and Dogma [42] from soil samples in the Dominican Republic, where it was isolated from the oogonia of *Achlya flagellata*. This would be the first report on the occurrence of *Phlyctochytrium parasitans* in the aquatic environment. The growth of *Phlyctochytrium reinboldtae* was noted in December in the River Biebrza at the site by the road, being the first aquatic site of that fungus. It was isolated from soil samples by Persiel [43] and later also from soil [44-45]. *Rhizidium richmondense*, another new species to Polish waters, was found in December on cellophane at site VI Kanał Rudzki. This fungus was first described by

Willoughby [46] also from soil, with cellophane as bait. It also grows well on onion skin [47]. *Rhizophyidium ampullaceum*, never before encountered in Polish waters, was found in site VI Kanał Rudzki in December. In literature it is described as a parasite of filiform green algae [12]. *Rhizophyidium coronum* is a plant saprophyte first described from grass leaves [48]. In Biebrza National Park we found this fungus in site VI Kanał Rudzki. *Saprolegnia semihypogyna*, another species new to Polish waters, was found in September 2002 at site IV Osowiec (in bottom of shore). This fungus was first described by Inaba and Tokumasu [49] from soil in Japan. *Lagenidium destruens* is a parasite of other aquatic fungi, particularly those belonging to the genus *Achlya*, and was first described by Sparrow [7]. In the River Biebrza valley it was observed at site IX Olszowe Pole - the old river-bed. And finally, *Phytophthora cactorum*, also new to Polish waters, was found in October at site VI Kanał Rudzki.

Among the conidial fungus species which appear new to Polish waters *Dactylaria gracilis* and *Dactylaria candidula* are predacious species catching nematodes.




Fig. 2. Clustering of river Biebrza according of water chemistry data and to number fungus species (autumn 1999).




Fig. 3. Clustering of river Biebrza according of water chemistry data and to number fungus species (spring 2000).

The former was first reported from soil by Duddington [50], the latter by Bhatt and Kendrick [51]. In the present study they were both found in the autumn; *Dactylaria gracilis* at site VI Kanał Rudzki and *Dactylaria candidula* at site XII Burzyn. *Bidentacula cannae* was found to grow only in the spring months in the River Biebrza in Osowiec (site IV) and in the water of moat Osowiec (site V). It was observed in land conditions in Africa by Deighton [52] and reported from Rio Itaya, an Amazon tributary [22]. *Filosporella exilis* was first described from the fragments of horsetail *Equisetum fluviatilis* submerged in the water of Busianka stream in Belarus [53]. We found it in the old river-bed closed in the vicinity of Dolistowo (site II) and site IX Olszowe Pole (the old river-bed connected with the River Biebrza), and in the site VII River Biebrza at the site of Olszowe Pole. Species new to Polish waters, such as *Hyalobelemnospora amazonica* and *Ramichloridium clavulisorum* were first described and known from the Amazon Basin. In the present study *Hyalobelemnospora amazonica* was found to grow in the spring and autumn at site VII (Olszowe Pole), while the other species at the site also in the autumn. It was described by Alcorn [54] and encountered in the Amazon tributaries [22]. And finally, *Pseudoanguillospora gracilis* was reported from the State of Alabama in the United States from *Platanus occidentalis* leaves fallen to water [55]. We found this fungus in site IV River Biebrza at the site of Osowiec in the spring and in the old river-bed closed at the site I Dolistowo.

Like in the cases of the National Park [56] and Narew National Park [57], the waters of Biebrza National Park provide a favourable habitat for numerous groups of conidial fungus species belonging to the so called "Amazon fungi", first described at the beginning of the 1990s in South America in the Amazon Basin by Japanese mycologist Matsushima [22]. Twenty three conidial species found in the present study belong to this group. This is a relatively large group of conidial fungus species of which

some inhabit springs in Knyszyn Forest [58] or certain lakes in the Suwałki District [38]. This would suggest that Podlasie waters still create favourable conditions for conidial fungus species variety.

There are no geographic barriers for aquatic fungi and therefore the same species can be encountered under different climatic conditions. The group of conidial fungus species described from the Amazon and its tributaries, and recently detected in the waters of northeastern Poland seems to confirm this phenomenon. Winds, air currents and migratory birds are all involved in the intercontinental dissemination of aquatic fungi. Physicochemical factors of a particular water reservoir determine the occurrence of a particular species in the aquatic environment. With a few exceptions, both zoosporic and conidial aero-aquatic and aquatic fungi belong to the group of cosmopolitan organisms [59,60].

The cluster analysis of the respective parameters carried out in water of river Biebrza during study on fungi species has revealed that in spring the number of fungi species is associated with hardness in Mg of water and in autumn with temperature of water. According to literature data an increase in water hardness in certain water-courses leads to a decrease in the number of zoosporic fungi [61] and conidial species [62]. It is generally accepted that the zoosporic and conidial fungi prefer low water temperature [32, 57, 63].

Acknowledgements

Thanks are due J. Kurkus and M. Orłowska for help in the field and in the laboratory.

References

1. CZECZUGA B., WORONOWICZ L., BRZOZOWSKA K. Aquatic fungi of the lowland river Biebrza. *Acta Mycol.* **26**, 77, 1990.
2. CZECZUGA B., ORŁOWSKA M., MAZALSKA B. Wodne grzyby niższe Biebrzańskiego Parku Narodowego [Lower

- aquatic fungi of the Biebrza National Park]. W: Czeczuga B., Rybak I.(eds.) Materiały XVIII Zjazdu Hydrobiologów Polskich w Białymstoku, 4-8 września 2000, p. 45, **2000**. (In Polish).
3. CZECZUGA B. Rola grzybów niższych w zbiornikach wodnych [Function of lower aquatic fungi in water bodies]. W: Czeczuga B., Rybak I.(eds.) Materiały XVIII Zjazdu Hydrobiologów Polskich w Białymstoku, 4-8 września 2000, pp. 7-11, **2000**. (In Polish).
 4. CZECZUGA B. Rzadkie gatunki grzybów konidialnych (Hyphomycetes) w wodach Biebrzańskiego Parku Narodowego [Some rare species Hyphomycetes in north-eastern Poland]. Reg. Stud. **2**, 55, **2001**. (In Polish).
 5. GREENBERG A.E., CLESCERI L.S., EATON A.D. Standard Methods for the Examination of Water and Wastewater. American Public Health Association, Washington, pp. 1193, **1992**.
 6. JOHNSON T.W. The genus *Achlya*. Morphology and Taxonomy. Ann Arbor, Michigan, pp. 180, **1956**.
 7. SPARROW F.K. Aquatic Phycomycetes. Ann Arbor, Michigan, pp. 1187, **1960**.
 8. WATERHOUSE G.M. Key to *Pythium* Pringsheim. Mycol. Pap. **109**, 1, **1967**.
 9. WATERHOUSE G.M. The genus *Pythium* Pringsheim. Mycol. Pap. **110**, 1, **1968**.
 10. SEYMOUR R.L. The genus *Saprolegnia*. Nova Hedwigia **19**, 1, **1970**.
 11. BEDENEK T. Fragmenta Mycologica. I. Some historical remarks of the development of „hairbaiting” of Tomarling- Vanbreuseghem (the To-Ka-Vahairbaiting method). Mycopathol. Appl. **68**, 104, **1972**.
 12. BATKO A. Zarys hydromikologii [Hydromycology - an overview]. PWN, Warszawa, pp. 478, **1975**.
 13. KARLING J.S. Chytridiomycetorum Iconographia. J. Cramer, Vaduz, pp. 414, **1977**.
 14. DICK M.W. Keys to *Pythium*. Reading, U.K., College Estate Management, Whiteknights, pp. 66, **1990**.
 15. PYSTINA K.A. Genus *Pythium* Pringsh. Nauka, Sankt Petersburg, pp. 126, **1998**.
 16. NILSSON S. Freshwater Hyphomycetes. Taxonomy, morphology and ecology. Symb. Bot. Uppsal. **18**, 1, **1964**.
 17. DUDKA I.O. Wodni hifomiceti Ukraini [Aquatic Hyphomycetes of Ukraine]. Naukova Dumka, Kiev, pp. 239, **1974**. (In Ukrainian).
 18. DUDKA I.A. Vodnyje nesovershennyje griby SSSR [Aquatic Fungi imperfecti from the Soviet Union]. Naukova Dumka, Kiev, pp. 187, **1985**. (In Russian).
 19. INGOLD C.T. An illustrated guide to aquatic and water born Hyphomycetes (Fungi Imperfecti) with notes on the their biology. Fresh. Biolog. Assoc. Sci. Publ. **30**, 1, **1975**.
 20. CARMICHAEL J.W., KENDRICK W.B., CONNERS I.L., SIGLER L. Genera of Hyphomycetes. Alberta Press: Edmonton, pp. 386, **1980**.
 21. BRÄTHEN I. The aquatic stauroconidial Hyphomycetes of Norway with notes on the Nordic species. Nord. J. Bot. **4**, 375, **1984**.
 22. T. MATSUSHIMA Mycological Memoirs No 7. Published by the author, Kobe, pp. 75, **1993**.
 23. PODANI J. Introduction to the exploration of multivariate biological data. Backhuys Publishers BV, Leiden, pp. 407, **2000**.
 24. CZECZUGA B., ORŁOWSKA M. Investigations on the joint occurrence of *Anabaena spiroides* Klebahn and Hyphomycetes in various types of water bodies. Acta hydrochim. hydrobiol. **28**, 162, **2000**.
 25. CZECZUGA B. Grzyby niższe w wodzie rzeki Narwi i jej dopływach na odcinku od Suraża do Tykocina [Lower fungi in water of the river Narew and its tributaries in the stretch from Suraż to Tykocin. W: Banaszuk H., Czeczuga B. (eds.) Narwiański Park Krajobrazowy i okolice. Zagad. Przyrodn. Gospod. OBN, Białystok: 129, **1990**. (In Polish).
 26. CZECZUGA B. Aquatic fungi of the river Pisa and its tributary the river Skroda. Acta hydrochim. hydrobiol. **19**, 57, **1991**.
 27. CZECZUGA B. Mycoflora of the river Węgorapa and its tributary, the river Gołdapa-Jarka. Acta hydrochim. hydrobiol. **19**, 517, **1991**.
 28. CZECZUGA B. Mycoflora of the Narew River and its tributaries and the stretch between Tykocin and Ostrołęka. Acta Mycol. **30**, 181, **1995**.
 29. CZECZUGA B. Mycoflora of the Supraśl river and its tributaries. Acta Mycol. **31**, 13, **1996**.
 30. CZECZUGA B. Aquatic fungi of the Narew river and its tributaries in the stretch from Siemianówka to Doktorce. Ann. Acad. Med. Bialostocensis **41**, 245, **1996**.
 31. CZECZUGA B., MUSZYŃSKA E. Aquatic fungi of the Rudawka river. Ann. Med. Univer. Białystok **38**, 7, **1993**.
 32. CZECZUGA B., SNARSKA A. *Pythium* species in 13 various types of water bodies of N-E Poland. Acta Soc. Bot. Pol. **70**, 61, **2001**.
 33. CZECZUGA B., PRÓBA D., BRZOZOWSKA K. Grzyby wodne rzeki Narwi na odcinku Suraż-Tykocin oraz w ujściu rzeki Turoślanki i Supraśli na tle zróżnicowanego środowiska [Water fungi of the river Narew in the stretch from Suraż to Tykocin and the mouths of the rivers Turoślanka and Supraśl with reference to differences in the environment]. Ann. Acad. Med. Bialostocensis **29**, 77, **1984**. (In Polish).
 34. CZECZUGA B., WORONOWICZ L., BRZOZOWSKA K. Grzyby wodne rzeki Białej na stanowiskach o różnym stopniu i charakterze zanieczyszczeń [Water fungi of the river Biała at sites on differences and quality of pollutions]. Ann. Acad. Med. Bialostocensis **31**, 49, **1986**. (In Polish).
 35. CZECZUGA B., WORONOWICZ L., BRZOZOWSKA K. Aquatic fungi of two forest brooks. Nova Hedwigia **43**, 459, **1986**.
 36. CZECZUGA B., BRZOZOWSKA K., WORONOWICZ L. Mycoflora of the upper course of the river Szeszupa. Ann. Acad. Med. Bialostocensis **33**, 115, **1989**.
 37. CZECZUGA B., BRZOZOWSKA K., WORONOWICZ L. Mycoflora of the river Czarna Hańcza and its tributary - the river Marycha. Int. Revue ges. Hydrobiol. **75**, 245, **1990**.
 38. CZECZUGA B., ORŁOWSKA M., WORONOWICZ L. Fungi in water reservoirs of the Western Suwałki Lake District and in some adjacent water basins. Ann. Acad. Med. Bialostocensis **42**, 43, **1997**.
 39. CZECZUGA B., KIZIEWICZ B., ORŁOWSKA M. Zoosporic and conidial fungi within the Podlasie stretch of the River Bug. Ann. Acad. Med. Bialostocensis **47**, 40, **2002**.
 40. SPARROW F.K. Observations on operculate chytridiaceous fungi collected in the vicinity of Ithaca, New York. Amer. J. Bot. **20**, 63, **1933**.
 41. KARLING J.S. Brazilian chytrids. X. New species with sunken opercula. Mycologia **39**, 56, **1947**.
 42. SPARROW F.K., DOGMA I.J. Zoosporic Phycomycetes from Hispaniola. Arch. Mikrobiol. **89**, 177, **1973**.
 43. PERSIEL I. Über *Phlyctochytrium reinboldtae* n.sp. Arch. Mikrobiol. **32**, 411, **1959**.
 44. BARR D.J.D. *Phlyctochytrium reinboldtae* (Chytridiales): morphology and physiology. Can. J. Bot. **48**, 479, **1970**.
 45. WILLOUGHBY L.G., RIGG E. Lower aquatic fungi in the soils of Majorka. Nova Hedwigia. **37**, 369, **1983**.

46. WILLOUGHBY L.G. Studies on soil chytrids. I. *Rhizidium richmondense* sp. nov. and its parasites. Trans. Brit. Mycol. Soc. **39**, 125, **1956**.
47. KARLING J.S. Some zoosporic fungi of New Zealand. IV. Polyphlictis gen. nov., *Phlyctochytrium* and *Rhizidium*. Sydowia **20**, 86, **1967**.
48. HANSON A.M. A morphological, developmental and cytological study of four saprophytic chytrids. II. *Rhizophydium coronum* Hanson. Amer. J. Bot. **32**, 479, **1945**.
49. INABA S., TOKUMASU S. *Saprolegnia semihypogyna* sp. nov., a saprolegniaceous oomycete isolated from soil in Japan. Mycoscience **43**, 73, **2002**.
50. DUDDINGTON C.L. Further records of predacious fungi. II. Trans. Brit. Mycol. Soc. **34**, 194, **1951**.
51. BHATT G.C., KENDRICK W.B. The generic concept of *Diplorhinotrichum* and *Dactylaria*, and anew species of *Dactylaria* from soil. Can. J. Bot. **46**, 1253, **1968**.
52. DEIGHTON F.C. Four leaf-spotting Hyphomycetes from Africa. Trans. Brit. Mycol. Soc. **59**, 419, **1972**.
53. GULIS V.I., MARVANOVÁ L. *Filosporella exilis* sp. nov. on submerged plant debris from Belarus. Mycotaxon **68**, 313, **1998**.
54. ALCORN J.L. *Monodisma*, a new genus of Hyphomycetes. Trans. Brit. Mycol. Soc. **65**, 139, **1975**.
55. SINCLAIR R.C., MORGAN-JONES G. Notes on Hyphomycetes. XXII. Five new aquatic species. Mycotaxon **9**, 469, **1979**.
56. CZECZUGA B., ORŁOWSKA M., KOZŁOWSKA M. Hyphomycetes of lakes of varied trophic state in the Wigry National Park (North-Eastern Poland). Pol. J. Ecol. **49**, 201, **2001**.
57. CZECZUGA B., KIZIEWICZ B., GODLEWSKA A., ORŁOWSKA M. Further studies on aquatic fungi in the River Narew within the Narew National Park. Ann. Acad. Med. Bialostocensis **47**, 58, **2002**.
58. CZECZUGA B., ORŁOWSKA M. Hyphomycetes in twenty springs of the Knyszyn-Białystok Forest in various seasons. Int. Revue ges. Hydrobiol. **81**, 417, **1996**.
59. WOOD-EGGENSCHWILER S., BÄRLOCHER F. Geographical distribution of Ingoldian fungi. Verh. Inter. Vercin. Limnol. **22**, 2780, **1985**.
60. CZECZUGA B., KIZIEWICZ B. Dissemination on fungi in the aquatic environment. Ecol. Quest. **2**, in press, **2003**.
61. CZECZUGA B., PRÓBA D. Mycoflora of the upper part of the river Narew and its tributaries in a differentiated environment. Nova Hedwigia **44**, 151, **1987**.
62. BÄRLOCHER F. The Ecology of Aquatic Hyphomycetes. Springer-Verlag, Berlin et al., pp. 225, **1992**.
63. CZECZUGA B., ORŁOWSKA M. Hyphomycetes in the river Supraśl in various seasons of the year with reference to environmental conditions. Int. Revue ges. Hydrobiol. **78**, 611, **1993**.

Learning FEATURES you get in EVERY Self-Study Course!

OBJECTIVES

At the beginning of each lesson, you'll find a list of clearly defined learning objectives which gives you a complete overview of what will be covered and what you can expect to learn. These objectives steer you towards your goal of greater knowledge and proficiency in the subject area.

EXERCISES

Section-by-section exercises and review questions are featured in each lesson to increase your retention and help you apply the concepts to your own situation. Many sections are also footnoted, providing you with additional references and resources to answer other questions you may have.

SELF-TESTS

Concluding each lesson is a short multiple-choice self-test that covers the key points you should retain from your learning. Answers are provided separately so you can continually check your progress. Should you have any questions about the course content, your instructor is always available by e-mail or fax.

MASTER EXAM

A comprehensive, challenging Master Exam at the end of the course measures your knowledge of every aspect of the materials covered. You'll feel confident knowing that you've received a thorough knowledge base to help you achieve regulatory compliance, improve your management systems, and meet your professional development goals.