

Original Research

Remediating Cd-Contaminated Soils Using Natural and Chitosan-Introduced Zeolite, Bentonite, and Activated Carbon

Na Yi, Yaoguo Wu*, Lin Fan, Sihai Hu

School of Natural and Applied Sciences, Northwestern Polytechnical University, Xi'an, Shaanxi, China

Received: 28 December 2017

Accepted: 27 March 2018

Abstract

The effects of in-situ immobilization of heavy metals by applying natural and chitosan-introduced zeolite, bentonite, and activated carbon (AC) were systematically studied to remediate cadmium (Cd)-contaminated soils in a pot experiment using *Brassica juncea* as the indicator plant. The results show that zeolite, bentonite, and its chitosan composites can increase soil pH and reduce the biological effectiveness of heavy metals. The *Brassica juncea* dry weight increased with increasing of amendment dosage. Highest values were found for CS-AC, followed by CS-bentonite, CS-zeolite, AC, bentonite, and zeolite. With an amendment dosage of 75 g per pot, *Brassica juncea* dry weight increased by 41.91%, 39.00%, 27.64%, 35.93%, 23.78%, and 23.58%, respectively, for CS-AC, CS-bentonite, CS-zeolite, AC, bentonite, and zeolite, compared to the control. Cadmium uptake by *Brassica juncea* was lowest for this dosage. With a dosage of 75 g, 50 g, 75 g, 75 g, 50 g, and 75 g per pot for CS-AC, CS-bentonite, CS-zeolite, AC, bentonite, and zeolite, respectively, Cd uptake decreased by 21.89%, 19.88%, 19.48%, 18.67%, 17.47%, and 13.85%, respectively. Similarly, bioavailable Cd content decreased by 27.38%, 19.29%, 22.83%, 23.22%, 15.74%, and 8.66%, respectively, compared to the control.

Keywords: Cd contamination, remediation, chitosan, *Brassica juncea*, amendments

Introduction

The heavy metal cadmium (Cd) has long been recognized as a more hazardous pollutant for plants, animals, and humans than any other heavy metal [1-2]. Soil contamination with toxic heavy metals had been a worldwide challenge for food security and health as heavy metals do not degrade, adversely affecting the biota via the food chain [3-6]; therefore,

the accumulation of heavy metals in soil can result in a decrease of soil fertility and soil microbial activities and biodiversity, is collectively responsible for crop losses, and has even more serious implications for animal and human health [7-8]. Studies estimate that the area of soil contaminated with Cd covers about 1.3×10^5 ha and accounts for 1/5 of the total farmland area in China [9]. And more than 11 provinces and 25 districts are subjected to Cd-contaminated soil. In 2007, the direct economic loss from farmland metal pollution reached \$3.2 billion as calculated by the Ministry of Land and Resource in China [10]. Therefore, the issue of scientific and economic mitigation of Cd-contaminated

*e-mail: wuygal@nwpu.edu.cn

soils requires urgent clarification. Many remediation techniques have been applied for the removal of heavy metals from soils. Chemical methods of soil remediation may destroy soil structure and result in secondary pollution [11]. Therefore, more economic, effective, and environmentally friendly metal chelators are required. The enhancement of chemical immobilization by the addition of soil amendments is increasingly seen as a valuable alternative strategy that can effectively reduce the bioavailability of contaminants [12-13]. Two mechanisms are responsible for the efficiency of this remediation action: 1) the increase in metal sorption of the resulting soil+material mixture and 2) dilution of the contaminant concentration when large material doses are used [14].

A potential "green" metal chelator is chitosan (CS). It is the second most abundant biopolymer in the natural environment and is obtained from the deacetylation of chitin extracted from shellfish processing waste [15-16]. At present, CS is receiving considerable attention as an excellent natural adsorbent to remove numbers of heavy metal ions due to the presence of amine (-NH₂) and amounts of reactive hydroxyl (-OH) function groups on the main chain, which act as chelation sites for metal ions [15, 17-18]. The largely spread hydroxyl and amine groups serve as effective chelating sites for heavy metals through ion exchange, adsorption, and complexation. Of these, complexation is the major, stable, and irreversible form [19]. Although the adsorption of chitosan on Cd²⁺ and other heavy metal ions has been widely studied [20-21], the studies of chitosan on Cd²⁺-contaminated soils are too rare.

Clay minerals have larger interior and exterior-specific surface areas, many reactive functional groups, and the hydration and dehydration of interlaminal surface [22]. The high metal adsorption capacity and low cost of clay minerals makes them one of the most common remediation options for metal adsorption [23-25]. They therefore have an ability to bind heavy metals, preventing their transportation from the soil into the plant. Zeolite is an important clay mineral and has been used in numerous studies on Cd-polluted soils [26-28]. Similarly, bentonite is a type of expandable clay composed primarily of montmorillonite and characterized by high permanent negative charges and a large surface area [29]; it is also an efficient adsorbent for Cd [28, 30]. Activated carbon (AC) with large surface area has been used for a long time as an efficient amendment to remediate Cd-contaminated soil; the capability of AC to bind Cd has been confirmed by Břendová et al. [31].

Cruciferae canola is a plant with strong absorption characteristics and heavy metal enrichment ability [32]. *Brassic juncea* is a variety of cruciferae canola. *Brassica juncea* was selected as the experimental species because of its short life stage, high biomass, and the hyperaccumulation ability of heavy metal [33]. And *Brassica juncea*, a widely consumed species,

is very sensitive to the presence of toxic elements in soil. In addition, the great commercial value of *Brassica juncea* makes it necessary to protect it from Cd contamination. It has great potential applications in phytoremediation. Considering ecology, toxicology, and health, it is particularly important to study the contents of effective heavy metals in contaminated soil, because these components are easily magnified through biont. In order to reduce the amount of heavy metals in plants, it will be an effective way to improve the contaminated soil by heavy metals, and adding amendments into the soil reduces the effectiveness of heavy metals. In recent years, extensive research have been conducted on adjusting the soil parameters to reduce the effectiveness of heavy metal through adding a hardening agent (activated carbon, zeolite, bentonite, etc.) into the contaminated soil by heavy metal. The method of heavy metal pollution treatment can be widely applied due to the advantages of economically viable and non-destructive soil structures [34]. In addition to reducing the dissolubility of heavy metals, the amendments can effectively reduce the bioavailability of heavy metals and the negative effects on health [35]. The objective of this study was to evaluate the efficiency of Cd-contaminated soil by natural and chitosan-introduced zeolite, bentonite, and AC in the remediation of Cd-contaminated soils. We determined the effects of these substances on soil pH, dry weight of individual *Brassica juncea*, Cd uptake of *Brassica juncea*, and bioavailable Cd.

Experimental Section

Soil and Amendments

Soil (0-20 cm depth) was collected from a greenhouse in Yangling, Shaanxi Province, China. Phosphate was applied to the soil over a period of 30 years. The soil was air-dried, thoroughly mixed, and passed through a 20-mm mesh sieve to remove gravel. The soil was classified as loess soil, and some of the physical and chemical properties were listed in Table 1. The pH was measured with a pH meter (FE28-Meter, Mettler Toledo instrument co., LTD, China) in a 1:5 (w/w) soil-H₂O slurry. Soil organic matter content was determined using the potassium dichromate-outside heating method. Bioavailable Cd was extracted for 2 h at room temperature with DTPA (DTPA-TEA-CaCl₂, pH = 7.3) buffer solution. Heavy metal content was determined using a flame atomic absorption spectrophotometer (Z-2000, Hitachi, Japan.).

Zeolite, bentonite, and activated carbon were obtained from the Zhongchuan mineral powder processing plant, Lingshou, Hebei province, China. Industrial-grade chitosan was obtained from Guangzhou Shengxun chemical industry co. LTD, and it had the following properties: yellowish-white powder with 100 meshes, 93% degree of deacetylation, and a molecular weight of 2*10⁶.

Table 1. Physiochemical properties of the 0-20 cm soil layer in the experimental area. Values represent mean \pm standard deviation (n = 3).

Parameter	Unit	Value
Soil type		Heavy loam
pH		7.81 \pm 0.10
Total nitrogen	g kg ⁻¹	1.69 \pm 0.05
Olsen phosphate	mg kg ⁻¹	303.6 \pm 8.47
Available potassium	mg kg ⁻¹	722.7 \pm 14.12
Soil organic matter	g kg ⁻¹	36.15 \pm 1.10
Cd	mg kg ⁻¹	1.55 \pm 0.04
Available Cd	mg kg ⁻¹	0.29 \pm 0.03
Fe	%	3.11 \pm 0.21
Cu	mg kg ⁻¹	24.6 \pm 1.98
Zn	mg kg ⁻¹	69.5 \pm 3.48
Ca	%	4.79 \pm 0.22
Mg	%	1.28 \pm 0.02

Incubation and Pot Experiments

Different dosage (25 g, 50 g, and 75 g) amendments (zeolite, bentonite, AC) and chitosan-introduced composites (the application ratio of CS was 6% in chitosan-introduced composites. CS-zeolite, CS-bentonite, CS-AC) were thoroughly mixed with 5 kg of Cd-contaminated soil; subsequently, the amended soil and the control soil samples were packed into respective pots. A control treatment with no amendment was also prepared. All soil samples were incubated for one month.

Seeds of *Brassica juncea* were sowed in the middle of the greenhouse after mixing the unpolluted surface soils using a rotary machine. The cultivation area was covered with polyethylene plastic film to ensure adequate moisture. After three weeks, three equally sized seedlings of *Brassica juncea* (10 cm in height) were transplanted into each pot. Soil water was maintained at 60% WHC by adding tap water daily.

Sampling Pretreatment and Metal Analysis

On July 4, 2015, plant samples were harvested and rinsed with tap water to remove surface soil. The plants were then carefully washed with deionized water and dried of surface water. The fresh samples were weighed

and subsequently oven-dried at 105°C for 30 min, then at 70°C until constant weight. The dried samples were crushed and ground to a fine powder using a high-speed universal pulverizer (FW100, Zhongxing Weiye instrument co., LTD, China) and stored in plastic bags until analysis.

For analysis, 0.5 g of the dried plant sample was placed into a 100 mL beaker covered with a funnel. Then 15.0 mL of solution containing 87% of concentrated HNO₃ and 13% of concentrated HClO₄ (v/v) was added. The sample was digested at 50-80°C overnight and then at 180°C for 8 h to near dryness and allowed to cool. The digested solution was diluted with 2% (v/v) HNO₃ to a volume of 25.0 mL before filtration. The concentration of Cd in the digested plant sample was determined using a flame atomic absorption spectrometer at a wavelength of 228.8 nm [36]. Standard material, GBW-10015, was bought from a standard material center in Beijing, China to control analysis quality. The recovery rate was 95 \pm 75%.

Statistical Analysis

All analyses were performed using SigmaPlot 12.5(Systat Software Inc.), Excel 2000, and Origin 9.0 software. One-way ANOVA and multiple comparisons (Fisher LSD Method) were used to compare the difference in the dry weight of individual *Brassica juncea*, Cd uptake of *Brassica juncea*, and bioavailable Cd content; significance levels were 0.01 and 0.05. Data represent mean values of three replicates.

Results and Discussion

Effects of Different Amendments on Soil pH

Soil pH is an important factor of heavy metal adsorption as it affects the biological effectiveness of heavy metal ions in the soil. The pH values of different amendments and its chitosan composites are shown in Table 2. All of amendments were alkaline. Table 3 shows the pH value effect of the soil sample treated with different amendments. It is well known that biological effectiveness of heavy metal ions in soil decreased with increasing soil pH [37]. And the concentration of heavy metals in soils was influenced by soil pH [38]. However, some amendments showed nonspecific adsorption. Heavy metal adsorption varied with changing soil pH values. At low soil pH we observed competitive adsorption between heavy metals and a high concentration of H⁺ in the soil, impeding heavy metal adsorption [39]. The adsorbed metal ions increased

Table 2. pH values of different amendments.

Amendments	CS	Zeolite	Bentonite	AC	CS-Z	CS-B	CS-AC
pH	7.823	7.802	8.975	10.133	8.411	9.063	9.876

Table 3. pH value effects of the soil samples treated with different amendments.

Amendments	Added contents of different amendments(g/pot)			
	CK	25	50	75
Zeolite	7.786	7.883	7.874	7.872
Bentonite	7.786	7.955	7.898	7.874
AC	7.786	7.987	7.896	7.898
CS-Zeolite	7.786	7.793	7.925	7.912
CS-Bentonite	7.786	7.788	7.957	7.835
CS-AC	7.786	7.909	7.937	7.923

with increasing soil pH. Zeolite, bentonite, AC, and their chitosan composites showed an alkaline pH; their application significantly affected soil pH. This, in turn, can reduce the biological effectiveness of heavy metals to a certain extent. Besides, soil has intrinsic buffer properties and the pH therefore remained alkaline.

Effects of Different Amendments on the Dry Weight of *Brassica juncea*

The impacts of different amendments on the dry weight of individual *Brassica juncea* plants are shown in Table 4, along with the ANOVA results. The results showed that the addition of the amendments significantly increased the dry weight of individual *Brassica juncea* plants. However, there were no significant differences between the different pots in the same dosage. Plant dry weight increased with increasing amendment dosage (25 g, 50 g, and 75 g). The highest increase in dry weight was observed with the addition of CS-AC, followed by CS-bentonite, CS-zeolite, AC, bentonite, and zeolite. With an amendment dose of 75 g per pot, individual *Brassica juncea* dry weight increased by 41.91%, 39.00%, 27.64%, 35.93%, 23.78%, and 23.58% for CS-AC, CS-bentonite, CS-zeolite, AC, bentonite, and zeolite, respectively, compared with the control.

Cadmium contamination can significantly reduce soil pH, negatively impacting crop growth and development. The addition of different soil amendments

can improve soil pH and growth conditions, thereby decreasing soil Cd activity. Therefore, the adverse soil conditions to the growth and development of *Brassica juncea* can be alleviated, abating the negative impacts of excessive soil cadmium. This may explain the higher dry weight values for individual *Brassica juncea* in treatments with added amendments compare to control. The higher dry weights in chitosan composite-treated soils might be attributed to the introduced chitosan and increased soil pH [40]. One reason is that chitosan has a large amount of amino and hydroxyl groups with a high affinity for Cd ions, making it highly suitable for Cd removal [20, 41-43]. The *Brassica juncea* dry weight in chitosan composite-treated soil was higher than in soil with other amendments; these results are agreement with the findings of Shaheen et al. [40]. So the addition of different amendments had beneficial effects on the growth of *Brassica juncea* because all treatments with amendments had a higher dry weight than the control.

Effects of Different Amendments on Cd Uptake by *Brassica juncea*

Table 5 shows the effects of different amendments on Cd uptake by *Brassica juncea* and the ANOVA results. Cadmium uptake was significantly decreased, albeit to different degrees, with the addition of various amendments. Different dosages resulted in significant differences in terms of Cd uptake by *Brassica juncea*.

Table 4. *Brassica juncea* dry weight in soils treated with different amendments.

Amendment	Dry weight of individual <i>Breassica jincea</i> (g)			
	0 (g/pot)	25(g/pot)	50(g/pot)	75(g/pot)
Zeolite	8.058±0.41Ab	9.381±0.62Aa	9.669±0.57Aa	9.958±0.25Aa
Bentonite	8.058±0.41Ab	9.513±0.43Aa	9.894±0.88Aa	9.974±0.53Aa
AC	8.058±0.41Ab	9.611±0.58 Aab	10.545±0.45Aa	10.953±1.45Aa
CS-zeolite	8.058±0.41Ab	9.582±0.31Aa	9.913±0.54Aa	10.285±0.34Aa
CS-bentonite	8.058±0.41Ac	9.681±0.38Ab	10.331±0.38Ab	11.201±0.19Aa
CS-AC	8.058±0.41Ac	9.806±0.41Ab	11.367±0.79Aa	11.435±0.40Aa

Table 5. Cd content of plants in soil treated with different amendments.

Amendments	The Cd content (mg/kg)			
	0 (g/pot)	25(g/pot)	50 (g/pot)	75(g/pot)
Zeolite	4.98±0.15 Aa	4.45±0.06 Ab	4.31±0.09 Ab	4.29±0.04 Ab
Bentonite	4.98±0.15 Aa	4.33±0.07ABCb	4.24±0.11 ABb	4.11±0.11 ABb
AC	4.98±0.15 Aa	4.38±0.06 ABb	4.17±0.07 ABb	4.05±0.14 BCb
CS-Zeolite	4.98±0.15 Aa	4.38±0.06ABb	4.12±0.16 ABb	4.01±0.15 BCb
CS-Bentonite	4.98±0.15 Aa	4.28±0.09 BCb	4.07±0.06 BCb	3.99±0.15 BCb
CS-AC	4.98±0.15 Aa	4.23±0.10 Cb	3.91±0.13 Cb	3.89±0.10 Cb









Fig. 1. Regression curves between Cd uptake by *Brassica juncea* and added amount of different soil amendments.

The lower the content of Cd in *Brassica juncea*, the better the effect of the amendment-fixed Cd. The highest effects were found for CS-AC, followed by CS-bentonite, CS-zeolite, AC, bentonite, and zeolite, which means that the effect of the fixed Cd of the amendments was followed by CS-AC, CS-bentonite, CS-zeolite, AC, bentonite, and zeolite. The effect of dosage on Cd uptake was also studied for various amendment dosages (25 g, 50 g, and 75 g). With a dosage of 75 g per pot, the Cd uptake by *Brassica juncea* was lower than the other dosages. Cadmium uptake decreased by 21.89%, 19.88%, 19.48%, 18.67%, 17.47%, and 13.85%, respectively, for CS-AC, CS-bentonite, CS-zeolite, AC, bentonite, and zeolite. The results clearly show that remediation efficiency was better than the treatments without added chitosan, most likely due to the addition of chitosan with a large number of adsorption sites. Chitosan contain large numbers of chemical groups, such as amine and hydroxyl groups, and these groups have the ability to bind the Cd metals through several mechanisms, including chemical interactions such as chelation, electrostatic interactions, such as ion exchange, or the formation of ion pairs [44]. Besides, the effect of amendments on Cd uptake by *Brassica juncea* was CS>bentonite>zeolite. It may be that the AC itself has a larger surface area and stronger adsorption capacity than other two materials. The relative percentage of Cd desorbed from bentonite was higher than from zeolite. This means that bentonite has a stronger and more specific bonding of Cd compared to zeolite. These results may be attributed to the differences in the amendments' characteristics and structures. Although natural zeolite also has good adsorption performance, because the channels of zeolite are often blocked by some impurities and uneven arrangement these faults limit its adsorption effect. Bentonite contains a high proportion of swelling clays [45] and has the tendency to increase in volume (swell) [46] and provide a much higher surface area [28].

In order to understand the effects of different amendments on Cd uptake by *Brassica juncea*, regression equations and the corresponding correlation curves were fitted and are shown in Fig. 1 (amendment dosage as independent variable and Cd uptake as dependent variable).

The relationship between Cd uptake and the addition of different amendments followed a reverse "parabola"-type function (Fig. 1). Correlation coefficient ($R^2>0.92$) of samples treated with chitosan was higher than that in samples without chitosan. With an amendment dosage of 50-75 g per pot, Cd uptake was the correspondingly lowest. Hence 50-75 g/pot amendments can be used for preliminary dosage to repair Cd-polluted soil. To sum up, the dosage of various amendments treated with Cd-polluted soil is appropriate for 50-75 g/pot. The amendment can reduce by 14% the Cd uptake by *Brassica juncea* at least. In this case, CS-AC has the greatest potential for reducing Cd uptake by *Brassica juncea*.

Effects of Different Amendments on Bioavailable Cd Content

According to the ANOVA results (Table 6), the addition of different amendments significantly decreased bioavailable Cd content compared to the control. Differences were also found between the different amendment dosages. And the results show that the bioavailable Cd content has a different degree-reducing effect treated with different amendments. In other words, all six kinds of amendments on the heavy metal Cd have a certain fixed effect. With a dosage of 75 g CS-AC per pot, bioavailable Cd content was lowest and significantly decreased by 27.38% compared to the control. Based on the results of multiple comparisons, adding 25 g and 50 g of CS-AC per pot did not result in a significant difference in terms of bioavailable Cd content. With a dosage of 75 g, 50 g, 75 g, 75 g, 50 g, and 75 g per pot for CS-AC, CS-bentonite, CS-zeolite, AC, bentonite, and zeolite, respectively, bioavailable Cd content decreased by 27.38%, 19.29%, 22.83%, 23.22%, 15.74%, and 8.66%, respectively, compared to the control. Bioavailable Cd content was higher in treatments with the addition of chitosan compared to treatments without chitosan (Table 6), which has a large number of active sites and, consequently, a high affinity for Cd ions. On the other hand, activated carbon has a larger surface area and a strong adsorption capacity. The effectiveness of heavy metals was decreased through the following two effects: 1) The metals are directly adsorbed by activated carbon and 2) The addition

Table 6. Contents of bioavailable in soil treated with different amendments.

Amendments	The bioavailable Cd content (mg/kg)			
	0	25	50	75
Zeolite	2.54±0.07Aa	2.46±0.09 Aa	2.38±0.04Ab	2.32±0.10Ab
Bentonite	2.54±0.07Aa	2.21±0.05Bb	2.14±0.12Bb	2.16±0.14Ab
AC	2.54±0.07Aa	2.28±0.10Bb	2.11±0.07 Bc	1.95±0.10 Bc
CS-Zeolite	2.54±0.07Aa	2.15±0.09 Bb	2.19±0.11Bb	1.96±0.09Bc
CS-Bentonite	2.54±0.07Aa	2.13±0.09Bb	2.05±0.11Bb	2.08±0.09ABb
CS-AC	2.54±0.07Aa	2.03±0.11BCb	2.01±0.10BCbc	1.85±0.08BCc

of activated carbon can increase the content of soil organic carbon, thereby indirectly increasing heavy metal fixation [47]. Based on the above-described results, adding chitosan as a soil amendment results in a significant reduction of bioavailable Cd.

Conclusion

We investigated changes in soil pH, *Brassica juncea* dry weight, Cd content of *Brassica juncea*, and available soil Cd content with the addition of natural and chitosan-introduced zeolite, bentonite, and activated carbon. Our results show that natural and chitosan-introduced zeolite and bentonite can increase soil pH and reduce the biological effectiveness of heavy metals to a certain extent. Although activated carbon has an acidic pH value, the soil could buffer this effect and stay alkaline. With increasing amendment dosage, *Brassica juncea* dry weight increased. With an amendment dosage of 75 g per pot, Cd uptake by *Brassica juncea* was lowest. With a dosage of 75 g, 50 g, 75 g, 75 g, 50 g, and 75 g per pot for CS-AC, CS-bentonite, CS-zeolite, AC, bentonite, and zeolite, respectively, Cd uptake decreased by 21.89%, 19.88%, 19.48%, 18.67%, 17.47%, and 13.85%, respectively. Similarly, bioavailable Cd content decreased by 27.38%, 19.29%, 22.83%, 23.22%, 15.74%, and 8.66%, respectively, compared to the control.

The results of this study are of considerable significance in demonstrating the practical application of chitosan in the successful remediation of Cd-contaminated soil.

Acknowledgements

The authors gratefully acknowledge financial support by the Ministry of Agriculture of the People's Republic of China (No. 201203045), the Seed Foundation of Innovation and Creation for Graduate Students in Northwestern Polytechnic University (Z2017192), the National Natural Science Foundation of China (41502240), the Natural Science Basic Research Plan in Shaanxi Province of China (2017JM4005), the Fundamental Research Funds for the Central Universities (3102017zy056), and the Key Laboratory of Groundwater Contamination and Remediation, China Geological Survey (CGS) and Hebei Province (KF201610).

Conflict of Interest

The authors declare no conflict of interest.

References

1. BASZYŃSKI T. Interference of Cd²⁺ in functioning of the photosynthetic apparatus of higher plants. *Acta Societatis Botanicorum Poloniae*, **55** (2), 291, **1986**.
2. HE H., NORA F.Y.T., YAO A., QIU R., LI, W.C., YE Z. Growth and Cd uptake by rice (*Oryza sativa*) in acidic and Cd-contaminated paddy soils amended with steel slag. *Chemosphere*, **189**, 247, **2017**.
3. VAREDA J.P., VALENTE A.J.M., DURÃES L. Heavy metals in Iberian soils: Removal by current adsorbents/amendments and prospective for aerogels. *Advances in Colloid and Interface Science*, **237**, 28, **2016**.
4. MAHAR A., WANG P., ALI A., GUO Z.Y., AWASTHI M.K., LAHORI A.H., WANG Q., SHEN F., LI R.H., ZHANG Z.Q. Impact of CaO, fly ash, sulfur and Na₂S on the (im)mobilization and phytoavailability of Cd, Cu and Pb in contaminated soil. *Ecotoxicology and Environmental Safety*, **134**, 116, **2016**.
5. ALI A., GUE D., MAHAR A., WANG P., SHEN F., LI R.H., ZHANG Z. Mycoremediation of potentially toxic trace elements- a biological tool for soil cleanup: a review. *Pedosphere*, **27** (2), 205, **2017**.
6. TICA D., UDOVIC M., LESTAN D. Immobilization of potentially toxic metals using different soil amendments. *Chemosphere*, **85**(4) 577, **2011**.
7. PÉREZ-ESTEBAN J., ESCOLÁSTICO C., MOLINER A., MASAGUER A., RUIZ-FERNÁNDEZ J. Phytostabilization of metals in mine soils using *Brassica juncea* in combination with organic amendments. *Plant and Soil*, **377** (1-2), 97, **2014**.
8. HUSSAIN L.A., ZHANG Z., GUO Z., MAHAR A., LI R., KUMAR A.M., ALI S.T., KUMBHAR F., WANG P., SHEN F., ZHAO J., HUANG H. Potential use of lime combined with additives on (im)mobilization and phytoavailability of heavy metals from Pb/Zn smelter contaminated soils. *Ecotoxicology and Environmental Safety*, **145**, 313, **2017**.
9. RAFIQ M.T., AZIZ R., YANG X., XIAO W., RAFIQ M.K., ALI B., LI T. Cadmium phytoavailability to rice (*Oryza sativa* L.) grown in representative Chinese soils. A model to improve soil environmental quality guidelines for food safety. *Ecotoxicology and Environmental Safety*, **103** (1), 101, **2014**.
10. LI G., QIU J., YIN C. Study on calculating losses of cropland degradation. *Chinese Agricultural Science Bulletin*, **25** (3), 230, **2009**.
11. YAN H., LIN G. Usage of chitosan on the complexation of heavy metal contents and vertical distribution of Hg (II) and Cr (VI) in different textural artificially contaminated soils. *Environmental Earth Sciences*, **73** (5), 2483, **2015**.
12. GONZÁLEZ-NÚÑEZ R., ALBA M.D., ORTA M.M., VIDAL M., RIGOL A. Remediation of metal-contaminated soils with the addition of materials – Part I: Characterization and viability studies for the selection of non-hazardous waste materials and silicates. *Chemosphere*, **85** (9), 1511, **2011**.
13. SHAHEEN S.M., TSADILAS C.D., RINKLEBE J. 2015. Immobilization of soil copper using organic and inorganic amendments. *Journal of plant nutrition and soil science*, **178** (1), 112, **2015**.
14. GONZÁLEZ-NÚÑEZ R., ALBA M.D., ORTA M.M., VIDAL M., RIGOL A. Remediation of metal-contaminated soils with the addition of materials – Part II: Leaching tests to evaluate the efficiency of materials in the remediation of contaminated soils. *Chemosphere*, **87** (8), 829, **2012**.
15. HUANG R., YANG B., LIU Y. Simultaneous adsorption of aniline and Cr (VI) ion by activated carbon/chitosan composite. *Journal of applied polymer*, **131** (4), 1, **2014**.

16. CRINI G. Recent developments in polysaccharide-based materials used as adsorbents in wastewater treatment. *Progress of Polymer Science*, **30** (1), 38, **2005**.
17. TRIPATHI N., CHOPPALA G., SINGH R.S. Evaluation of modified chitosan for remediation of zinc contaminated soils. *Journal of Geochemical Exploration* <http://dx.doi.org/10.1016/j.gexplo.2016.08.011>, **2016**
18. ZHANG L., ZENG Y.X., CHENG Z.J. Removal of heavy metal ions using chitosan and modified chitosan: A review. *Journal of Molecular Liquids*, **214**, 175, **2016**.
19. MUZZARELLI R.A.A. *Industrial Production and Application*. Chitin, Pergamon Press, Oxford, England, **207**, **1977**.
20. LI M., ZHANG Z., LI R., WANG J., ALI A. Removal of Pb (II) and Cd (II) ions from aqueous solution by thiosemicarbazide modified chitosan. *International Journal of Biological Macromolecules*, **86**, 876, **2016**.
21. BORNET A., TEISSEGRE P.L. Chitosan, chitin-glucan and chitin effects on minerals (iron, lead, cadmium) and organic (ochratoxin A) contaminants in wines. *European Food Research and Technology*, **226** (4), 681, **2008**.
22. XU Y., LIANG X.F., QIN X., HUANG Q.Q., WANG L., SUN Y.B. Remediation of heavy metal-polluted agricultural soils using clay minerals: A review. *Pedosphere*, **27** (2), 193, **2017**.
23. SUN Y., WU Q.T., LEE C.C.C., LI B.Q., LONG X.X. Cadmium sorption characteristics of soil amendments and its relationship with the cadmium uptake by hyperaccumulator and normal plants in amended soils. *International Journal of Phytoremediation*, **16** (5), 486, **2014**.
24. FU F., WANG Q. Removal of heavy metals from wastewaters: a review. *Journal of Environmental Management*, **92**, 407, **2011**.
25. KUMARARAJA P., MANJIAH K.M., DATTA S.C., SARKAR B. Remediation of metal contaminated soil by aluminium pillared bentonite: Synthesis, characterisation, equilibrium study and plant growth experiment. *Applied Clay Science*, **137**, 115, **2017**.
26. LI P., AN Z., ZHAO T., LIU B., ZHANG C., ZHANG J., YANG M., ZHANG D., LI Y. Effects of natural zeolite addition on Cd in soil and tomato biomass. *Ecology and Environmental Sciences*, **20** (6-7), 1147, **2011**.
27. LIU X., ZHAO X., MA Z. Application of bentonite and zeolite in dealing soil contaminated by Cd. *Journal of Soil and Water Conservation*, **21** (6), 83, **2007**.
28. HAMIDPOUR M., KALBASI M., AFYUNI M., SHARIATMADARI H., HOLM P., HANSEN G. Sorption hysteresis of Cd (II) and Pb (II) on natural zeolite and bentonite. *Journal of Hazardous of Materials*, **181** (1-3), 686, **2010**.
29. SUN Y., LI Y., XU Y., LIANG X., WANG L. *In situ* stabilization remediation of cadmium (Cd) and lead (Pb) co-contaminated paddy soil using bentonite. *Applied Clay Science*, **105-106**, 200, **2015**.
30. LI J., LI Y., MENG Q. Removal of nitrate by zero-valent iron and pillared bentonite. *Journal of Hazardous Materials*, **174** (1-3), 188, **2010**.
31. BŘENDOVÁ K., ZEMANOVÁ V., PAVLÍKOVÁ D., TLUSTOŠ P. Utilization of biochar and activated carbon to reduce Cd, Pb and Zn phytoavailability and phytotoxicity for plants. *Journal of Environmental Management*, **181**, 637, **2016**.
32. EBBS S.D., LASAT M.M., BRADY D.J., CORNISH J., GORDON R., KOCHIAN L.V. Phytoextraction of cadmium and zinc from a contaminated soil. *Journal of Environmental Quality*, **26** (5), 1424, **1997**.
33. EBBS S.D., KOCHIAN L.V. Phytoextraction of Zinc by oat (*Avena sativa*), barley (*Hordeum vulgare*), and Indian mustard (*Brassica juncea*). *Environmental Science and Technology*, **32** (6), 802, **1998**.
34. CHEN H., ZHENG C., TU C., SHEN Z. Chemical ethods and phytoremediation on soil contaminated with heavy metals. *Chemosphere*, **41** (1-2), 229, **2000**.
35. TRIPATHI R.D., VAJPAYEE P., SINGH N., RAI U.N., KUMAR A., ALI M.B., KUMAR B., YUNUS M. Efficacy of various amendments for amelioration of fly-ash toxicity: growth performance and metal composition of *Cassia siamea* Lamk. *Chemosphere*, **54** (11), 1581, **2004**.
36. JI P., SUN T., SONG Y., ACKLAND M.L., LIU Y. Strategies for enhancing the phytoremediation of cadmium-contaminated agricultural soils by *Solanum nigrum* L. *Environmental Pollution*, **159** (3), 762, **2011**.
37. ZHANG Q., LI J., XU M., SONG Z., ZHOU S. Effect of amendments on bioavailability of cadmium and zinc in compound contaminated red soil. *Journal of Agro-Environment Science*, **25** (4), 861, **2006**.
38. BOLAN N.S., CHOPPALA G., KUNHIKRISHNAN A., PARK J., NAIDU R. Microbial transformation of trace elements in soils in relation to bioavailability and remediation. *Reviews of Environmental Contamination and Toxicology*, **225**, 1, **2013**.
39. YANG L., CHEN Z., LIU Y., WANG Y. Effects of lime and activated carbon on remedying chromium contaminated soil. *Acta Pedologica Sinica*, **49** (3), 518, **2012**.
40. SHAHEEN S., RINKLEBE J. Impact of emerging and low cost alternative amendments on the (im)mobilization and phytoavailability of Cd and Pb in a contaminated floodplain soil. *Ecological Engineering*, **74** (1), 319, **2015**.
41. GUO Z., HU X., AO Y. Effect of chitosan on the available contents and vertical distribution of Cu²⁺ and Cd²⁺ in different textural soils. *Journal of Hazardous Materials*, **167** (1-3), 1148, **2009**.
42. BORSAGLI F.G.L.M., MANSUR A.A.P., CHAGAS P., OLIVERIRA L.C.A., MANSUR H.S. O-carboxymethyl functionalization of chitosan: Complexation and adsorption of Cd (II) and Cr (VI) as heavy metal pollutant ions. *Reactive and Functional Polymers*, **97**, 37, **2015**.
43. ZENG L., CHEN Y., ZHANG Q., GUO X., PENG Y., XIAO H., CHEN X., LUO J. Adsorption of Cd (II), Cu(II) and Ni(II) ions by cross-linking chitosan/rectoritenano-hybrid composite microspheres. *Carbohydrate Polymers*, **130**, 333, **2015**.
44. EROSA M.S.D., EROSA T.I.S., MEDINA R.N., MENDOZA M.A., RODRIGUEZ, GUIBAL E. Cadmium sorption on chitosan sorbents: kinetic and equilibrium studies. *Hydrometallurgy*, **61** (3), 407, **2011**.
45. NGAH W.S.W., TEONG L.C., HANAFIAH M.A.K.M. Adsorption of dye and heavy metal ions by chitosan composite: A review. *Carbohydrate Polymers*, **83** (4), 1446, **2011**.
46. KAYA A., ÖREN A.H. Adsorption of zinc from aqueous solutions to bentonite. *Journal of Hazardous of Materials*, **125** (1-3), 183, **2005**.
47. HU Z., ZHANG G., WANG G., ZHAO Q., LIU X., CAO X., CAO Z. Effect of soil amendments on cadmium and lead contents in tobacco. *Acta Pedologica Sinica*, **43** (2), 233, **2006**.